


FOR THE PEOPLE
FOR EDUCATION
FOR SCIENCE

LIBRARY
OF
THE AMERICAN MUSEUM
OF
NATURAL HISTORY

Bound at
A.M.N.H.
1924



日本鳥學會編纂
日本鳥類目錄

59 x 25
**A HAND-LIST OF THE
JAPANESE BIRDS.**



創立第十週年記念出版

日本鳥學會

1922

A Hand-List of the Japanese Birds.

By

N. Kuroda, F. M. B. O. U., H. F. A. O. U. (Order Colymbi—Order Alectorides, pp. 1—59).

Viscount Y. Matsudaira (Order Limicolæ—Order Columbæ, pp. 59—91).

Prince N. Taka-Tsukasa (Order Cuculi—Order Pici, pp. 91—111).

S. Uchida, C. F. A. O. U. (Order Passeres, pp. 111—184).

(Issued for the Commemoration of the 10th Year of the Foundation of the
Ornithological Society of Japan).

1922.

The Ornithological Society of Japan.

c/o Zoological Institute Science College,
Tokyo Imp. Univ.

凡　例

(一) 本目録ハ日本鳥學會創立第十週年ヲ記念セシガ爲メ刊行セルモノニシテ本邦全土(滿洲及南洋諸島ハ除外セリ)ニ産スル既知鳥類ノ凡テヲ掲載セリ

(二) 本目録ハ下ノ部類ニ分チ下記四氏編輯ヲ擔當セラレタリ
1 阿比日ヨリ鶴型目迄九目

理學士 黒田長禮氏 (一ヨリ五九頁)

2 鶴型目ヨリ鳩鵠目迄五目

子爵 松平頼孝氏 (五九ヨリ九一頁)

3 杜鵑目ヨリ啄木鳥目迄十目

理學士 公爵 鷹司信輔氏 (九一ヨリ一一一頁)

4 燕雀目

獸醫學士 内田清之助氏 (一一一ヨリ一八四頁)

(三) 本目録ノ編纂ハ前記諸氏數回協議ノ上大體方針ヲ定メ各分擔部類ノ編纂ニ着手シ大正九年四月ヨリ大正十年八月ニ至ル十七ヶ月間ニ脱稿セリ

(四) 本目録ニハ目科屬種及亞種ヲ掲ゲ亞目亞科亞屬ハ之レヲ省略セリ

(五) 本目録ニハ學名和名及ビ之ガしのにむ並ニ英名及產地ヲ掲載セリ

(六) 學名ノ直下ニ記セシ書名ハ該學名ノ發表セラレタル書名頁數及基產地ヲ示ス

(七) 學名ノしのにむハ次ノ場合ニ限り之ヲ掲載セリ

A 内田清之助氏著「日本鳥類圖說」初版及再版ニ採用セル學名ニシテ本目録ノモノト相異セル場合

B Hartert: Die Vögel der paläarktischen Fauna ニ用ヒラレタル學名ニシテ本目録ノモノト相異セル場合

C 従來本邦ニテ廣く使用セラレ或ハ其他特ニ重要ナルしのにむト認メタルモノ

(八) 和名ハ主トシテ日本鳥類圖說ニ採用セラレタルモノヲ用ヒタルモ二三不適當ナルモノハ變更ヲ加ヘタリ但シ重要ナルモノ

ハ前者ノしのにむトシテ之ヲ掲ゲタリ從來和名ナキ種類ニハ凡テ
新ニ命名セリ

(九) 英名ハ主トシテ飯島博士編纂「日本ノ鳥目録」 Dresser,
A manual of Palaeartic Birds, Fauna of British India 等ニ採用ノモ
ノヲ用ヒ若シ之等ノ書物中ニナキ場合ニハ編輯者協議ノ上新ニ命
名セリ

(十) 產地記述ノ順序ハ大體樺太, 千鳥 北海道, 本道, 伊豆七島
小笠原群島, 四國, 九州, 對馬, 薩南諸島, 琉球, 臺灣, 朝鮮, 濟洲島,
鬱陵島等トシ分布局限セル種類ニアリテハ特ニ其地名ヲ記セリ

(十一) 本目録ノ編輯ニ當リ日本鳥學會々員糸山德太郎氏ハ諸
種ノ助力ヲ與ヘラレタリ記シテ之ヲ鳴謝ス

(十二) 從來發行セラレタル本邦產鳥類目錄下ノ如シ

1. Temminck & Schlegel: Siebold's Fauna Japonica. Aves, 1845-1850.
2. Blakiston & Pryer: A Catalogue of the Birds of Japan. Ibis, 1878, pp. 209-250.
3. Blakiston & Pryer: Birds of Japan. Trans. As. Soc. Japan, 1882, pp. 84-186.
4. Blakiston: Amended List of the Birds of Japan. 1884.
5. Seeböhm, H.: The Birds of the Japanese Empire. 1890.
6. Iijima, I.: Nippon no Torinokuroku (List of the Birds of Japan). "Dobutsugaku Zasshi" (Tokyo Zool. Mag.), Vol. III, 1891.
7. Ogawa, M.: A Hand-List of the Birds of Japan. Annot. Zool. Japan., Vol. VI, pp. 337-420, 1908.
8. Uchida, S.: Nihon san Chōrui Mokuroku (List of the Birds of Japan). "Nihon Chōrui Zusetsu" (The Birds of Japan), Vol. II, Suppl., pp. 1-36, 1914.

大正十一年九月

日本鳥學會

Order COLYBMI.

Family COLYMBIDÆ.

Colymbus L. (1758).

1. ***Colymbus stellatus*** Pontoppidan. *Albi.*
1763.—*Colymbus stellatus* Pontopp., Danske Atlas, I, p. 621
(Denmark).
Colymbus septentrionalis L.
Red-throated Diver.
Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku,
Kiusiu, Formosa, Corea, Quelpart Is.
2. ***Colymbus arcticus viridigularis*** (Dwight). *Ōhamu.*
1918.—*Gavia viridigularis* Dwight, Auk, p. 189 (North-east
Siberia).
Colymbus arcticus (nec L.), Uchida.
Siberian Black-throated Diver.
Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku,
Kiusiu, Tsushima, Corea, Quelpart Is.
3. ***Colymbus arcticus pacificus*** Lawrence *Shirocri-ōhamu*
1860.—*Columbus pacificus* Lawrence, Baird's B. N. Amer., p.
889 (California).
Pacific Diver.
Hab. Hokkaido, Hondo (Sagami, Suruga), Shikoku (Sanuki).
4. ***Colymbus adamsii*** Gray. *Hashijiro-abi.*
1859.—*Colymbus adamsii* Gray, Proc. Zool. Soc. London, p. 167.
(Alaska).
White-billed Diver.
Hab. Northern Kurile Is., Hokkaido (Hakodate), Hondo (Sen-
dai Bay, Sagami), Kiusiu, Corea.

Family PODICIPIDÆ.

Podiceps Latham (1787).

5. **Podiceps ruficollis japonicus** Hartert. *Kaitsuburi.*
1920.—*Podiceps ruficollis japonicus* Hartert, Vög. pal., II, p.
1455 (Tokio).
Podicipes fluviatilis philippensis (nec Bonnat.), Uchida.
Japanese Little Grebe.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Amami-oshima,
Corea, Quelpart Is.
6. **Podiceps ruficollis philippensis** (Bonnaterre).
..... *Taiwan-kaitsuburi.*
1790.—*Colymbus Philippensis* Bonnat., Tabl. Enc. et Méth. I,
p. 58. (Philippine Is.).
Philippine Little Grebe.
Hab. Formosa.

Dytes Kaup (1829).

7. **Dytes auritus** (L.). *Mimi-kaitsuburi.*
1758.—*Colymbus auritus* L., Syst. Nat., ed. X, I, p. 135 (Sweden).
Podiceps auritus (L.).
Slavonian Grebe.
Hab. Hokkaido, Hondo, Kiusiu, Tsushima, Corea.

Otodytes Reichenbach (1852).

8. **Otodytes nigricollis nigricollis** (Brehm). ... *Hajiro-kaitsuburi.*
1831.—*Podiceps nigricollis* Brehm, Handb. Naturg. Vög.
Deutschl., p. 963 (Germany).
Podicipes nigricollis Brehm.
Black-necked Grebe.
Hab. Sakhalien?, Hokkaido, Hondo, Shikoku, Kiusiu, Corea,
Quelpart Is.

Pedetaithya Kaup (1829).

9. **Pedetaithya griseigena holboellii** Reinhardt.
..... *Aka-ri-kaitsuburi.*

1853.—*Podiceps Holboellii* Reinh., Videnskab. Meddelelser, p. 76 (Greenland).

Podicipes holboelli (Reinhardt).

Eastern Red-necked Grebe.

Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, Corea.

Lophaithyia Kaup (1829).

10. **Lophaithyia cristatus cristatus** (L.) *Kumatori-kaitsuburi.*

1758.—*Colymbus cristatus* L., Syst. Nat., ed. X. I, p. 135, (Sweden).

Podicipes cristatus L.

Great Crested Grebe.

Hab. Hokkaido ?, Hondo (Awa), Tsushima ?, Kiusiu (Satsuma), Riu-kiu (Okinawajima), Formosa, Corea, Quelpart Is.

Order TUBINARES.

Family HYDROBATIDÆ.

Oceanodroma Reichenbach (1852).

11. **Oceanodroma leucorrhœa leucorrhœa** Vieillot.
..... *Koshijiro-u-mitsubame.*

1817.—*Procellaria leucorrhœa* Vieillot, N. Dict. d'Hist. Nat., XXV, p. 422 (North Atlantic).

Oceanodroma leachii (Temminck).

Leach's Petrel.

Hab. Kurile Is., Hokkaido.

12. **Oceanodroma castro** (Harcourt) *Kuro-koshijiro-u-mitsubame.*

1851.—*Thalassidroma castro* Harc., Sketch of Madeira, p. 123 (Madeira).

- Madeiran Fork-tailed Petrel.
- Hab. Hondo (Nikkō).
- 13. Oceanodroma markhami owstoni** (Mathews & Iredale).
 *Owston-umitsubame*.
- 1890.—? *Oceanodroma tristrani* Salvin, Cat. B. Brit. Mus., XXV, p. 354 (Sendai Bay).
- 1915.—*Cymochorza owstoni* Math. & Ired., Ibis, p. 581 (Okinose, Sagami).
- Oceanodroma melanaria* (nec Bp.), Uchida.
- Oceanodroma markhami* (nec Salvin), Uchida.
- Oceanodroma fuliginosa* (nec Gmelin), Uchida.
- Owston's Fork-tailed Petrel.
- Hab. Hondo (Sendai Bay, Uraga Channel, Sagami Bay), Torishima.
- 14. Oceanodroma melania** (Bonaparte). *Kuro-umitsubame*.
- 1854.—*Procellaria melania* Bp., C. R., XXXVIII, p. 662 (Nicaragua to California).
- Black Fork-tailed Petrel.
- Hab. Off the coast of Sagami. (Five specimens were obtained in May, 1921).
- 15. Oceanodroma monorhis monorhis** (Swinhoe).
 *Hime-kuroumitsubame*.
- 1867.—*Thalassidroma monorhis* Swinh., Ibis, p. 386 (Amoy, China).
- Oceanodroma monorhis* (Swinhoe).
- Swinhoe's Fork-tailed Petrel.
- Hab. Hondo (Mutsu), Riu-Kiu, North-east Formosa ?.
- 16. Oceanodroma furcata** (Gmelin). *Haïiro-umitsubame*.
- 1788.—*Procellaria furcata* Gm., Syst. Nat. I, 2, p. 561. ("Habitat in glacie maris, Americam et Asiam interfluentis").

Grey Fork-tailed Petrel.

Hab. Kurile Is., Hokkaido, Hondo (Sagami, Suruga, Kobe).

Family PUFFINID.E.

Puffinus Prisson (1760).

17. **Puffinus leucomelas** (Temminck). *O-nizunagidori.*

1836.—*Procellaria leucomelas* Temm., Pl. Col. 587, livr. 09 c.
("Les mers du Japon").

1915.—*Calonectris leucomelas* (Temm.). Math. & Ired., Ibis, p.
594.

Streaked Shearwater.

Hab. Kurile Is., Hokkaido (Hakodate), Hondo (Yokohama,
Daishi, Suruga, Saikawa, Kōyasan), Seven Is. (Mikura-
jima), Shikoku, Hoyo Channel, Kiusiu (Chikuzen, S.
Kiusiu), South Is. of Satsuma (Iwogashima, Tanegashima,
Yerabujima), Riu-kiu (Iriomotejima); Formosa, Corea,
Dagelet Is.=Utsurioto.

18. **Puffinus cuneatus** Salvin. *Onaga-mizunagidori.*

1888.—*Puffinus cuneatus* Salvin, Ibis, p. 353 (Krusenstern
Island, Pacific Ocean).

Salvin's Shearwater.

Hab. Hondo (Sagami Bay), Bonin Is., Sulphur Is., Marcus Is.,
Formosa (Pescadores Is.).

19. **Puffinus tenuirostris** tenuirostris (Temminck).
..... *Hashiboso-mizunagidori.*

1835.—*Procellaria tenuirostris* Temm., Pl. Col. text to Pl. 587
(Japanese Seas).

Puffinus tenuirostris (Temminck).

Slender-billed Shearwater.

Hab. Sakhalien, Kurile Is., Tsugaru Strait, Hondo (Tokyo Bay,

- Owari, Yoshino, Yamato), Bonin Is., Kiusiu (Nagasaki), Corea.
20. **Puffinus nativitatis** Streets. *Mizunagidori*.
 1877.—*Puffinus (Neotris) nativitatis* Streets, Bull. U. S. Nat. Mus., No. 7, p. 29 (Christmas Is.).
 Christmas Island Shearwater.
 Hab. Bonin Is., Marcus Is.
21. **Puffinus griseus** (Gmelin). *Haiire-mizunagidori*.
 1788.—*Procellaria grisca* Gmelin, Syst. Nat., I, 2, p. 564
 ("Habitat in Hemisphaerio australi, inter 35° et 50°").
 1915.—*Neonetris griseus pescadoresi* Math. & Ired., Ibis, p. 652 (Pescadores Is., Formosa).
 1915.—*Neonetris griseus missus* Math. & Ired., Ibis, p. 603.
 (Kurile Is.).
 Sooty Shearwater.
 Hab. Kurile Is., Hondo (Chōshi, Sagami Bay), Formosa
 (Pescadores Is.).
22. **Puffinus carneipes** Gould. *Alkaashi-mizunagidori*.
 1844.—*Puffinus carneipes* Gould, Ann. & Nat. Hist. XIII, p. 365 (The islands off Cape Leeuwin, W.-Australia).
 1912.—*Puffinus carneipes hakodate* Math., B. Austr., II, p. 90
 (Hakodate).
 Pink-footed Shearwater.
 Hab. Hokkaido (Hakodate), Hondo (Sagami Bay).
23. **Puffinus assimilis bannermani** Mathews & Iredale.
 *Ogasawara-mizunagidori*.
 1915.—*Puffinus bannermani* Math. & Ired., Ibis, p. 594 (Bonin Is.).
 Bannerman's Shearwater.
 Hab. Bonin Islands, Sulphur Islands.

Pterodroma Bonaparte (1856).

24. **Pterodroma brevipes** (Peale). *Kō-shirohara-nizunagidori.*
1848.—*Procellaria brevipes* Peale, U. S. Expl. Exp., VIII, pp.
294, 337, Pl. 80 (“Pacific Ocean 68° S., Long. 95° W.”).
Estrelata desolata (nec Gm.), Uchida.
White-throated Fulmar.
Hab. Hondo (? Mutsu, ? Hida).
Remark.—“Godmans und andere Angaben vom Vorkommen
in Japan beruhen auf Missverständnis” (Hartert).
25. **Pterodroma hypoleuca** (Salvin). *Shio-oharamizuna-židori.*
1888.—*Estrelata hypoleuca* Salvin, Ibis, p. 359 (Krusenstern
Island).
Bonin Island Fulmar.
Hab. Hondo (Sagami, Mino), Seven Is. (Miyakejima), Bonin-
Islands, Riu-Kiu.
26. **Pterodroma longirostris** (Stejneger).
..... *Hime-shiroharamizuna-židori.*
1893.—*Estrelata longirostris* Stejn., Proc. U. S. Nat. Mus.,
XVI, p. 618 (Mutsu, Hondo).
Estrelata leucoptera (nec Gould), Stejn.
Estrelata longirostris Stejn.
Japanese Fulmar.
Hab. Hondo (Mutsu).

Bulweria Bonaparte (1842).

27. **Bulweria bulwerii bulwerii** (Jardine & Selby). *Anadori.*
1828.—*Procellaria Bulwerii* Jard. & Selby, Ill. Orn., II, Pl.
65 and text (“Madeira or the small islands adjacent”).
1915.—*Bulweria bulweri pacifica* Math. & Ired., Ibis, p. 607
(Sulphur Is.).

Bulweria bulweri (Jard. & Selby).

Bulweria macgillivrayi (nec Gray), Kuroda.

Bulwer's Fulmar.

Hab. Hondo (Nikko, Ōsaka Bay), Bonin Is., Sulphur Is., Riū-kiū (Iriomotejima?), Formosa (Kelung, Menkato, Horisha).

Fulmarus Stephens (1820).

28. **Fulmarus glacialis rodgersii** Cassin. *Furuma-kamome*.

1862.—*Fulmarus Rodversii* Cassin, Proc. Acad., Philadelphia, p. 325 ("South Indian Ocean." Error). Type from North Pacific Ocean or Ice Sea).

1884.—*Fulmarus glacialis glupischa* Stejneger, Auk, I, p. 234 (Commander Islands).

Fulmarus glacialis (nec L.), Uchida.

Pacific Fulmar.

Hab. Sakhalien (Robben Is.), Kurile Is., Hokkaido, Hondo (Kawagoye, Shinano, Yechigo).

Family DIOMEDEIDÆ.

Diomedea L. (1758).

29. **Diomedea albatrus** Pallas. *Ahōdori*.

1769. *Diomedea albatrus* Pallas, Spic. Zool., V, p. 28 (Kamchatka & Kuril).

1829 & 1835.—*Diomedea brachiora* Temm., in text to livr. 75 and Pl. Col. Pl. 554 (Japanese Seas and near Riukiu Is.).

1873.—*Diomedea derogata* Swinhoe, Proc. Zool. Soc. London, p. 786 (East coast of China—young.)

Steller's Albatross.

Hab. Kurile Is. (Iturup), Hokkaido (Hakodate), Hondo (Tokyo

bay, Sagami), Seven Is. (Torishima), Bonin Is., Kiushu (Nagasaki, Kagoshima), Tsushima, Rukiu Is., Formosa (Formosan Channel, Agincourt Is., Pescadores Is.), Corean Channel.

30. **Diomedea immutabilis** Rothschild. *Kōakidōri*.
1893.—*Diomedea immutabilis* Rothschild. Bull. Brit. Orn. Club, Vol. I, p. xlvi (Laysan Is.).

Laysan Albatross.

Hab. Hondo (Sagami Bay), Marcus Island.

31. **Diomedea nigripes** Audubon. *Kuroishi shōdōri*.
1839.—*Diomedea nigripes* Audubon, Orn. Biogr., V, p. 327
("Pacific Ocean, in lat. $30^{\circ} 44'$ N., long. $140^{\circ} 1'$ ").

Black-footed Albatross.

Hab. Kurile Is., Tsugaru Strait (erroneously recorded as Hakodate), Hondo (Tokyo City, Ōiso, Odawara), Marcus Island, Formosa (Formosan Channel, Pescadores Is.).

Order STEGANOPODES.

Family PHAETHONTIDÆ.

Scæophaëthon Mathews (1913).

32. **Scæophaëthon rubricauda brevirostris** Mathews.
..... *Akao-nettaichō*.
1914-15.—*Scæophaëthon rubricauda brevirostris* Math., B. Austr., IV, p. 303 (Bonin Islands).
Phaëthon rubricauda (nec Bodd.), Uchida.
Bonin Island Red-tailed Tropic Bird.

Hab. Hondo (Rikuchū, Musashi, Mino), Bonin Is., Sulphur Is., Marcus Island.

Leptophæthon Mathews (1913).

33. **Leptophæthon lepturus dorotheæ** Mathews *Shirao-nettaichō*.
1913.—*Leptophæthon lepturus dorotheæ* Math., Ibis, p. 554.
(E. Australian Seas and Pacific Ocean).

Phaethon candidus (nec Temminck), Uchida.

White-tailed Tropic Bird.

Hab. Hondo (Shinano, Kaga), Bonin Is., Sulphur Is., Marcus Is.
Formosa.

Family SULIDÆ.

Sula Brisson (1760).

34. **Sula leucogaster plotus** (Forster). *Katsuwodori*.
1844.—*Pelicanus plotus* Forster, Descr. Anim., ed. Licht., p.
278. (Near New Caledonia)

Sula leucogastra (nec Bodd.), Uchida.

Sula sula (nec L.), Uchida.

New Caledonian Brown Gannet.

Hab. Bonin Is., Sulphur Is., Marcus Islands, Kiusiu (Nagasaki, Riu-Kiu Is. (Iriomote), Formosa.

Piscatrix Reichenbach (1852).

35. **Piscatrix sula rubripes** (Gould). *Akuashi-katsuwodori*.
1838.—*Sula rubripes* Gould, Synops. B. Austr., pt. IV, App.,
p. 7 (Raine Island, N. Queensland).

Sula piscator (nec L.), Uchida.

Australian Red-legged Gannet.

Hab. Marcus Island.

Parasula Mathews (1913).

36. **Parasula dactylatra personata** (Gould). *Iwotsura-katsutcodori.*

1846.—*Sula personata* Gould, Proc. Zool. Soc. London, p. 21
(Raine Island, N. Queensland).

Sula cyanops (nec Sundevall), Uchida.

Australian Masked Gannet.

Hab. Marcus Island, Riu-Kiu Islands.

Family PHALACROCORACIDÆ.

Phalacrocorax Brisson (1760).

37. **Phalacrocorax carbo** [(?) *sinensis* (Shaw & Nodder)]. *Umi-u.*

1801.—*Pelecanus sinensis* Shaw & Nodd., Nat. Misc. XIII, Pl. 529 (China).

Phalacrocorax carbo (nec L.), Uchida.

Chinese Cormorant.

Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima,
Formosa, Corea, Quelpart Is.

38. **Phalacrocorax capillatus** (Temminck & Schlegel). ... *Kawa-u.*

1850.—*Carbo capillatus* Temm. & Schl., Siebold's Fauna
Japonica, Aves, Pl. 83, 83 B.

1850.—*Carbo filamentosus*, iid., op. cit., p. 129 (Japan).

Temminck's Cormorant.

Hab. Hokkaido, Hondo, Kiusiu, Tsushima, Dagelet Is. = Utsu-
riōtō, Quelpart Is., Corea, South Is. of Satsuma (Tanega-
shima, Amami-ōshima).

Urile Bonaparte (1855).

39. **Urile urile** (Gmelin). *Chishima-u-garasu.*

1789.—*Pelecanus Urile* Gm., Syst. Nat., I, 2, p. 575. (Kamt-
schatka).

1827.—*Phalacrocorax bicristatus* Pallas, Zoogr. Rosso-Asiat., II, p. 103 (Kamtschatka, Kurile Is. and Islands till America).

Bare-faced Shag.

Hab. Kurile Is., Hondo (Echigo).

40. **Urile pelagicus pelagicus** (Pallas). *Ugarasu, Hime-ut.*
1827. *Phalacrocorax pelagicus* Pallas, Zoogr. Rosso-Asiat., II, p. 503 (Eastern Kamtschatka & Aleutian Islands).

Pelagic Shag.

Hab. Kurile Is., Hokkaido, Hondo, Kiushiu, Corea, Quelpart Is., Northern Formosa.

Family FREGATIDÆ.

Fregata Lacepédë (1799).

41. **Fregata aquila palmerstoni** (Gmelin). *S-gunkandori.*
1788.—*Pelecanus palmerstoni* Gm., Syst. Nat., I, 2, p. 573.
(Palmerstone Is., Pacific Ocean).

Fregata aquila (nec L.). Uchida.

Pacific Fregate Bird.

Hab. Sulphur Is., Marcus Island.

42. **Fregata ariel** [(?) *ariel* (Gould)]. *Gunkandori.*
1845. *Alagen ariel* Gray, Genera Brd., Vol. III, p. 669, Pl.
183 (Raine Island, N. Queensland).

Fregata ariel (Gould).

Lesser Fregate Bird.

Hab. Hokkaido (Hakodate), ? Formosa.

Family PELECANIDÆ.

Pelecanus L. (1758).

43. **Pelecanus crispus** Bruch. *Garancho.*

1832.—*Pelecanus crispus* Bruch, Isis, p. 1109 (Dalmatia, Austria).

Pelecanus sp., Uchida.

Dalmatian Pelican.

Hab. Kiusiu (Satsuma), Formosa, Corea (Keiki Distr.).

Order GRESSORES.

Family ARDEIDÆ.

Ardea L. (1758).

44. *Ardea cinerea* jouyi Clark. *Ake-sagi*.

1907.—*Ardea cinerea* *jouyi* Clark, Proc. U. S. Nat. Mus., XXXII, p. 408 (Seoul, Korea).

Ardea cinerea (nec L.), Uchida.

Eastern Grey Heron.

Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, South Is. of Satsuma (Amami-ōshima), Riukiu, ? Dagelet Is., Quelpart Is., Corea, Formosa.

Pyrrherodias Finsch & Hartlaub (1870).

45. *Pyrrherodias purpureus* manillensis (Meyen). *Murasaki-sagi*.

1831.—*Ardea purpurea* var. *manillensis* Meyen, Acta Acad. Leop. Carol., XVI, suppl., p. 102; id., Reise um die Erde, III, p. 226, 1831 (Philippine Is.).

Phœx manillensis (Meyen).

Eastern Purple Heron.

Hab. Hondo (Kanagawa, Shizuoka), Riu-Kiu (Ishigakijima), Formosa, Corea.

Egretta Forster (1817).

46. *Egretta garzetta* garzetta (L.) *Kosagi*.

1766.—*Ardea Garzetta* L., Syst. Nat., ed. XII, I, p. 237
("Habitat in Oriente").

Herodias garzetta (L.).

Little Egret.

Hab. Hokkaido, Hondo, Kiusiu, Riukiu (Okinawa, Miyako, Iriomote), Formosa.

Mesophoyx Sharpe (1894).

47. **Mesophoyx intermedia intermedia** (Wagler). *Chū-sagi*.

1829.—*Ardea intermedia* Wagler, Isis, p. 659 (Java).

Egretta intermedia intermedia (Wagler), Hartert.

Herodias intermedia (Wagler).

Plumed Egret.

Hab. Hokkaido, Hondo (Tokyo, Haneda, Yokohama, Suruga), Seven Is. (? Ōshima), Kiusiu (Nagasaki), Corea, Formosa.

Casmerodus Gloger (1842).

48. **Casmerodus albus albus** (L.). *Dai-sagi*.

1758.—*Ardea alba* L., Syst. Nat., ed. X, I, p. 144 ("Habitat in Europa").

Herodias timoriensis (Cuvier), Uchida (pt.).

Great White Egret.

Hab. Hokkaido, Hondo (Tokyo, Yokohama), Riukiu, Corea.

49. **Casmerodus albus modesta** (Gray). *Kōmomojiro*.

1831.—*Ardea modesta* Gray, Zool. Misc., p. 19 (India).

Ardea Torra Franklin.

Ardea timoriensis Lesson.

Egretta alba modesta (Gray), Hartert.

Herodias timoriensis (Cuvier), Uchida (pt.).

Herodias terra Gigl. & Salv., Uchida.

Eastern Great White Egret.

Hab. Kurile Is. (Iturup), Hokkaido (Hakodate), Hondo (Saitama, Chiba, Tokyo), ? Shikoku, Kiusin (Chikuzen, Nagasaki), Riu-Kiu (Okinawa), Formosa, Corea.

Hemigarzetta Mathews (1914).

50. **Hemigarzetta euphotes** Swinhoe. *Kara-shirasa_{gi}*.

1860.—*Herodias euphotes* Swinhoe, Ibis, p. 64 (Amoy, China).

Egretta euphotes (Swinh.), Hartert.

Demiegretta euphotes (Sw.).

Chinese Egret.

Hab. ? Tsushima, Quelpart Is., Corea, Formosa.

Demigretta (corr. **Demiegretta**) Blyth (1840).

51. **Demiegretta sacra ringeri** Stejneger.

..... *Kurosagi*, *Shirekuro-sagi*.

1887.—*Demiegretta ringeri* Stejn., Proc. U. S. Nat. Mus., X, p. 300 (Tsushima, Gotō, Riu-Kiu Is.).

Demiegretta jugularis (nec Wagler), Uchida.

Demiegretta jugularis greyi (nec Gray), Uchida.

Demiegretta sacra (nec Gm), Hartert (pt.).

Japanese Reef-Heron.

Hab. Hondo (Shimōsa, Sagami, Izu), Shikoku, Kiusiu (Chikuzen), Tsushima, Gotō Is., South Is. of Satsuma (Tanegashima, Amamīōshima), Riu-Kiu (Ishigaki, Iriomote), Formosa (near Kōshun, Botol-tobago, Pescadores Is.), Corea (N. Chūsei Distr., Chōsen Bay), Quelpart Is.

Bubuleus Bonaparte (1855).

52. **Bubuleus ibis coromandus** (Boddart). *Amasagi, Shōjōsagi.*

1783.—*Caverota Coromanda* Bodd., Tabl. Pl. Enl., p. 54
("Crabier de Coromandel").

Bubuleus coromandus (Boddart).

Indian Cattle Egret.

Hab. Hondo (Saitama, Tokyo, Haneda, Sakai near Ōsaka),
Kiusiu (Nagasaki). Riu-Kiu (Ishigaki, Iriomote), Tsushima,
Formosa, Corea.

Ardeola Boie (1822).

53. **Ardeola bacchus** (Bonaparte). *Akagashira-sagi.*

1855.—*Euphus bacchus* Bonaparte, Conspl. Gen. Av., II, p. 127
(Malacea).

Ardeola prasinocincta Swinhoe.

Chinese Pond Heron.

Hab. Hokkaido (Hakodate), Hondo (Kawanakajima in Shinano,
Kobe), Riu-Kiu (Miyakojima), Formosa (Taihoku).

Butorides Bluth (1852).

54. **Butorides striatus javanicus** (Horsfield). *Hime-sasagoi.*

1821.—*Ardea javanica* Horsf., Trans. Linn. Soc. London,
XII, p. 190 (Java).

Javan Green Heron.

Hab. Riu-Kiu (Okinawa), Formosa.

55. **Butorides striatus amurensis** Schrenck. *Sasagoi.*

1860.—*Ardea (Butorides) virescens* var. *amurensis* Schrenck,
Reis. Amur-Lande, I, 2, p. 441 (Amurland).

Butorides amurensis (Schlenk) errore, Uchida.

Amur Green Heron.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, South Is. of Satsuma
(Yakushima, Amamiōshima), Bonin Is., Corea, Formosa.

Nycticorax Forster (1817).

56. **Nycticorax nycticorax nycticorax** (L.)..... *Goi-sagi*.
1758.—*Ardea Nycticorax* L., Syst. Nat., ed. X, 1, p. 142
("Habitat in Europa australi").
Nycticorax nycticorax (L.).
Night-Heron.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, South Is. of Satsuma
(Amamiōshima), Riu-Kiu, Formosa, Quelpart Is., Dagelet
Is. (=Utsurio-tō).
57. **Nycticorax caledonicus crassirostris** Vigors.... *Hashibuto-goi*.
1839.—*Nycticorax crassirostris* Vigors, Voy. Blossom, Zool.,
p. 27 (Bonin Is.).
Bonin Island Night-Heron.
Hab. Bonin Is.

Gorsachius Bonaparte (1855).

58. **Gorsachius goisagi** (Temminck)..... *Mizo-goi*.
1835.—*Nycticorax Goisagi* Temm., Pl. Col., 582 (Japan).
Japanese Night-Heron.
Hab. Hondo, Seven Is. (Ōshima, Miyakejima), Shikoku, Kiusiu,
Riu-Kiu (Okinawa), Formosa.
59. **Gorsachius melanolophus** (Raffles)..... *Taiwan mizo-goi*.
1822.—*Ardea melanlopha* Raffl., Trans. Linn. Soc., XIII, p.
326 (Sumatra).
Butio melanlopha (Raffl.).
Malay Bittern.
Hab. Riu-Kiu (Ishigaki), Formosa.

Ixobrychus Billberg (1828).

60. **Ixobrychus sinensis sinensis** (Gmelin). *Yoshi-gvi.*

1789.—*Ardea Sinensis* Gm., Syst. Nat., I, 2, p. 642 (“Habitat in Sina”).

1887.—*Ardea Inteola* Stejneger, Proc. U. S. Nat. Mus., X, pp. 289—290 (Yezo and Hondo).

Ardetta sinensis (Gm.).

Chinese Little Bittern.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, South Is. of Satsuma (Amauiōshima), Formosa, Corea.

61. **Ixobrychus cinnamomeus** (Gmelin). *Riukiu-yoshigoi.*

1789.—*Ardea cinnamomea* Gm., Syst. Nat., I, 2, p. 643 (“Habitat in Sina”).

1905.—*Nannocnus ijimai* Ogawa, Annot., Zool. Japan., V, p. 215 (Yagachi, Okinawa, Ishigaki).

Cinnamon Bittern.

Hab. Riu-Kiu (Yagachi, Okinawa, Ishigaki), Formosa.

Nannocnus Stejneger (1887).

62. **Nannocnus eurythmus** (Swinhoe). *Ō-yoshigoi.*

1873.—*Ardetta erythra* Swinhoe, Ibis, p. 74, pl. 2 (Amoy and Shanghai).

Ixobrychus erythrus (Swinh.), Hartert.

Schrenck's Little Bittern.

Hab. Hokkaido, Hondo, Riukiu (Ishigaki), Formosa, Corea.

Botaurus Stephens (1819).

63. **Botaurus stellaris stellaris** (L.). *Sankano-gvi.*

1758.—*Ardea stellaris* L., Syst. Nat., ed. X, I, p. 144 (Sweden).

Botaurus stellaris (L.).

Bittern.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

Dupetor Heine & Reichenbach (1890).

64. **Dupetor flavigollis flavigollis** (Latham). ... *Takasago-kurosa*.

1790.—*Ardea flavigollis* Lath., Index Ornith., II, p. 701.

Dupetor flavigollis (Latham).

Ixobrychus (Dupetor) flavigollis flavigollis (Lath.), Hartert.

Black Bittern.

Hab. Formosa.

Family CICONIHDÆ.

Ciconia Brisson (1760).

65. **Ciconia ciconia boyaciana** Swinhoe. *Kōnotori*.

1873.—*Ciconia boyaciana* Swinhoe, Proc. Zool. Soc. London, p. 513 (Yokohama).

Japanese Stork.

Hab. Hondo (Shimōsa, Yokohama, Izushi and Kinosaki in Tajima), Shikoku (Tokushima), South Is. of Satsuma (Tanegashima), Formosa (Tainan), Corea, Tsushima.

Melanopelargus Reichenbach (1853).

66. **Melanopelargus niger** (L.) *Nabe-kō*.

1758.—*Ardea nigra* L., Syst. Nat., ed. X, I, p. 142 (North Europe).

Ciconia nigra (L.).

Black Stork.

Hab. Hondo (Tokyo, Nishitama in Musashi), Corea, Formosa.

Family IBIDIDÆ.

Nipponia Reichenbach (1852).

67. **Nipponia nippon** (Temminck). *Toki*.

1835.—*Ibis Nippon* Temm., Pl. Col., 551 (Japan).

Nipponia Temminckii Reichenb. (New name for *Ibis nippon*).

Ibis sincensis David (Grey phase!).

Japanese Crested Ibis.

Hab. Hokkaido (Hakodate), Hondo (Miyako in Rikuchū, Nishitama in Musashi, Yokohama, Mino, Echigo), Seven Is., Shikoku (Tokushima), Kiusu, Riu-Kiu (Okinawa), Formosa, Corea.

Threskiornis Gray (1842).

68. **Threskiornis melanocephalus** (Latham). *Kurotoki*.

1790.—*Tantalus melanocephalus* Lath., Index Ornith., II, p. 709 (India).

Ibis melanoccephala (Latham).

White Ibis.

Hab. Hokkaido, Hondo (Sagiyama at Saitama, Kameido in Tokyo, Irumagawa, Shinhamra in Shimōsa, Yokohama), Riukiu (Okinawa), Formosa.

Platalea L. (1758).

69. **Platalea leucorodia major** Temminck & Schlegel.... *Herasagi*.

1849.—*Platalea major* Temm. & Schl., Siebold's Fauna Japonica, Aves, p. 119, pl. 75 (Japan).

Platalea leucorodia (nec L.), Uchida.

Japanese Spoonbill.

Hab. Hokkaido, Hondo (Shimōsa, Yokohama, Yamashiro), Shikoku (Tokushima), Kiusu (Ariake Bay, Kagashima in Yatsushiro Bay, Higo), Riu-Kiu (Okinawa), Formosa, Corea.

70. **Platalea minor** Temminck & Schlegel.... *Kurotsura-herasagi*.

1849.—*Platalea minor* Temm. & Schl., Siebold's Fauna Japonica, Aves, p. 120, pl. 76 (Japan).

1877.—*Platalea japonica* Reichenow, Journ. f. Orn., p. 159.

(New name for *P. minor*).

Black-faced Spoonbill.

Hab. Hondo (Shimōsa), Kiusiu (Nagasaki), Gotō Is., Riū-Kiū (Okinawa), Formosa, Corea.

Order ANSERES.

Family ANATIDÆ.

Mergus L. (1758).

71. **Mergus merganser** *merganser* L. *Kawa-aisa*.

1758.—*Mergus Merganser* L., Syst. Nat., ed. X, I, p. 129
(Sweden).

Goosander.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Corea, Quelpart Is.

72. **Mergus merganser** *orientalis* Gould. *Ko-kawaaisa*.

1845.—*Mergus orientalis* Gould, Proc. Zool. Soc. London, p. 1
(Amoy in China).

Merganser comatus Salvadori.

Eastern Goosander.

Hab. Hondo (Mouth of Rokugō River, Koshiba near Yokohama), Kiusiu (Chikuzen).

73. **Mergus serrator** L. *Umi-aisa*.

1758.—*Mergus Serrator* L., Syst. Nat., ed. X, I, p. 129
(Sweden).

Serrator serrator (L.).

Red-breasted Merganser.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Riū-Kiū (Okinawa), Formosa, Corea, Quelpart Is.

Mergellus Selby (1840).

74. **Mergellus albellus** (L.)..... *Miko-aisa.*
1758.—*Mergus Albellus* L., Syst. Nat., ed. X, I, p. 129
("Habitat in Europa").

Smew.

Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Corea,
Quelpart Is.

Erionetta Coues (1884).

75. **Erionetta spectabilis** (L.)..... *Kewata-gamo.*
1758.—*Anas spectabilis* L., Syst. Nat., ed., X, I, p. 123
("Habitat in Canada, Svecia").

Somateria spectabilis (L.).

King Eider.

Hab. Kurile Islands.

Polysticta Eyton (1836).

76. **Polysticta stelleri** (Pallas)..... *Ko-kewata-gamo.*
1769.—*Anas stelleri* Pall., Spicil. Zool., pt. VI, p. 35 (Kamtschatka).
Iniconetta stelleri (Pallas).
Steller's Eider.
Hab. Kurile Islands, Hokkaido (Nemuro, Akiyuru Is.).

Histrionicus Lesson (1828).

77. **Histrionicus histrionicus pacificus** Brooks. *Shinori-gamo.*
1915.—*Histrionicus histrionicus pacificus* Brooks, Bull. Mus.
Comp. Zool. Harvard College, LIX, p. 393 (Kamtschatka).
Cosmonetta historionica (nec L.), Uchida.
Pacific Harlequin Duck.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Kawasaki, Yokohama, Banyū R., Nishiura in Izu), Kiusiu (Chikuzen), Corea, Quelpart Is.

Oidemia Fleming (1822).

78. **Oidemia nigra americana** Swainson..... *Kuro-gamo*.
1832.—*Oidemia Americana* Swainson, Swainson & Richardson,
Fauna Bor. Amer., II, p. 450 (Hudson Bay).

American Black Scoter.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Hitachi, Shimosa, Tokyo Bay, Yokohama, Enoshima in Sagami Bay, Shizuura in Suruga Bay), Corea.

Melanitta Boie (1822).

79. **Melanitta fusca stejnegeri** (Ridgway)..... *Birōdokinkuro*.
1887.—*Oidemia stejnegeri* Ridgway, Man. N. Amer. B., p. 112
("Kamtschatka to Japan").
Oidemia fusca stejnegeri Ridgway.
Oidemia fusca (nec L.), Dresser (pt.).

Eastern Velvet Scoter.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Teganuma in Shimōsa, Haneda, Misaki in Sagami, Mikawa Bay), Shikoku, Kiusiu (Chikuzen, Nagasaki), Corea.

Clangula Leach (1819).

80. **Clangula hyemalis** (L.)..... *Kōri-gamo*.
1758.—*Anas hyemalis* L., Syst. Nat., ed. X, I, p. 126 (North Sweden).
Harelda hyemalis (L.).

Long-tailed Duck.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Miyako in Rikuchū, Teganuma in Shimōsa), Corea.

Bucephala Baird (1860).

81. **Bucephala clangula clangula** (L.). *Hōjirō-gamo*.

1758.—*Anas Clangula* L., Syst. Nat., ed. X, I, p. 125 (Sweden).

1758.—*Anas Glaucion* L., op. cit., p. 126 (Sweden—♀).

Clangula glaucion (L.).

Golden-Eye.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Tsushima, Corea, Quelpart Is., Formosa.

Marila Reichenbach (1852).

82. **Marila marila mariloides** (Vigors). *Suzu-gamo*.

1839.—*Fuligula Mariloides* Vigors, Zool. Beechey's Voy.
Blossom, p. 31 (Northern Pacific Ocean).

Fuligula marila (nec L.), Uchida.

Nyroca marila mariloides (Vig.), Hartert.

Eastern Scaup-Duck.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
South Is. of Satsuma (Tanegashima), Riukiu, Formosa,
Corea, Quelpart Is.

Fuligula Stephens (1824).

83. **Fuligula fuligula** (L.). *Kinkuro-hajiro*.

1766.—*Anas fuligula* L., Syst. Nat., ed. X, I, p. 128 (Sweden).

Nyroca fuligula (L.), Hartert.

Tufted Duck.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Kiusiu, South Is.
of Satsuma (Yakushima), Riukiu (Ishigakijima), Formosa,
Corea.

Nyroca Fleming (1822).

84. **Nyroca ferina ferinoides** Kuroda. *Hoshi-hajiro*.

1920.—*Nyroca ferina ferinooides* Kuroda, "Dōbutsugaku Zasshi" (Tokyo Zool. Mag.), XXXII, pp. 243, 245 (Haneda near Tokyo).

Nyroca ferina (nec L.), Uchida.

Nyroca ferina ferina (nec L.), Hartert (pt.).

Eastern Pochard.

Hab. Hokkaido, Hondo (Shimosa, Haneda, Yokohama, Suruga), Kiusiu (Satsuma), Corea (Kōkai Distr., Keiki Distr.).

85. **Nyroca nyroca bæri** Radde, *Alkahajiro*.

1803.—*Anas (Fuligula) Bæri* Radde, Reise Sud. v. Ost-Sibirien, II, p. 376, pl. 15 (Amur).

Nyroca bæri Radde.

Nyroca nyroca nec Guld., Uchida.

Siberian White-eyed Duck.

Hab. Hokkaido, Hondo (Haneda, Yokohama), Corea (Keiki Distr.).

Spatula Boie 1822.

86. **Spatula clypeata** (L.) *Hashibiro-gamo*.

1758.—*Anas clypeata* L., Syst. Nat., ed. X, I, p. 124 (South Sweden).

1913.—*Spatula clypeata indianus* Math., Austr. Av. Rec. I, p. 194 (India).

Shoveller.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, South Is. of Satsuma (Tanegashima), Riukiu (Okinawa-jima) Formosa, Corea, Quelpart Is.

Daf'a Stephens (1824).

87. **Dafila acuta acuta** L.) *Onaga-gamo*.

1758.—*Anas acuta* L., Syst. Nat. ed. X, I, p. 126 (Sweden).

Dafila acuta (L.).

Anas acuta acuta L., Hartert.

Pintail.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
South Is. of Satsuma (Tanegashima, Formosa, Corea,
Quelpart Is.)

Eunetta Bonaparte (1856).

88. **Eunetta falcata** (Georgi).....*Yoshi-gamo.*
1775.—*Anas falcata* Georgi, Reise Russ. Reichs, I, p. 167
(Asiatic Russia).

Falcated Teal.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Formosa, Corea, Quelpart Is.

Mareca Stephens (1824).

89. **Mareca penelope** (L.)*Hidori-gamo.*
1758.—*Anas Penelope* L., Syst. Nat., ed. X, I, p. 126 (Sweden).
Wigeon.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
South Is. of Satsuma (Tanegashima), Formosa, Corea,
Quelpart Is.

90. **Mareca americana** (Gmelin)*Amerika-hidori.*
1789.—*Anas americana* Gmelin, Syst. Nat., I, 2, p. 526
(Louisiana).

American Wigeon.

Hab. Hondo (Uaneda, Shinhamai in Shimosha).

Querquedula Stephens (1824).

91. **Querquedula querquedula** (L.)*Shimajii.*

1758.—*Anas Querquedula* L., Syst. Nat., ed. X, I, p. 126
(Sweden).

Querquedula cincia (L.).

Garganey.

Hab. Sakhalien, Hondo (Haneda, Yokohama), Shikoku (Tokushima), Kiusiu (Nagasaki), Formosa (Tamsui), Corea.

Nettion Kaup (1829).

92. **Nettion formosum** (Georgi). *Tomew-gamo*.

1775.—*Anas formosa* Georgi, Reise Russ. Reichs, p. 168
(Lake Baikal).

Spectacled Teal.

Hab. Sakhalien, Hokkaido, Hondo, Kiusiu, Tsushima, Corea,
Formosa.

93. **Nettion crecca crecca** (L.). *Kō-gamo*.

1758.—*Anas Crecca* L., Syst. Nat., ed. X, I, p. 125 (Sweden).

Anas crecca crecca L., Hartert.

Teal.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Tsushima, South Is. of Satsuma (Tanegashima, Amami-
shima), Riukiu (Ishigakijima, Formosa, Corea, Quelpart
Is.).

94. **Nettion crecca carolinense** (Gmelin). *Amerika-kogamo*.

1789.—*Anas carolinensis* Gm., Syst. Nat., I, 2, p. 533 ("A
Carolina ad sinum Hudsonis").

Nettion carolinense (Gmelin), Kuroda.

Anas crecca carolinensis Gm., Hartert.

American Teal.

Hab. Hondo (Haneda near Tokyo).

Chaulelasmus Bonaparte (1838).

95. **Chaulelasmus streperus** (L.) *Okayoshi-gamo.*
1758. *Anas strepera* L., Syst. Nat., ed. X, I, p. 125 (Sweden).
Gadwall.
Hab. Kurile Is., Hokkaido, Hondo, ?Shikoku, Kiusiu, Corea
(Keiki Distr.), Quelpart Is.

Polionetta Oates (1899).

96. **Polionetta poecilorhyncha zonorhyncha** Swinhoe).
..... *Karu-gamo.*
1863. *Anas zonorhyncha* Swinh., Ibis, p. 394 (Ningpo).
Anas poecilorhyncha zonorhyncha Swinh., Harttert.
Dusky Mallard.
Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Tsushima, South Is. of Satsuma (Tanegashima), Riukiu
(Okinawa, Ishigaki, Iriomote), Formosa, Corea, Quelpart
Island.

Anas L. (1758).

97. **Anas platyrhyncha platyrhyncha** L. *Magamo.*
1758. *Anas platyrhynchos* L., Syst. Nat., ed. X, I, p. 125
(Sweden—♀).
1758. *Anas Boschas* L., op. cit., p. 127 (Sweden—♂).
Mallard.
Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Seven Is. (Hachijōjima), Bonin Is., Shikoku, Kiusiu, South Is. of Satsuma
(Tanegashima, Amamiōshima), Formosa, Corea, Quelpart
Is.

Pseudotadorna Kuroda (1917).

98. **Pseudotadorna cristata** Kuroda. *Kammuri-tsukushigamo.*

1917.—*Pseudotadorna cristata* Kuroda, "Tori," Vol. I, no. 5, p. 1, fig. i. (Near Fusan, Corea; id., "Semman Chōrui Ippan," p. 45, with coloured plate).

Pseudotadorna cornuta (errone) Kuroda, Hartert.

Crested Sheld-ake.

Hab. Corea (Naktung River or Rakutokō near Fusan).

Casarca Bonaparte (1838).

99. **Casarca ferruginea** (Pallas). *Aka-tsukushi-gamo*.

1764.—*Anas ferruginea* Pallas, Vroeg's Cat., Adumbratiuncula, p. 5 (Tartary).

Tadorna casarca (L.).

Casarca rutila (Pall.).

Ruddy Sheldrake.

Hab. Hondo (Uzen, Sendai, Shimosha, Haneda, Kiusiu (Higo, Nagasaki), Formosa, Corea, Quelpart Is.).

Tadorna Fleming (1822).

100 **Tadorna tadorna** L. *Tsukushi-gamo*.

1758.—*Anas Tadorna* L., Syst. Nat., ed. X, I, p. 122 (Sweden).

Tadorna cornuta (S. G. Gmelin).

Sheldrake.

Hab. Hokkaido (Akkeshi in Kushiro), Hondo (Tōkyō,? Suruga, Haneda), Kiusiu (Hizen, Ariyakenoumi), Riukiu (Ishigaki), Formosa, Corea, Quelpart Is.

Dendrocygna Swainson (1837).

101. **Dendrocygna javanica javanica** (Horsfield). *Riukiu-gamo*.

1821.—*Anas javanica* Horsf., Trans. Linn. Soc. London, XIII, p. 199 (Java).

Indian Tree Duck.

Hab. Riukiu (Okinawa, Ishigaki, Iriomote), Formosa.

Aix Boie (1828).

102. **Aix galericulata** (L.)..... *Oshidori*.

1758.—*Anas galericulata* L., Syst. Nat., ed. X, I, p. 128
(China).

1914.—*Aix galericulata brunneiceps* Clark, Proc. Biol. Soc. Washington, XXVII, p. 87 (Kiusiu, Japan; Shanghai).

Aix galericulata (L.).

Mandarin Duck.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, South Is. of Satsuma (Tanegashima, Amami-ōshima), Riukiu (Okinawa), Formosa, Corea, Quelpart Is.

Branta Scopoli (1769).

103. **Branta bernicla bernicla** (L.)..... *Nezumi-gan*.

1758.—*Anas Bernicla* L., Syst. Nat., ed. X, I, p. 124
("Habitat in Europa boreali; migrat supra Sveciam").

Branta bernicla (Linn.), Dresser.

Brent Goose.

Hab. ? Hokkaido, Japan (recorded by Blakiston & Pryer; Dresser).

104. **Branta bernicla nigricans** (Lawrence). *Koku-gan*.

1846.—*Anser nigricans* Lawr., Ann. Lyce. Nat. Hist., New York, IV, p. 171, pl. XII (New Jersey).

Black Brent.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Pref. Chiba, Tokyo Bay), Kiusiu (Chikuzen, Nagasaki), Corea.

Leucoblepharon Baird (1860).

105. **Leucoblepharon canadensis hutchinsii** (Richardson).
..... *Shijukara-gan.*
1831.—*Anser Hutchinsii* Richards., in Swainson & Richards.,
Fauna Bor. Amer. II, Birds, p. 470 (Melville Peninsula).
Branta canadensis hutchinsii (Rch.), Hartert.
Hutchins's Goose.
Hab. Kurile Is., Hokkaido (Hakodate), Hondo (Kazusa,
Shimōsa, Tokyo Bay).

Chen Boie (1822).

106. **Chen cærulescens cærulescens** (L.). *Haku-gan.*
1758.—*Anas cærulescens* L., Syst. Nat., ed. X, I, p. 124
("Habitat in Canada"). (Grey phase!).
1769.—*Anser hyperboreus* Pallas, Spicilegia Zool. fasc. VI, p.
25 (Coasts of Ice Seas). (White phase!).
Anser cærulescens cærulescens (L.), Hartert.
Anser hyperboreus nivalis (nec Forster), Uchida.
Snow-Goose.
Hab. Hokkaido, Hondo (Teganuma in Shimōsa, Matsudo,
Tokyo Bay, Yokohama), Kiusiu (Nagasaki), Corea (S.
Zenra Distr.).

Melanonyx Buturlin (1901).

107. **Melanonyx fabalis sibiricus** Alphéraky. *Ō-hishikui.*
1905.—*Melanonyx arvensis sibiricus* Alphéraky, Geese Europe
and Asia, p. 140, pl. 10 & 23 (Taimyr-Peninsula to
Chukchiland, Kamtschatka and Commander Is.).
Anser arvensis sibiricus (Alph.).
Anser fabalis sibiricus (Alph.), Hartert.

Middendorff's Goose.

Hab. Sakhalien, Hokkaido, Hondo (Hitachi, Shimōsa, Echigo),
Corea (Keiki Distr., S. Chusei Distr., Zenra Distr.).

108. *Melanonyx fabalis mentalis* (Oates). *Hashibuto-hishikui*.

1899.—*Anser mentalis* Oates, Man. Game-B. India, II, p. 77
(Yokohama).

Melanonyx segetum mentalis (Oates).

Anser fabalis serrirostris Swinh., Hartert (pt.).

Thick-billed Goose.

Hab. Hondo (Yokohama), Corea (Keiki Distr.).

109. *Melanonyx fabalis serrirostris* (Swinhoe). *Hishikui*.

1871.—*Anser segetum* var. *serrirostris* Swinh., Proc. Zool.
Soc. London, p. 417 (China).

Melanonyx segetum serrirostris (Sw.).

Anser fabalis serrirostris Swinh., Hartert (pt.).

Eastern Bean-Goose.

Hab. Kurile Is., Hokkaido, Hondo, ? Shikoku, Formosa,
Corea.

110. *Melanonyx fabalis fabalis* (Latham). *Hime-hishikui*.

1787.—*Anas fabalis* Latham, Gen. Synops., suppl., I, p. 297
(Great Britain).

Melanonyx segetum segetum (Gm.).

Anser fabalis fabalis (Lath.), Hartert.

Bean-Goose.

Hab. ? Hokkaido, Hondo, Corea, Quelpart Is.

Anser Brisson (1760).

111. *Anser erythropus* (L.). *Karigane, Ko-karigane*.

1758.—*Anas erythropus* L., Syst. Nat., ed. X, I, p. 123
("Habitat in Europa septentrionale").

Marilochen erythropus (L.).

Lesser White-fronted Goose.

Hab. Hokkaido, Hondo (Ogawaranuma in Mutsu, Kashima in Hitachi, Yokohama), Corea.

112. **Anser albifrons gambelli** Hartlaub..... *Ō-magan.*

1852.—*Anser Gambelli* Hartlaub, Rev. & Mag. zool., p. 7
(Texas and Southern United States).

North American White-fronted Goose.

Hab. ? Hondo, Corea.

113. **Anser albifrons albifrons** (Scopoli). *Magan.*

1769.—*Branta albifrons* Scopoli, Annus I Histor. Nat., p. 69.
(Probably from North Italy).

Anser albifrons (Scopoli).

White-fronted Goose.

Hab. Kurile Is., Hokkaido, Hondo, ? Shikoku, Kiusiu,
Formosa, Corea, Quelpart Is.

114. **Anser anser** (L.) *Haiiro-gan, Komakarigane.*

1758.—*Anas Anser* L., Syst. Nat., ed. X, I, p. 123 (Sweden).

1871.—*Anser cinereus* var. *rubrirostris* Swinhoe, Proc. Zool.
Soc. London, p. 416 (Shanghai).

Grey-Lag Goose.

Hab. Hondo (Teganuma in Shimōsa), Quelpart Is.

Cygnopsis Brandt (1836).

115. **Cygnopsis cygnoides** (L.). *Sakatsura-gan.*

1758.—*Anas Cygnoides* L., Syst. Nat., ed. X, I, p. 122
("Habitat in Asia").

Chinese Goose.

Hab. Kurile Is., Hokkaido, Hondo (Kazusa, Shimōsa, Yokohama), Formosa, Corea.

Cygnus Bechstein (1803).

116. **Cygnus olor** (Gmelin). *Kobu-hakuchō*.

1789.—*Anas Olor* Gmelin, Syst. Nat. I, 2, p. 501 (“Habitat in Russia, Sibiria, Persico etiam littore maris caspii, hospitatur per omnem Europam”).

Mute Swan.

Hab. Corea (Genzan, Mok-po).

Olor Wagler (1832).

117. **Olor bewickii jankowskii** (Alphéraky). *Hakuchō*.

1904.—*Cygnus bewickii jankowskii* Alph., Priroda i Okhota (Nature and Sport), p. 10 (Russian ! Ussuriland).

Cygnus bewicki (nec Yarrell), Uchida.

Eastern Bewick's Swan.

Hab. Hokkaido, Hondo (Tokyo Bay, Kanamachi in Musashi), Corea, Quelpart Island.

118. **Olor cygnus** (L.). *Ō-hakuchō*.

1758. *Anas Cygnus* L., Syst. Nat., ed. X, I, p. 122 (Sweden).

Cygnus musicus Bechstein.

Cygnus cygnus (L.), Haërtet.

Wooper Swan.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo (Awomori Bay, Kominato, Yabuki in Iwaki, Tokyo Bay), Kiusiu (Chikuzen, Yanagawa, Nagasaki, Yasaka in Bungo), Corea,

Order ACCIPITRES.

Family VULTURIDÆ.

Ægyptius Savigny (1809).

119. **Ægyptius monachus** (L.) *Hage-washi*.
1766.—*Vultur Monachus* L., Syst. Nat., ed. XII, I, p. 122
("Habitat in Arabia").
Cinereous Vulture.
Hab. Corea, Formosa (Tōen, Kagi).

Gypaëtus Storr (1784).

- 120 **Gypaëtus barbatus grandis** Storr. *Hige-washi*.
1784.—*Gypaëtus grandis* Storr, Alpenreise vom Jahre 1781,
p. 69 (Switzerland).
Gypaëtus barbatus (nec L.), Kuroda.
Bearded Vulture.
Hab. Corea (S. Kankyo Distr.).

Family FALCONIDÆ.

Aquila Brisson (1760).

121. **Aquila chrysaëtos chrysaëtos** (L.) *Inu-washi*.
1758.—*Falco Chrysactes* L., Syst. Nat., ed. X, I, p. 88
(Europe).
Aquila chrysactus (L.).
Golden Eagle.
Hab. Hokkaido, Hondo, Shikoku, Corea.
122. **Aquila heliaca heliaca** Savigny. *Katashiro-washi*.
1809.—*Aquila heliaca* Savig., Descr. Egypte, Syst. Ois., p. 82,
pl. 12 (Upper Egypt).
Imperial Eagle.

Hab. Hondo (Miyako in Rikuchu), Corea (Kōkai Distr., Keiki Distr., Mok-po), Formosa.

123. *Aquila clanga* Pallas..... *Karafuto-washi*.
1827.—*Aquila Clanga* Pallas, Zoogr. Rosso-Asiat. I, p. 351.
("In Rossia Siberiaque universa").
Aquila maculata (Gm.), Hartert.
Great Spotted Eagle.
Hab. Sakhalien, Corea. (Keiki Distr.).

Spizaetus Vieillot (1816).

124. *Spizaetus nipalensis* nipalensis (Hodgson)
..... *Taiwan-kumataka*.
1836.—*Nisactus Nipalensis* Hodgs., Journ. As. Soc. Bengal V, p. 229 (Nepal).
Himalayan Hawk-Eagle.
Hab. Formosa.
125. *Spizaetus nipalensis orientalis* Temminck & Schlegel.....
..... *Kumataka*.
1844—1845.—*Spizaetus orientalis* Temm. & Schl., Siebold's Fauna Japon., Aves, p. 7, pl. III (text 1844, pl. 1845) (Japan).
Japanese Hawk-Eagle.
Hab. Hokkaido, Hondo, Shikoku, Corea, ? Kiushiu.

Spilornis Gray (1840).

- 126 *Spilornis cheela hoya* Swinhoe..... *O-kammuritewashi*.
1866.—*Spilornis hoyi* Sw., Ibis, p. 304 (Formosa).
Spilornis cheela (nec Lath.), Uchida.
Formosan Serpent-Eagle.
Hab. Formosa.
127. *Spilornis cheela* (Lath.), subsp..... *Kammuritewashi*.

- 1905.—*Spilornis pallidus* (nec Walden), Ogawa, Annot. Zool. Japon., V, pt. 4, p. 213 (Yayeyama Is.).
- 1920.—*Spilornis cheela* subsp. ?, Swann, Syn. List Accipitres, p. 84. (Iriomote, Yayeyama, S. Loo Choo Is.).
- Loo Choo Serpent-Eagle.
- Hab. Riukiu (Ishigaki, Iriomote).

Butastur Hodgson (1843).

128. **Butastur indicus** (Gmelin). *Sashiba*.
- 1788.—*Falco indicus* Gm., Syst. Nat., I, 1, p. 264 (Java).
- Eastern Buzzard-Hawk.
- Hab. Hondo, Kiusiu, South Is. of Satsuma (Yakushima, Amamiōshima), Riukiu (Okinawa, Ishigaki), Formosa, Corea (Keiki Distr.).

Haliæetus Savigny (1809).

129. **Haliæetus albicilla** (L.). *Ojiro-washi*.
- 1758.—*Falco Albicilla* L., Syst. Nat., ed. X, I, p. 89 (Sweden).
- 1870.—*Haliaetus brooksi* Hume, Ibis, p. 438 (Upper India; without ordinary description).
- White-tailed Eagle.
- Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea, Quelpart Is.

Thallasoaëtus Kaup. (1844).

130. **Thallasoaëtus pelagicus pelagicus** (Pallas). *Ō-washi*.
- 1827.—*Aquila pelagica* Pall., Zoogr. Rosso-Asiat. I, p. 343 (Islands between Kamtschatka and America : Bering Is.).
- Haliaetus pelagicus* (Pallas).
- Steller's Sea-Eagle.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, South Is. of Satsuma (Amamiōshima), Riukiu, Corea (S. Kankyō Distr., Keiki Distr.).

131. **Thallasoëetus pelagicus niger** (Heude).....*Chosen-owashi*.
1887.—*Haliaëetus niger* Heude, Naturaliste, p. 95 (Corea).
1888.—*Haliaëetus branickii* Taczanowski, Proc. Zool. Soc. London p. 451 (Corea).
Black Sae-Eagle.
Hab. Corea (S. Heian Distr., Kōgen Distr., Keiki Distr.).

Milvus Lacépède (1799).

132. **Milvus lineatus lineatus** (Gray).....*Tobi*.
1832.—*Haliaëetus lineatus* Gray, Hardwicke's Ill. Ind. Zool., I, p. 1, pl. 18 (China).
1844—1845.—*Milvus melanotis* Temm. & Schl., Fauna Jap., Aves, p. 14, pl. 5, 5 B (1844, pl. 1845). (Japan).
Milvus atter melanotis (Temm. & Schl.), Uchida.
Milvus lineatus (Gray), Hartert.
Black-eared Kite.
Hab. Kurile Is., Hokkaido, Hondo, Seven Is., Shikoku, Kiusiu, Tsushima, South Is. of Satsuma (Tanegashima), Corea, Quelpart Is.

133. **Milvus lineatus formosanus** Kuroda.*Hime-tobi*.
1920.—*Milvus lineatus formosanus* Kuroda, "Dōbutsugaku Zasshi" (Tokyo Zool. Mag.), XXII, pp. 243, 245 (Formosa).

Milvus atter govinda (nec Sykes), Kuroda.
Formosan Black-eared Kite.
Hab. Formosa.

Pernis Cuvier (1817)

134. **Pernis apivorus orientalis** Taczanowski.....*Hachikuma*.

1891.—*Pernis apivorus orientalis* Temm., Faune Orn. Sib. Or., I, p. 50 (Eastern Siberia).

Pernis apivorus (nec L.), Uchida.

Siberian Honey Buzzard.

Hab. Hondo (Shinano, Nikko), Kiusiu, Corea, Formosa.

Buteo Lacépède (1799).

135. **Buteo ferox hemilasius** Temminck & Schlegel. *O-nosuri*.

1844.—*Buteo leucoccephala* Hodgson, Gray's Zool. Misc., p. 81. (Nomen nudum).

1844—1845.—*Buteo hemilasius* Temm. & Schl., Siebold's Faun. Japon., Aves, p. 16, pl. VII (text 1844, pl. 1845) (Japan).

Buteo leucoccephalus (Hodgs.), Uchida.

Upland Buzzard.

Hab. Kiusiu (Nagasaki), Tsushima, Corea.

136. **Buteo buteo japonicus** Temminck & Schlegel. *Nosuri*.

1844.—*Circus plumipes* Hodgson, Gray's Zool. Misc., p. 81 (Nomen nudum).

1844—1845.—*Falco buteo japonicus* Temm. & Schl., Siebold's Faun. Japon., Aves, p. 16, pl. VI & VIb (text 1844, pl. 1845). (Japan).

Buteo vulgaris (nec Leach), Uchida.

Japanese Buzzard.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Seven Is. (Miyakejima, Hachijōjima), Bonin Is., Shikoku, Kiusiu, Soth Is. of Satsuma (Amamiōshima), Tsushima, ? Dagelet Is., Quelpart Is., Corea, Formosa.

Archibuteo Brehm (1828).

137. **Archibuteolagopus pallidus** Menzbier. *Kashii-nosuri*.

- 1888.—*Archibuteo pallidus* Menzb., Orn. Turkestan, I, p. 163.
 (Siberia, Turkestan, Kamtschatka, Ussuriland).
Archibuteo lagopus (nec Gmelin), Uchida.
Buteo lagopus pallidus (Menzb.), Hartert.
 Siberian Rough-legged Buzzard.
 Hab. Hokkaido, Hondo (Shinano), Corea.

Circus Lacépède (1806).

138. **Circus aeruginosus aeruginosus** (L.).....*Chūhi*.
 1758.—*Falco aeruginosus* L., Syst. Nat., ed. X, I, p. 91
 (Sweden).
Circus aeruginosus (L.).
 Marsh-Harrier.
 Hab. Hokkaido, Hondo, Shikoku, Formosa, Corea.
139. **Circus spilonotus** Kaup.....*Shiberia-chūhi*.
 1850.—*Circus spilonotus* Kaup, Jardine's Contr. Orn., p. 59
 (Asia).
 Eastern Marsh-Harrier.
 Hab. Japan (locality unknown), Formosa (Kagi, Horisha),
 Corea (Keiki Distr.).
140. **Circus cyaneus cyaneus** (L.).....*Haiiro-chūhi*.
 1766.—*Falco cyaneus* L., Syst. Nat., ed. XII, I, p. 126
 (Region of London).
Circus cyaneus (L.).
 Hen-Harrier.
 Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Corea,
 Formosa.
141. **Circus melanoleucus** (Forster).....*Madara-chūhi*.
 1781.—*Falco melanoleucus* Forster, Indisch. Zool., p. 12, pl.
 XI (Ceylon).
 Pied Harrier.

Hab. Corea (Kōgen Distr., Keiki Distr.).

Astur Lacépède (1801).

- 142. *Astur gentilis schvedowi* Menzbier.** *Ō-taka*.
1882.—*Astur palumbarius Schvedowi* Menzb., Orn. Geogr.
Eur. Russia, p. 439 (Transbaikalia).
Astus palumbarius (nec L.), Uchida.
Accipiter gentilis schvedowi (Menzb.), Hartert.
Siberian Goshawk.
Hab. Sakhalien, Hokkaido, Hondo, Seven Is. (Hachijōjima),
Shikoku, Tsushima, Corea.
- 143. *Astur trivirgatus rufiinctus* McClelland.** *Taiwan-ōtaka*.
1839.—*Spizaetus rufiinctus* McClell., Proc. Zool. Soc.
London, p. 153 ('Burhampooter and other rives in
Assam').
Lophospiza trivirgata (nec Temm.), Uchida.
Large crested Goshawk.
Hab. Formosa (Horisha, Taitō).
- 144. *Astur badius poliopsis* (Hume).** *Takasago-daka*.
1874.—*Micronisus poliopsis* Hume, Stray Feathers, II, p. 325
(Northern Pegu).
Accipiter badius poliopsis (Hume), Hartert.
Hume's Shikra.
Hab. Formosa.
- 145. *Astur soloensis soloensis* (Latham).** *Akahara-daka*.
1821.—*Falco soloensis* Lath., Gen. Hist., I, p. 209.
Astur soloensis (Latham).
Accipiter soloensis Horsf., Hartert (pt.).
Horsfield's Goshawk.
Hab. Corea, (Seoul, Suigen), Formosa.

146. **Astur soloënsis cuculoides** (Temminck). *Wakijiro-daka*.
 1823.—*Falco cuculoides* Temm., Pl. Col., I, pl. 110, 129 (Java).
Astur cuculoides (Temm.).
Accipiter soloënsis Horsf., Hartert (pt.).
 Chinese Goshawk. ,
 Hab. Corea (Seoul, Seiryōri, Suigen).
- Accipiter** Brisson (1760).
147. **Accipiter nisus nisisimilis** (Tickell).
 *Haitaka* (♀), *Konori* (♂).
 1833.—*Falco Nisosimilis* Tick., Journ. As. Soc. Bengal, II, p. 571 (India).
Accipiter nisus (nec L.), Uchida.
 Indian Sparrow-Hawk.
 Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Seven Is.,
 Shikoku, Kiusiu, South Is. of Satsuma (Tanegashima),
 Tsushima, Corea.
148. **Accipiter nisus pallens** Stejneger. *Haiiro-haitaka*.
 1893.—*Accipiter pallens* Stejn., Proc. U. S. Nat. Mus., XVI,
 p. 625 (Prov. Hitachi, Japan).
 Kamtschatkan Sparrow-Hawk.
 Hab. Hondo (Hitachi, Rikuzen).
149. **Accipiter virgatus affinis** Hodgson. *Taiwan-tsumitaka*.
 1844.—*Accipiter affinis* Hodgs., Gray's Zool. Misc., p. 81.
 (Himalayas).
 Larger Besra Sparrow-Hawk.
 Hab. Formosa.
150. **Accipiter gularis** (Temminck & Schlegel)
 *Isumi* (♀), *Issai* (♂).
 1844—1845.—*Astur gularis* Temm. & Schl., Siebold's Faun.
 Japon., Aves, p. 5, pl. 2 (1844, pl. 1845). (Japan).

Accipiter virgatus (nec Temm.), Uchida.

Japanese Sparrow-Hawk.

Hab. Hokkaido, Hondo, Kiusiu, South Is. of Satsuma (Amami-Ōshima), Riukiu (Okinawa, Ishigaki), Formosa, Corea.

Falco L. (1758).

151. **Falco peregrinus calidus** Latham.....*Hayabusa*.
1790.—*Falco calidus* Latham, Ind. Orn., I, p. 41 (India).
Falco peregrinus (nec Tunstall), Uchida.
Siberian Peregrine Falcon.
Hab. Kirile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Seven Is. (? Miyakejima, Hachijōjima), South Is. of Satsuma (Amami-Ōshima), Riukiu (Iriomote). Formosa, Corea, Quelpart Is.
152. **Falco peregrinus pealei** Ridgway.....*O-hayabusa*.
1873.—*Falco communis* var. *pealei* Ridgw., Bull. Essex Inst., V, p. 201 (Oregon, N. America).
1929.—? *Falco rudolfi* Kleinschmidt, Falco, V, p. 19 (Hokodate).
Peale's Falcon.
Hab. Kurile Is., Hokkaido, Hondo (Sagauni), Volcano Is. (Minamiiwōjima), Kiusiu (Kagoshima).
153. **Falco peregrinus peregrinator** Sundevall.....
.....*Muncaka-hayabusa*.
1837.—*Falco peregrinator* Sundev., Physiogr. Sallskapets Tidskr. Lund, I, p. 177, pl. 4 (Indian Ocean, off Nicobar Is.).
Indian Peregrine Falcon.
Hab. Formosa (Horisha).
154. **Falco cherrug milvipes** Jerdon.....*Wakisuzi-hayabusa*.

- 1844.—*Falco milvipes* Hodgson, Gray's Zool. Misc., p. 81.
 (Nom. nud.).
- 1871.—*Falco milvipes* Jerdon, Ibis, p. 240 (Umballa in India)
Falco hendersoni Hume.
 Hodgson's Saker.
 Hab. Corea.
155. **Falco subbuteo jakutensis** Buturlin. *Chigo-hayabusa*.
 1910.—*Hypotriorchis subbuteo jakutensis* But., Nasoga Ochota,
 4, 6, p. 71 (N. E. Siberia).
 Kamtschatkan Hobby.
 Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Corea.
156. **Falco columbarius insignis** (Clark)..... *Kō-chōgenbō*.
 1907.—*Aesalon regulus insignis* Clark, Proc. U. S. Nat. Mus.,
 XXXII, p. 470 (Fusan, Korea).
Falco aesalon (nec Tunstall), Uchida.
 Asiatic Merlin.
 Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Corea.
- Hierofalco** Cuvier (1817).
157. **Hierofalco rusticola candicans** (Gmelin).... *Shiro-hayabusa*.
 1788.—*Falco candicans* Gm., Syst. Nat., I, 1, p. 275
 (Greenland).
 1882.—*Hierofalco uralensis* Sewertzoff & Menzbier, in
 Menzbier, Orn. Geogr. Europ. Russl., p. 288, pl. III (Ural
 Mountains).
Falco gryfalco (nec L.), Uchida.
 Greenland Falcon.
 Hab. Sakhalien, Kurile Is., Hokkaido.
- Erythropus** Brehm (1828).
158. **Erythropus vespertinus amurensis** (Radde)....
 *Akaashi-chōgenbō*.

1853.—*Falco vespertinus* L. var. *amurensis* Radde, Reisen,
i, S. von O. Sibir., II, p. 102, pl. I, fig. 2 (Amur).

Falco vespertinus amurensis Radde.

Eastern Red-footed Falcon.

Hab. Corea (N. Heian Distr.).

Cerchneis Boie (1826).

159. **Cerchneis tinnuncula japonica** (Temminck & Schlegel).....
.....*Chōsen-chōznbō*.

1844. —*Falco tinnunculus japonicus* Temm. & Schl., in Siebold's
Fauna Japon., Aves, p. 2, pl. I, and Ib. (Japan).

Japanese Kestrel.

Hab. Hondo, Shikoku, Kiushiu, South Is. of Satsuma
(Tanegashima), Riukiu, Formosa, Corea, Quelpart Is.

160. **Cerchneis tinnuncula perpallida** Clark.....*Chōsen-chōznbō*.
1907.—*Cerchneis perpallida* Clark, Proc. U. S. Nat. Mus.,
XXXII, p. 470 (Fusan, Korea).

1920.—*Cerchneis tinnuncula dorriesi* Swann, Syn. List
Accipitres, p. 146 (Sidemi & Amur River).

Falco tinnunculus perpallidus (Clark).

Falco tinnunculus tinnunculus (nec L.), Hartert (pt.).

Siberian Kestrel.

Hab. Corea (Fusan).

Family PANDIONIDÆ.

Pandion Savigny (1809).

161. **Pandion haliaetus haliaetus** (L.).....*Misago*.
1758.—*Falco Haliaeetus* L., Syst. Nat., ed. X, I, p. 91 (Sweden).

Pandion haliaeetus (L.).

Osprey.

Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
South Is. of Satsuma (Tanegashima, Amamiōshima),
Tsushima, Corea, Formosa.

Order HEMIPODII.

Family TURNICIDÆ.

Turnix Bonnaterre (1790).

162. **Turnix dussumieri** (Temminck)..... *Hime-mifuuzura*.
1828.—*Hemipodius dussumieri* Temm., Pl. Col., V, pl. 454,
fig. 2 (Bengal).
Little Button-Quail.
Hab. Formosa (Takow).
163. **Turnix tanki blanfordi** Blyth..... *Chōsen-mifuuzura*.
1863.—*Turnix Blanfordi* Blyth, Journ. As. Soc. Bengal,
XXXII, p. 80 (Thayetmyo, Burma).
Burmese Hemipode.
Hab. Corea (Keiki Distr.).

Areoturnix Bonaparte (1856).

164. **Areoturnix javanica blakistoni** (Swinhoe)..... *Mifu-usura*.
1871.—*Areoturnix blakistoni* Swinhoe, Proc. Zool. Soc.
London, p. 401 (Canton).
Chinese Hemipode.
Hab. Kiusiu (Satanomisaki in Ōsumi), South Is. of Satsuma
(Tanegashima), Riukiu (Okinawa, Okinoerabushima,
Miyakojima, Ishigakijima).
165. ? **Areoturnix javanica rostrata** (Swinhoe).....
..... *Taiwan-mifu-usura*.
1865.—*Turnix rostrata* Sw., Ibis, p. 543 (Formosa).
Formosan Hemipode.
Hab. Formosa (Takow).

Order GALLI.

Family PHASIANIDÆ.

Phasianus L. (1766).

166. *Phasianus colchicus karpowi* Buturlin. *Kōrai-kiji*.
1904.—*Phasianus karpowi* But., Orn. Monatsber, p. 3 (Te-lin
in S. Manchuria).
1907.—*Phasianus karpowi buturlini* Clark, Proc. U. S. Nat.
Mus., XXXII, p. 468 (Tsu-shima).
Phasianus torquatus (nec Gmelin), Uchida.
Corean Ring-necked Pheasant.
Hab. Corea, Quelpart Is., Tsushima. [Probably introduced
into Hondo (Chiba, Suruga, Mino) and Kiusiu (Ōshima in
Prov. Chikuzen)].
167. *Phasianus colchicus formosanus* Elliot. *Taiwan-kiji*.
1870.—*Phasianus formosanus* Elliot, Proc. Zool. Soc. London,
p. 406 (Formosa).
Formosan Ring-necked Pheasant.
Hab. Formosa.
168. *Phasianus versicolor versicolor* Vieillot. *Kiji*.
1825.—*Phasianus versicolor* Vieill., Gal. Ois., II, p. 23, pl. 205
(Japan).
1830.—*Phasianus Diardi* Temminck, Pl. Col., V, text to *P.
versicolor*.
Green Pheasant.
Hab. Hondo (excluding the north-western parts), ? Seven Is.
(Ōshima).
169. *Phasianus versicolor robustipes* Kuroda. *Hokuroku-kiji*.
1919.—*Phasianus versicolor robustipes* Kuroda, "Dōbutsugaku

Zasshi" (Tokyo Zool. Mag.), XXXI, pp. 299, 309 (Sado Is.).

Western Green Pheasant.

Hab. Hondo (Sado Is., including the north-western parts).

170. **Phasianus versicolor kiusiuensis** Kuroda.....*Kiusiu-kiji*.
1919.—*Phasianus versicolor kiusiuensis* Kuroda, "Dōbutsugaku Zashi" (Tokyo Zool. Mag.), XXXI, pp. 300, 309 (Kagoshima, Kiusiu).

Kiusiu Green Pheasant.

Hab. Island of Kiusiu.

171. **Phasianus versicolor tanensis** Kuroda.....*Shimakijū*.
1919.—*Phasianus versicolor tanensis* Kuroda, "Dōbutsugaku Zasshi" (Tokyo Zool. Mag.), XXXI, pp. 300, 310 (Tanegashima).

Southern Green Pheasant.

Hab. South Is. of Satsuma (Tanegashima, Yakushima).

Graphophasianus Reichenbach (1852).

172. **Graphophasianus sœmmerringii scintillans** Gould.....
.....*Yamadori*.
1866.—*Phasianus (Graphophasianus) scintillans* Gould, Ann.
& Mag. N. H. (3), XVII, p. 150 (Yokohama).
Phasianus scintillans Gould.

Hondo Copper Pheasant.

Hab. Hondo (north of about $35^{\circ} 20'$ N. lat.).

173. **Graphophasianus sœmmerringii intermedius** (Kuroda).....
.....*Shikoku-yamadori*.
1919.—*Phasianus sœmmerringii intermedius* Kuroda, "Dōbutsugaku Zasshi" (Tokyo Zool. Mag.), XXXI, pp. 304, 312
(Prov. Iyo, Shikoku).

Shikoku Copper Pheasant.

Hab. South-western parts of Hondo (Chūgoku), Shikoku.

174. *Graphphasianus sœmmerringii subrufus* (Kuroda).....

..... *U'suaka-yamadori*.

1919.—*Phasianus sœmmerringi subrufus* Kuroda, “Dōbutsu-gaku Zasshi” (Tokyo Zool. Mag.), XXXI, pp. 303, 311 (Prov. Suruga, Hondo).

Lesser Copper Pheasant.

Hab. Hondo (Pacific side of Hondo south of about 35° 20' N. lat.; S. Kazusa, S. Idu, coasts of Suruga, Iga, Ise, Nagato, etc.).

175. *Graphphasianus sœmmerringii sœmmerringii* (Temminck).....

..... *Aka-yamadori*.

1830.—*Phasianus sœmmerringii* Temm., Pl. Col., V, pls. 8, 9 (nos. 487, 488), Kiusiu).

Copper Pheasant.

Hab. North and Middle Kiusiu.

176. *Graphphasianus sœmmerringii ijimæ* (Dresser).....

..... *Koshijiro-yamadori*.

1902.—*Phasianus ijimæ* Dresser. Ibis, p. 655 (Prov. Hiuga, Kiusiu).

White-rumped Copper Pheasant.

Hab. Middle and South Kiusiu.

***Neocalophasis* Takatsukasa & Kuroda (1922).**

177. *Neocalophasis mikado* (Ogilvie-Grant)..... *Mikadokiji*.

1906.—*Calophasis mikado* O.-Grant, Bull. B. O. Club, XVI, p. 122 (Mt. Arizan, C. Formosa).

Mikado Pheasant.

Hab. Formosa (Central and Eastern mountainous parts of 6,000-9,000 feet above sea-level).

Hierophasis Elliot (1872).

178. **Hierophasis swinhoii** (Gould).....*Sanker.*
1862.—*Euplocamus swinhoii* Gould, Proc. Zool. Soc. London,
p. 284 (Formosa).
Gennaeus swinhoii (Gould).
Swinhoe's Kaleege.
Hab. Formosa (mountainous parts of 3,000-8,000 feet above
sea-level).

Bambusicola Gould (1862).

179. **Bambusicola sonorivox** Gould.....*Tekkei.*
1862.—*Bambusicola sonorivox* Gould, Proc. Zool. Soc.
London, p. 285 (Formosa).
Formosan Bamboo Pheasant.
Hab. Formosa.

Excalfactoria Bonaparte (1856).

180. **Excalfactoria chinensis chinensis** (L.).....*Hime-uszura.*
1766.—*Tetrao chinensis* L., Syst. Nat., I, p. 277 (China).
Painted Quail.
Hab. Formosa (Takow).

Coturnix Moehring (1752).

181. **Coturnix coturnix japonica** Temminck & Schleger....*Uzura.*
1844.—*Coturnix vulgaris japonica* Temm. & Schl., Siebold's
Fauna Japon., Aves, p. 103, pl. 61 (Japan).
Coturnix japonica Temm. & Schl.
Coturnix communis (nec Bonaparte), Uchida.
Japanese Quail.
Hab. Sakhalien, Kurile Is., Hokkaido, Hondo, Seven Is.
(? Miyakejima, Hachijojima, Shikoku, Kiusu, South Is.
of Satsuma Amami-ōshima), Tsushima, Corea, Formosa.

Arboricola Hodgson (1844).

- 182. Arboricola crudigularis** (Swinhoe), *Miyama-tekki*.
1864.—*Orcoperdix crudigularis* Swinh., Ibis, p. 426 (Formosa).
Formosan Tree-Partridge.
Hab. Formosa (mountainous parts of 3,000-8,000 feet above
sea-level).

Tetrastes Keys & Blasius (1840).

- 183. Tetrastes bonasia vicinitas** Riley..... *Yezo-raichō*.
1915.—*Tetrastes bonasia vicinitas* Riley, Proc. Biol. Soc.
Washington, Vol. XXVIII, p. 161 (Hakodate, Yezo,
Japan).
Tetrastes bonasia (nec L., Uchida).
Japanese Hazel Grouse.
Hab. Hokkaido.

- 184.? Tetrastes bonasia kolymensis** Buturlin.. *Karafuto-yezoraichō*.
1916.—*Tetrastes bonasia kolymensis* But., Mess. Ora. Dez.,
p. 226. (Middle Kolyma and Yakutsk).
Sakhalien Island Hazel Grouse.
Hab. Sakhalien Is.

- 185.? Tetrastes bonasia amurensis** Riley..... *Chōsen-yezoraichō*.
1916.—*Tetrastes bonasia amurensis* Riley, Proc. Biol. Soc.
Washington, Vol. XXIX, p. 17, Jan., 1916 (Near I-mien-
po, N. Kirin, Manchuria).
1916.—*Tetrastes bonasia ussuricensis* Buturlin, Messager
Ornith., Vol. 7, no. 4, pp. 222, 227, Dec., 1916 (S.
Ussuriland).

Amur Hazel Grouse.

Hab. Corea (N. Kankyo Distr., S. Kankyo Distr., N. Heian
Distr., Kōgen Distr., Keiki Distr.).

Falcipennis Elliot (1864).

- 186. Falcipennis falcipennis** (Hartlaub), *Kamabane-raichō*.

1855.—*Tetrao falcipennis* Hartl., Journ. f. Orn., p. 39 (Stanovoi Range).

Siberian Spruce Grouse.

Hab. Sakhalien Is.

Tetrao L. (1766).

187. **Tetrao parvirostris parvirostris** Bonaparte. *Ō-raichō*.

1856.—*Tetrao parvirostris* Bp., C. R., XLII, p. 880 (E. Siberia).

1884.—*Tetrao urogalloides* var. *sachalinensis* Bogdanow, Conspr. Av. Ross. fasc. i, p. 122 (Isl. of Saghalien).

Tetrao urogallus (nec L.), Uchida.

Siberian Capercailly.

Hab. Sakhalien Is. (Vicinity of the eastern boundary between Japan and Russia).

Lagopus Brisson (1760).

188. **Lagopus mutus japonicus** Clark. *Raichō*.

1907.—*Lagopus japonicus* Clark, Proc. U. S. Nat. Mus., Vol. XXXII, p. 469. (Mountains of Central Hondo, Japan).

Lagopus mutus (nec Montin), Uchida.

Japanese Ptarmigan.

Hab. Hondo (Norikura, Yarigatake, Ontake in Kiso, Rengesan, ? Hakusan, ? Nikko).

- 189.? **Lagopus mutus** (Montin), subsp. *Chishima-raichō*.

1898.—*Lagopus, species?*, Stejneger, Proc. U. S.

Nat. Mus., Vol. XXI, p. 285. (Northern Kurils).

Lagopus mutus subsp.?, Hartert.

Kamtschatkan Ptarmigan.

Hab. Kurile Is. (Simusir, Paramushir).

- 190. *Lagopus lagopus* lagopus (L.)** *Karafuto-raichō*.
 1758.—*Tetrao Lagopus* L., Syst. Nat., ed. X, I, p. 159
 (Swedish Lappland).
Lagopus albus (nec Gmelin), Uchida.
 Willow Grouse.
 Hab. Sakhalien Is., Kurile Is. (Recorded by Hartert).

Order ALECTORIDES.

Family RALLID.E.

Rallus L. (1766).

- 191. *Rallus aquaticus* indicus** Blyth..... *Kuina*.
 1849.—*Rallus indicus* Blyth, Journ. Asiat. Soc. Bengal,
 XVIII, p. 280 (Bengal).
 Eastern Water-Rail.
 Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Corea
 (Keiki Distr.).

Hypotænidia Reichenbach (1852).

- 192. *Hypotænidia striata* (L.)**, subsp. *Hashinaga-kuina*.
 1766.—*Rallus striatus* L., Syst. Nat., I, p. 262 (Philippine Is.).
 Blue-breasted Banded Rail.
 Hab. Formosa.

Rallina Reichenbach (1846).

- 193. *Rallina euryzonoides* sepiaria** (Stejneger).... *Riukiu-ōkuina*.
 1887.—*Euryzona sepiaria* Stejneger, Proc. U. S. Nat. Mus.,
 X, p. 395 (Liu kiu Islands).
Crex sepiaria (Stejneger).

Loo Choo Crake.

Hab. Riukiu (Ishigaki, Iriomote).

194. **Rallina euryzonoides formosana** Seebold....*Taiwan-ōkuina*.
1895.—*Rallina formosana* Seeb., Ibis, pp. 146, 211 (Formosa).
Formosan Crake.
Hab. Formosa.

Porzana Vieillot (1816).

195. **Porzana pusilla pusilla** (Pallas).*Hime-kuina*.
1776.—*Rallus pusillus* Pall., Reis. Russ. Reichs., III, p. 700
(Dauria).
Porzana pusilla (Pallas).
Pallas's Crake.
Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Riukiu
(Ishigaki), Corea.

Ortygops Heine (1890).

196. **Ortygops noveboracensis exquisita** (Swinhoe)...*Shima-kuina*.
1873.—*Porzana exquisita* Swinh., Ann. & Mag. Nat. Hist. (4),
XII, p. 376 (Chefoo).
Swinhoe's Crake.
Hab. Hokkaido, Hondo, Shikoku, Korea (Keiki Distr.).

Poliolimnas Sharpe (1893).

197. **Poliolimnas cinereus brevipes** (Ingram).*Jamijiro-kuina*.
1912.—*Porzana cinerea brevipes* Ingram, Bull. B. O. Club,
XXIX, p. 21 (Sulphur Is.).
Porzana cinerea (nec Vieill.), Uchida.
Sulphur Island Crake.
Hab. Sulphur Islands or Iwojima, south of Bonin Is.

Limnobænus Sundevall (1872).

198. **Limnobænus fuscus erythrothorax** (Temminck & Schleg.).
..... *Ili-kuina*.
1849.—*Gallinula erythrothorax* Temm. & Schl., Siebold's
Faun. Japon., Aves, p. 121, pl. LXVIII (Japan).
Porzana fusca (nec L.), Uchida.
Porzana fusca erythrothorax Temm. & Schleg., Hartert.
Japanese Ruddy Crake.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, South Is.
of Satsuma (Yakushima).
199. **Limnobænus fuscus phæopygus** (Stejneger).
..... *Riukiu-hikuina*.
1887.—*Porzana phæopyga* Stejneger, Proc. U.S. Nat. Mus., X,
p. 394 (Yayeyama Isl.).
Porzana fusca phæopyga Stejn., Hartert.
Loo-Choo Ruddy Crake.
Hab. South Is. of Satsuma (Amamiōshima), Riukiu (Okinawa-
jima, Ishigaki, Iriomote).
200. **Limnobænus fuscus fuscus** (L.). *Taiwan-hikuina*.
1766.—*Rallus fuscus* L., Syst. Nat., ed. XII, I, p. 262.
(Philippine Is.).
Porzana fusca fusca (L.), Hartert.
Ruddy Crake.
Hab. Formosa.
201. **Limnobænus paykullii** (Ljungh). *Kōrai-hikuina*.
1813.—*Rallus Paykullii* Ljungh, Kongl. Svenska. Vet. Akad.
Handlingar, p. 258, pl. V. (Java).
Porzana paykulli (Ljungh), Hartert.
Siberian Ruddy Crake.
Hab. Corea (N. Heian Distr., Keiki Distr.).

Amaurornis Reichenbach (1852).

202. **Amaurornis phoenicurus chinensis** (Boddäert).
..... *Shirohara-kuina*.
1783.—*Fulica chinensis* Bodd., Tabl. Pl. Enl., p. 54 (China).
Amaurornis phoenicurus (nec Forst.), Uchida.
Chinese White-breasted Water-hen.
Hab. Formosa.

Gallinula Brisson (1760).

203. **Gallinula chloropus parvifrons** Blyth. *Ban*.
1843.—*Gallinula parvifrons* Blyth, Journ. Asiat. Soc. Bengal,
XII, p. 180 (Calcutta).
Gallinula chloropus (nec L., Uchida.
Indian Water-hen.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, South Is. of Satsuma
(Amamiōshima). Riukiu (Okinawa, Ishigaki), Corea,
Dagelet Is.=Utsurioto.
204. ♀ **Gallinula chloropus lozanoi** Lletget *Takasago-ban*.
1918.—*Gallinula chloropus lozanoi* Lletget, Bol. R. Soc. Esp.
Hist. Nat., XVIII, p. 76 (Luzon, Philippine Is.).
Philippine Water-hen.
Hab. Formosa.

Gallicrex Blyth (1849).

205. **Gallicrex cinerea** (Gmelin). *Isuru-kuina*.
1789.—*Fulica cinerea* Gm., Syst. Nat., I, 2, p. 702 (China).
Water-cock.
Hab. Hondo (Ōsawa in Musashi, Namerikawa in Shimōsa,
Kamikamo in Yamashiro, Kōbe), Kiusiu (Nagasaki),
Riukiu (Iriomote), Formosa, Corea.

Fulica L. (1766).

- 206. Fulica atra atra** L. *Oban.*

1758.—*Flica atra* L., Syst. Nat. ed. X, I, p. 152 (Sweden).

1849.—*Fulica atra japonica* Temm. & Schl., Siebold's Faun.

Japon., Aves, p. 120, pl. LXXVII (Japan).

Coot.

Hab. Sakhalien, Hokkaido, Hondo, Shikoku, Kiusiu, Riukiu, Formosa, Corea.

Family BALEARICIDÆ. (Gruidæ).

Megalornis Gray (1841).

- 207. Megalornis grus lilfordi** (Sharpe) *Kuro-tsuru.*

1894.—*Grus lilfordi* Sharpe, Cat. Brd. Brit. Mus., Vol.

XXIII, p. 252 (E. Siberia, Turkestan, N.-W. India, etc.).

Grus communis (nec Bechstein), Uchida.

Eastern Common Crane.

Hab. Corea (Keiki Distr., Kōgen Distr.).

- 208. Megalornis nigricollis** (Przewalski) *Oguro-zuru.*

1876.—*Grus nigricollis* Przewalski, Mongolija i strana Tangut

II, p. 135 (1876); id. in Rowley's Orn. Misc., II, p. 436, pl. IX (1847) (Kokonor).

Black-necked Crane.

Hab. Corea (Keiki Distr.).

- 209. Megalornis japonensis** (P. L. S. Müller) *Tanchō.*

1776.—*Ardea (Grus) Japonensis* Müller, Syst. Nat. Suppl., p. 110 (Japan).

Grus japonicus (P. L. S. Müll.).

Japanese Crane or Mantchurian Crane.

Hab. Hokkaido (Iburi), Kiusiu (Nagasaki, Akune in prefect.

Kagoshima), Corea (Mok-po, Keiki Distr., S. Chusei Distr., S. Keishō Distr., Kokai Distr.).

210. *Megalornis monachus* (Temminck). *Nabe-zuru.*

1835.—*Grus monachus* Temm., Pl. Col., V, pl. 555 (Hokkaido and Corea).

Hooded Crane.

Hab. Hondo (Tokyo, Yokohama, Yashiro-mura in Prefect. Yamaguchi), Kiusiu (Nagasaki, Akune in Prefect. Kagoshima), Corea (Mok-po, Fusān, Keiki Distr., Genzan), Quelpart Is.

***Pseudogeranus* Sharpe (1893).**

211. *Pseudogeranus vipio* (Pallas). *Manazuru.*

1827.—*Grus Vipio* Pall., Zoogr. Rosso-Asiat., II, p. 111 (Near Nerchinsk, A. Russia).

1838.—*Grus leucachen* Temm., Pl. Col., V, pl. 449 (Japan).
Megalornis vipio (Pall.), Hartert.

White-naped Crane or White-necked Crane.

Hab. Hokkaido, Kiusiu (Nagasaki, Akune in Prefect. Kagoshima), South Is. of Satsuma (Tanegashima), Corea (Mokpo, Fusān, Seoul, N. Chusei Distr., Daidōkō), Quelpart Is.

***Sarcogeranus* Sharpe (1893).**

212. *Sarcogeranus leucogeranus* (Pallas). *Sodeguro-zuru.*

1773.—*Grus Leucogeranus* Pallas, Reis. Russ. Reichs, II, p. 714, tab. F. (Siberia).

Megalornis leucogeranus (Pall.), Hartert.

Siberian White Crane.

Hab. Spring and autumn visitor to Japan. (Seeböhm).

Anthropoides Vieillot (1816).

213. **Anthropoides virgo** (L.) *Ineha-zuru.*
1758.—*Irdea Virgo* L., Syst. Nat., ed. X, I, p. 141 (“Habitat
in Oriente”);
Grus virgo (L.).

Demoiselle Crane.

Hab. Hokkaido, Hondo (Izu).

Family OTIDIDÆ.

Otis L. (1766).

214. **Otis tarda dybowskii** Taczanowski..... *No-gan.*
1874.—*Otis Dybowskii* Tacz., Journ. f. Orn., p. 331 (Dauria).
Siberian Bustard.
Hab. Hokkaido, Hondo (Maiko, Kuwana, Inaba, Saku in
Shinano), Kiusiu, Yashai, Minamimatsuura-gōri in Prefect.
Nagasaki, Corea (Fusan, Chemulpo, Seoul, Kokai Distr.,
Shingishu).

Order LIMICOLÆ

Family CURSORIIDÆ.

Glareola Brisson (1760).

215. **Glareola pratincola maldivarum** Forster..... *Tsubame-chidori*
1795—*Glareola (Pratincola) Maldivarum* Forster, Faun.
Indica p. 11. (Maldive Islands).
Glareola orientalis Leach, Uchida.
Glareola maldivarum Forster, Hartert.

Oriental Pratincole.

Ha Hondo (Hitachi), Formosa.

Family CHARADRIIDÆ.

Charadrius L. (1758).

- 216. Charadrius placidus** Gray *Ikaru-chidori*
1863.—*Charadrius placidus* Gray, Cat. Mamm. B. & C. of
Nepal, Tibet in Bsit Mus., II. ed., p. 70. (Nepel).
Long-billed Ringed Plover.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Corea.
- 217. Charadrius dubius curonicus** Gmelin *Kō-chidori*
1789.—*Charadrius curonicus* Gmelin, S. N. I. 2, p. 692.
("Habitat in Curonia").
Charadrius minor Wolf & Meyer, Uchida.
Little Ringed Plover.
Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo
Ils., Formosa, Corea.
- 218. Charadrius dubius dubius** Scopoli *Minami kō-chidori*
1786.—*Charadrius dubius* Scopoli, Del. Fann. Flora. Insubr.
II, p. 93. (Luzon).
. *Egialitis dubia dubia* (Scop.), Kuroda.
Hab. Hondo, Formosa.
- 219. Charadrius hiaticula tundræ** (Lowe) *Hajiro-kō-chidori*
1915.—*Aegialitis hiaticola tundrac* Lowe, Bull. B. O. C.
XXXVI, p. 7. (Russia & Siberia).
Lowe's Ringed Plover.
Hab. Hondo (Sagami).
- 220. Charadrius alexandrinus alexandrinus** L. *Hashiboso-*
shiro-chidori.
1758.—*Charadrius alexandrinus* L., S. N. X. I, p. 150. (Egypt).
Kentish Plover.
Hab. Hokkaido, Hondo, Formosa, Corea.
- 221. Chardrius alexandrinus dealbatus** (Swinhoe) ... *Shiro-chidori*

1870.—*Aegialites dealbatus* Swinhoe, P. Z. S., p. 138. (S. China, Formosa & Hainan).

Charadrius cantarus Latham, Uchida.
Eastern Kentish Plover.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Amami-Ōshima,
Loo-choo, Formosa, Corea.

Cirrepidesmus Bonaparte (1856).

- 222. *Cirrepidesmus mongolus* mongolus** (Pallas) *Medai-chidori*.
1776.—*Charadrius mongolus* Pallas. Reise d. versch. Prov.
Russ. Reichs, III, p. 700. (Mongolia).
Mongolian Plover.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku,
Kiusiu, Loo-choo, Formosa, Corea.

Pagoa Mathews (1913).

- 223. *Pagoa leschenaultii* (Lesson).....** *Ō-medai-chidori*
1826.—*Charadrius Leschenaultii* Lesson, Dict. Sci. Nat. XLII,
p. 36. (Pondichery in India).
Charadrius geoffroyi Wagler, Uchida.
Geoffroy's Sand-Plover.
Hab. Saghalin, Hondo, Seven Islands (Izu), Kiusiu, Loo-choo
Isls., Formosa, Corea.

Eudromias Brehm (1831).

- 224. *Eudromias morinellus* (L.).....** *Kobashi-chidori*.
1758.—*Charadrius Morinellus* L., S. N. X, I, p. 150. (Sweden).
Dotterel
Hab. Saghalin, Hokkaido, Hondo (Sagami).

Eupoda Brandt (1845).

- 225. *Eupoda asiatica vereda* (Gould)** *Ō-chidori*

1848.—*Charadrius veredus* Gould, P. Z. S. p. 38. (North Australia).

Ochthodromus veredus (Gould). Uchida.

Eastern Dotterel

Hab. Corea, Japan.

Pluvialis Brisson (1760).

226. **Pluvialis dominicus fulvus** (Gm.) *Munaguro*.

1789.—*Charadrius fulvus* Gmelin, S. N. X, I, 2, p. 687. (Tahiti).
Eastern Golden Plover.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Islands
(Izu), Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.

Squatarola Cuvier (1816).

227. **Squatarola squatarola hypomelaena** (Pallas) *Daizen*

1776.—*Charadrius hypomelas* Pallas, Reise d. versch. Prov.
Russ. Reiches, III, p. 699. ("Colit paludes borealis orae").
Squatarola helvetica (L.), Uchida.

Hab. Saghalin, Kuril Isls., Hokkaido, Hondo, Shikoku, Loo-
choo Isls., Formosa, Corea.

Vanellus Brisson (1760).

228. **Vanellus vanellus** (L.) *Tur-geri*

1758.—*Tringa Vanellus* L., S. N. X, I, p. 148. (Sweden).

Vanellus vulgaris Bechstein, Uchida.

Lapwing.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Isls.,
Formosa, Corea.

Microsarcops Sharpe (1896).

229. **Microsarcops cinereus** (Blyth) *Keri*

1842.—*Pluvianus cinereus* Blyth, Jour. As. Soc. Bengal XI,
p. 584. (Calcutta).

Lobivanellus cinereus (Blyth), Uchida.

Hab. Hokkaido, Hondo, Kiusiu, Formosa, Corea.

Arenaria Brisson (1760).

230. **Arenaria interpres interpres** (L.) *Kyōjo-shigi*.

1758.—*Tringa Interpress* L., S. N. X, I, p. 148. (Sweden).

Turnstone.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Kiusiu, Loo-choo, Formosa, Corea.

Erolia Vieillot (1816).

231. **Erolia ferruginea** (Brünnich) *Saruhama-shigi*

1764.—*Tringa Ferruginea* Brünnich, Orn. Bor., p. 53. (Iceland
& Christiansö).

Tringa subarquata (Güldenstедт), Uchida.

Curlew-Sandpiper.

Hab. Hondo (Gyōtoku, Sagami) Formosa, Marcus Is.

Pelidona Cuvier (1817).

232. **Pelidona alpina sakhalina** (Vieillot) *Hama-shigi*

1816.—*Scolopax sakhalina* Vieillot, Nonv. Dict. d'His Nat.
III, p. 359. (Sakhalin Is.)

Tringa alpina pacifica (Couch), Uchida.

Pacific Dunlin.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Pescadres Isls., Corea, Quelpart Is.

Pisobia Billberg (1828).

233. **Pisobia minuta ruficollis** (Pallas) *Tōnen*

1776.—*Trynga ruficollis* Pallas, Reise d. versch. Prov. Russ.

Reichs, III, p. 700. ("Circa Lacus salsos Davuriae campestris").

Eastern Little Stint.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Islands (Izu), Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.

234. Pisobia minutilla subminuta (Middendorff).....*Hibari-shigi*.

1851.—*Tringa subminuta* Middendorff, Reise N. O. & O. Sibir. II, 2, p. 222, pl. XIX, fig. 6. (Stanovoi Range).

Tringa damaccensis Horsfield, Uchida.

Long-toed Stint.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Kiusiu, Formosa, Corea.

235. Pisobia temminckii (Leisler).....*Ojiro-tōnen*.

1812.—*Tringa Temminckii* Leisler, Nacht. z. Bechst. Naturg. Deutsch., p. 63–73. (Main, Germany).

Temminck's Stint

Hab. Kurile Isls., Hondo, Formosa.

Heteropygia Coues (1861).

236. Heteropygia acuminata (Horsfield).....*Uzura-shigi*.

1821.—*Totanus acuminatus* Horsfield, Tr. L. Soc. London XIII, p. 192. (Java).

Tringa acuminata Horsf., Uchida.

Siberian Pectoral Sandpiper.

Hab. Kurile Isl. Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

Canutus Brehm (1831).

237. Canutus canutus rogersi Mathews*Ko-oba-shigi*

1913.—*Canutus canutus rogersi* Mathews, B. Austr. III, p. 270. ("Siberia.....Australia in Winter")

Eastern Knot.

Hab. Hokkaido, Hondo (Gyōtoku), Formosa, Corea.

Anteliotringa Mathews (1913).

238. **Anteliotringa tenuirostris** (Horsfield) *Oba-shigi*.
1821.—*Totanis tenuirostris* Horsfield, Tr. L. Soc. London.
XIII, p. 192. (Java).
Tringa crassirostris Temm. & Schl., Uchida.
Great Knot.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Kiusiu,
Formosa, Corea, Quelpart Isl.

Arquatella Baird (1858).

239. **Arquatella maritima quarta** (Hartert) *Chishima-shigi*.
1920.—*Irolia maritima quarta* Hartert, Nov. Zool., p. 137.
(Commander Islands).
Tringa striata couesi (Ridgway), Uchida.
Hab. Kurile Islands.

Philomachus Anonymus (1804).

240. **Philomachus pugnax** (L.) *Irimaki-shigi*.
1758.—*Tringa Pugnax* L., S. N. X, I, p. 148. (Sweden).
Totanus pugnax (L.), Uchida.
Ruff & Reeve.
Hab. Hokkaido, Hondo (Rikuzen, Shimōsa, Sagami), Corea.

Tryngites Cabanis (1856).

241. **Tryngites subruficollis** (Vieillot) *Komon-shigi*.
1819.—*Tringa subruficollis* Vieillot, Nouv. Dict. d'Hist. Nat.
XXXIV, p. 415. (Paraguay).
Buff-breasted Sandpiper.
Hab. Hondo (Straggler!).

Crocethia Billberg (1828).

242. **Crocethia alba** (Pallas) *Miyubi-shigi*.

1766.—*Tringa alba* Pallas, Vroeg's Cat. Coll., Adumbratincula,
p. 7. (Holland).

Tringa arcuaria L., Uchida.

Sanderling.

Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Formosa, Corea,
Quelpart Is.

Limicola Koch (1816).

243. **Limicola falcinellus sibirica** (Dresser) *Kirai.*

1876.—*Limicola sibirica* Desser, P. Z. S., p. 674. (Siberia,
China).

Limicola platyrhyncha (Temm), Uchida.

Eastern Broad-billed Sandpiper.

Hab. Hokkaido, Hondo, Kiusiu, Formosa, Corea.

Eurynorhynchus Nilsson (1821).

244. **Eurynorhynchus pygmaeus** (L.) *Hera-shigi.*

1758.—*Platalea pygmaea* (sic !) L., S. N. X, I, p. 140. (Surinam-
errore !)

Tringa pygmaea (L.), Uchida.

Spoon-billed Sandpiper.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Kiusiu, Corea.

Macrorhamphus Forster (1817).

245. **Macrorhamphus semipalmatus** Blyth *O-hashi-shigi.*

1848.—*Macrorhamphus semipalmatus* "Jerdon" Blyth, Journ.
As. Soc. Bengal XVII, I. p. 252. (Near Calcutta).

Macrorhamphus griseus (nec Gmelin), Uchida.

Hab. Hokkaido, Hondo.

Erythroscelus Kaup (1829).

246. **Erythroscelus erythropus** (Pallas) *Tsuru-shigi.*

1764.—*Scolopax erythropus* Pallas, Vroeg's Cat. Coll., Adumbratinnula, p. 6. (Holland).

Totanus fuscus (L.), Uchida.

Spotted or Dusky Redshank.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Tsushima, Corea.

247. **Erythroscelus totanus eurhinus** (Oberholser)...*ika-ashi-shigi*.

1900.—*Totanus tetinus eurhinus* Oberholser, P. U. S. Nat. Mus. XXII, p. 207. ("Central and Eastern Asia").

Eastern Redshank.

Hab. Saghalin, Hondo, Formosa, Corea.

Iliornis Kaup (1829).

248. **Iliornis stagnatilis** (Bechstein)*Ko-awo-ashi-shigi*.

1803.—*Totanus stagnatilis* Bechstein, Orn. Taschent. II, p. 292. (Germany).

Marsh Sandpiper.

Hab. Hondo, Formosa, Corea.

Glottis Koch (1816).

249. **Glottis nebularius** (Gunnerus).....*Aeo-ashi-shigi*.

1767.—*Scolopax nebularia* Gunnerus, in Leem, Beskr. Finm. Lapp., p. 251. (Norway).

Totanus glottis (L.), Uchida.

Green-Shank.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

Pseudoglottis Stejneger (1884).

250. **Pseudoglottis guttifer** (Nordmann) *Karafuto-awo-ashi-shigi*.

1835.—*Totanus guttifer* Nordmann, Ermann Reise, Naturh.

Atlas., p. 17. (Ochotsk).

Nordmann's Greenshank.

Hab. Saghalin, Hondo (Rokugō), Kiusiu (Satsuma), Corea.

Tringa L. (1758).

251. **Tringa ochropus** L. *Kusa-shigi.*

1758.—*Tringa Ochromalus* L., S. N. X, I, p. 149. (Sweden).

Totanus ochropus (L.), Uchida.

Green Sandpiper.

Hab. Saghalin, Kuril Isls., Hokkaido, Hondo, Hachijō (Seven Islands, Izu), Shikoku, Kiusiu, Formosa, Corea.

Rhyacophilus Kaup (1829).

252. **Rhyacophilus glareolus** (L.) *Takabu-shigi.*

1758.—*Tringa Glareola* L., S. N. X, I, p. 149. (Sweden).

Totanus glareola (L.), Uchida.

Wood Sandpiper.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Kiusiu, Loochoo, Formosa, Corea.

Heteroscelus Baird (1858).

253. **Heteroscelus incanus brevipes** (Vieillot) *Kiashi-shigi.*

1816.—*Totanus brevipes* Vieillot, Nouv. Dict. d'Hist. Nat.

(nouv. éd) VI, p. 410. (Locality not given).

Totanus incanus (Gmelin), Uchida.

Asiatic Wandering Tattler.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Isls. (Izu), Bonin Islands, Shikoku, Kiusiu, Tsushima, Loochoo Isls., Formosa, Corea.

254. **Heteroscelus incanus incanus** (Gmelin) *Meriken-kiashi-shigi.*

1789.—*Scolopax incana* Gmelin, S. N. I, 2, p. 658 (Pacific

Ocean).

- Totanus incanus brevipes* (nec Vieillot), Uchida.
American Wandering Tattler.
Hab. Hondo, Bonin Is., Marcus Is., Shikoku, Loo-choo Isls.

Actitis Illiger (1811).

255. **Actitis hypoleucus** (L.) *Iso-shigi*.
1758.—*Tringa Hypoleucus* L., S. N. X, I, p. 149. (Sweden).
Totanus hypoleucus (L.), Uchida.
Common Sandpiper.
Hab. Saghalin, Kurile Isls, Hokkaido, Hondo, Bonin Islands,
Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.

Terekia Bonaparte (1838).

256. **Terekia cinerea** (Güldenstadt) *Sorihashi-shigi*.
1774.—*Scolopax cinerea* Güldenstadt, Nov. Comm. Petrop.
XIX, p. 473, pl. 19 (Coasts of Caspian Sea).
Totanus terekis (Latham), Uchida.
Terek Sandpiper.
Hab. Sagalin, Hokkaido, Hondo, Shikoku, Kiusiu, Formosa,
Corea.

Phalaropus Brisson (1760).

257. **Phalaropus fulicarius** (L.) *Haiiro-hireashi-shigi*.
1758.—*Tringa Fulicaria* L., S. N. X, I, p. 148. (Hudson Bay).
Red Phalarope.
Hab. Kurile Islands, Hondo.

Lobipes Cuvier (1817).

258. **Lobipes lobatus** (L.) *Akoyeri-hireashi-shigi*.
1758.—*Tringa lobata* L., S. N. X, I, p. 148. (Hudson Bay).

Phalaropus hyperboreus (L.), Uchida.

Red-necked Phalarope.

Hab. Sagalin, Kurile Isls., Hokkaido, Hondo, Loo-choo Isls., Formosa, Corea.

Himantopus Brisson (1760).

259. **Himantopus himantopus himantopus** (L.).....*Scitaka-shigi*.

1758.—*Charadrius Himantopus* L., S. N. X, I. p. 151. (Europe).

Himantopus himantopus, (L.) Uchida.

Black-winged Stilt.

Hab. Sulpher Islands, Formosa.

Recurvirostra L. (1766).

260. **Recurvirostra avosetta avosetta** L....*Sorihashi-scitaka-shigi*.

1758.—*Recurvirostra Avosetta* L., S. N. X, I. p. 151. (Europe). Avocet.

Hab. Formosa, Corea.

Limosa Brisson (1760).

261. **Limosa lapponica baueri** Naumann*O-sorihashi-shigi*.

1836.—*Limosa Baueri* Nanmann., Nat. Vög. Deutsch. VIII. p. 429. (New Zealand).

Limosa rufa uropygialis (Gould), Uchida.

Eastern Bar-tailed Godwit.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa Corea.

262. **Limosa limosa melanurooides** (Gould).....*O-suro-shigi*.

1846.—*Limosa melanurooides* Gould, P. Z. S., p. 84. (Port Essington).

Limosa melanura (nec Leisler), Uchida.

Eastern Black-tailed Godwit.

Hab. Saghalin, Hakkaido, Hondo, Kiusiu, Formosa, Corea.

Numenius Brisson (1760).

263. **Numenius arquatus** lineatus Cuvier. *Dri-shaku-shigi*.

1829.—*Numenius lineatus* Cuvier, Règne Anim. nouv. éd. I. p. 521. (India).

Indian Curlew.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.

264. **Numenius cyanopus** Vieillot *Tōroku-shigi*.

1817.—*Numenius cyanopus* Vieillot, Nouv. Dict. d'Hist. Nat. VIII, p. 306. (Australia).

Australian Curlew.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

Phaeopus Cuvier (1816).

265. **Phaeopus phaeopus** variegatus (Scopoli) *Chu-shaku-shigi*.

1786.—*Tantalus variegatus* Scopoli, Del. Flor. et Faun. Insubr., II. p. 92. (Luzon).

Numenius phaeopus variegatus Scopoli, Uchida.

Eastern Whimbrel.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Isls. (Izu), Bonin Isls., Shikoku, Kiusiu, Formosa, Corea.

266. **Phaeopus tahitiensis** (Gmelin) *Harimomo-chūshaku*.

1788.—*Scolopax tahitiensis* Gm., S. N., I, p. 656 (Tahiti).

Bristle-thighed Whimbrel.

Hab. Hondo (Yamashiro). (Straggler!).

Mesoscolopax Sharpe (1896).

267. **Mesoscolopax minutus** (Gould) *Ko-shaku shigi.*
1841.—*Numenius minutus* Gould., P. Z. S., p. 176. (New South Wales).

Least Whimbrel.

Hab. Hondo, Kiusiu, Formosa, Corea.

Scolopax L. (1766).

268. **Scolopax rusticola rusticola** L. *Kumri-shigi.*
1758.—*Scolopax Rusticola* L., S. N. X, I, p. 146. (Sweden).
Wood cock.
Hab. Sighalin, Hokkaido, Hondo, Seven Isls. (Izu.), Shikoku, Kiusiu, Satsunan Islands, Formosa, Corea, Quelpart Isl.
269. **Scolopax rusticola mira** Hartert *Amami-yama-shigi.*
1916.—*Scolopax rusticola miro* Hartert, Bull. B. O. C. XXX-VI, p. 64. (Amami-ōshima).
Hab. Amami-ōshima.

Gallinago Koch (1816).

270. **Gallinago gallinago raddei** Buturlin *Ta-shigi.*
1912.—*Scolopax (Gallinago) gallinago raddei* Buturlin, “Kuliki Rossieskoi Imperie-Premiyu-k-Journal”, p. 54. (East Siberia).
Gallinago celestris (Frenz.), Uchid i.
Hab. Saghulin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.

Note. Dr. Hartert said in his “Vög. paläarkt. Faun., p. 1658 & 1660” that the Common Snipe of ‘Far eastern Asia’ belongs to ‘raddei’ and not to ‘gallinago.’

Ditelmatias Mathews (1913).

271. **Ditelmatias hardwickii** (Gray) *Ō-jishigi.*

1831.—*Scolopax hardwickii* Gray, Zool. Misc., p. 16. (Tasmania).

Gallinago australis (Latham), Uchida.

Latham's Snipe.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiushiu, Formosa.

Subspilura Mathews (1913).

272. **Subspilura megala** (Swinhoe) *Chu-jishi*.

1861.—*Gallinago megala* Swinhoe, Ibis, p. 343. (Peking).

Swinhoe's Snipe.

Hab. Saghalin, Hokkaido, Hondo, Kiushiu, Tsushima, Loo-choo, Formosa, Corea.

Spilura Bonaparte (1856).

273. **Spilura stemura** (Bonaparte) *Hario-shigi*.

1830.—*Scolopax stemura* "Kuhl" Bonaparte, Ann. Stor. Nat. Bologna IV, p. 335. (Sunda Islands).

Gallinago stemura (Kuhl), Uchida.

Pin-tailed Snipe.

Hab. Saghalin, Hokkaido, Hondo, Formosa, Corea.

Neospilura Mathews (1913).

274. **Neospilura solitaria** (Hodgson) *Iwo-shigi*.

1831.—*Gallinago solitaria* Hodgson, Gleanings in Science III., p. 238. (Nepal).

Solitary Snipe.

Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiushiu, Loo-choo Irls., Corea.

Lymnocryptes Kaup (1829).

275. **Lymnocryptes gallinula** (L.) *Ko-shigi*.

1765.—*Scolopax Gallinula* L. S. N. XII, I, p. 244. (Europe).

Gallinago gallinula (L.), Uchida.

Jack-Snipe.

Hab. Kurile Isl. Hokkaido, Hondo (Suruga, Hitachi), Formosa, Corea.

Rostratula Vieillot (1816).

276. **Rostratula benghalensis benghalensis** (L.) *Tama-shigi*.

1758.—*Rallus benghalensis* L., S. N. X, I, p. 153. ("Asia")

Rostratula capensis (L.), Uchida.

Painted Snipe.

Hab. Hokkaido, Hondo, Shikoku, Kiusu, Loo-choo Isls., Formosa, Corea.

Hæmatopus L. (1766).

277. **Hæmatopus ostralegus osculans** Swinhoe. *miyakodori*.

1871.—*Hæmatopus osculans* Swinhoe, P. Z. S., p. 405. (China).

Hæmatopus ostralegus (L.), Uchida.

Eastern Oystercatcher.

Hab. Saghalin, Kurile Irls., Hokkaido, Hondo, Shikoku, Tsushima, Formosa, Corea.

278. **Hæmatopus niger bachmani** Audubon *Kuro-miyakodori*.

1838.—*Hæmatopus bachmani* Audubon, B. America IV. Pl. 427, fig. 1 (America).

Hæmatopus niger (nec Pallas), Uchida.

West American Black Oystercatcher.

Hab. Kuril Islands.

Hydrophasianus Wagler (1832).

279. **Hydrophasianus chirurgus** (Scopoli) *Renkaku*.

1786.—*Tringa chirurgus* Scopoli, Del. Flor. et Faun. Insubr.,

II, p. 92.

Luzonian Jacana.

Hab. Formosa.

Oder LARI.

Family LARIDÆ.

Chlidonia Rafinesque (1822).

280. **Chlidonia leucoptera** (Temminck).

..... *Haじro-kuroharu-ajisashi*.

1815.—*Sterna leucoptera* Temminck, Man. d'Orn., p. 483.
(Mediterranean Sea).

White-winged Black Tern.

Hab. Saghalin, Bonin Islands, Corea.

281. **Chlidonia leucopareia swinhonis** Mathews.....

..... *Kurchirra-ajisashi*.

1912.—*Hydrochelidon leucopareia swinhonis* Mathews, B.
Anstr., II, p. 320. (Futschau, China).

Hydrochelidon hybrida (nec Pallas), Uchid*i*.

Chinese Whiskered Tern.

Hab. Formosa.

Gelochelidon Brehm (1831).

282. **Gelochelidon nilotica addenda** Mathews... *Hashibuto-ajisashi*.

1912.—*Gelochelidon nilotica addenda* Mathews, B. Austr. II,
p. 331. (China).

Gelochelidon anglica (nec Moutagu), Uchida & Kuroda, 'Tori'
no. 6. 1918.

Chinese Gull-billed Tern.

Hab. Formosa.

Hydroprogne Kaup (1829).

283. **Hydroprogne tschegrava tschegrava** (Lepechin).
..... *Oni-ajisashi*.
1770.—*Sterna Tschegrava* Lepechin, Novi Comm. Acad. Petropol. XIV, I, p. 500. Pl. XIII. (Caspian Sea in S. Russia).
Hydroprogne caspia (Pall.), Uchida.
Caspian Tern.
Hab. Loo-choo Islands (Yayeyama Group), Formosa.

Thalassers Boie (1822).

284. **Thalasseus bergii cristatus** (Stephens).....
..... *Ô-ajisashi*.
1826.—*Sterna cristata* Stephens, Shaws Gen. Zool. XIII, p. 146. (China).
Sterna bergii Licht., Uchida.
Hab. Marcus Isl., Loo-choo Isl., Formosa.

Sterna L. (1758).

285. **Sterna longipennis** Nordmann.
..... *Ajisashi*.
1835.—*Sterna longipennis* Nordmann, Ermans Verz. Thieren u. Pflanzen, p. 17. (Near Ochotsk).
Nordmann's Tern.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku.
286. **Sterna dougallii bangsi** Mathews
..... *Boni-ajisashi*.
1912.—*Sterna dougallii bangsi* Mathews, B. of Austr. II, p. 364. (Riukiu Islands).
Sterna dougalli gracilis (nec Gould), Uchida.
Hab. Loo-choo Islands.
287. **Sterna aleutica** Baird
..... *Koshijiro-ajisashi*.
1869.—*Sterna aleutica* Baird, Tr. Chicago Acad. Sci. I, p. 321, pl. 31, fig. 1 (Kodiak Island near Alaska).

Aleutian Tern.

Hab. Saghalin, Hondo (Sagami).

Gygisterna Mathew (1912).

288. **Gygis**terna sumatrana sumatrana (Raffles)
..... *Iriguro-ajisashi*.
1821.—*Sterna sumatrana* Raffles, Tr. L. Soc. London XIII.
p. 329. (Sumatra).

Sterna melananchon Temm., Uchida.

Black-naped Tern.

Hab. Loo-choo Isls., Formosa.

Sternula Boie (1822).

289. **Sternula** albifrons sinensis (Gmelin). *Ko-ajisashi*.
1789.—*Sterna sinensis* Gmelin S. N. I. 2., p. 608. (China).
Asiatic Little Tern.
Hab. Hondo, Shikoku, Kiusiu, Loo-choo Isls., Formosa,
Corea.

Onychoprion Wagler (1832).

290. **Onychoprion** fuscatus infuscatus (Lichtenstein)
..... *Seguro-ajisashi*.
1823.—*Sterna infuscata* Lichtenstein, Verz. Dubl. Mus. Berlin, p. 81. (Eastern India).
Sterna fuliginosa (nec Gmelin), Uchida.
Sooty Tern.
Hab. Hondo, Marcus Is., Loo-choo Isls., Formosa.

Melanosterna Blyth (1846).

291. **Melanosterna** anaethetus anaethetus (Scopoli)
..... *Mamijiro-ajisashi*.

1786.—*Sterna Anachetus* Scopoli, Del. Flor. et Faun. Insb.

II, p. 92. (Panay, Philippines).

Bridled Tern, Panayan Tern.

Hab. Hokkaido, Hondo, Marcus Is., Loo-choo Isls., Formosa.

292 **Melanosterna lunata** (Peale) *Nanyō-mamijiro-ajisashi*.

1848.—*Sterna lunata* Peale, U. S. Expl., Birds, p. 277. (Vincennes Is., Paumotu Group).

Hab. Marcus Is.

Procelsterna Lafresnaye (1842).

293. **Procelsterna cærulea** (F. D. Bennett) *Haiiro-ajisashi*.

1840.—*Sterna cærulea* Bennett, Narr. Whaling-Voy., I, p. 385,
and II, p. 248. (Christmas Is., Fanning Group).

Hab. Marcus Is. or Minami-torishima.

Anous Stephens (1826).

294. **Anous stolidus pileatus** (Scopoli) *Kuro-ajisashi*.

1786.—*Sterna pileata* Scopoli, Faun. et Flor. Insubr., II, p.
92. (Philippine Isls.)

1901.—*Anous pullus* Bangs, Bull. Mus. Comp. Harvard,
XXXVI, p. 258. (Riukiu Islands).

Philippine Noddy.

Hab. Hondo, Marcus Is., Loo Choo Isls. Formosa.

Micranous Saunders (1895).

295. **Micranous tenuirostris marcusii** Bryan.... *Torishima-ajisashi*.

1903.—*Micranous marcusii* Bryan, Monogr. Marcus Is., p.
Marcus Island Tern.

Hab. Marcus Island or Minamitorishima.

Gygis Wagler (1832).

296. **Gygis alba kittlitzii** Hartert *Shiro-ajisashi.*
1891.—*Gygis alba kittlitzii* Hartert, Katal. Vogelsamml. Senckenb., p. 237. (Caroline Islands).
Gygis candida Gmelin, Uchida.

White Tern.

Hab. Kurile Isls., Hokkaido, Hondo, Bonin Isls., Sulpher Isls., Marcus Isl., Shikoku, Kiusiu, Loo-choo Isls.

Xema Leach (1819).

297. **Xema sabini** (Sabine) *Kubiza-Kamome.*
1818.—*Larus Sabini* Sabine Tr. L. Soc. London XII, p. 520.
(Near Disko, West Greenland).
Sabine's Gull.
Hab. Hondo (Rikuzen) (Straggler!).

Larus L. (1758).

298. **Larus argentatus vegae** Palmén *Seguro-kamome.*
1887.—*Larus argentatus* Brün. var. *Vegae* Palmén, Vega Exped. Vetensk. Arb. V, p. 370. (Coasts of Siberian Ice Sea).
Larus vegae Palmén, Uchida.
Hab. Hokkaido, Hondo, Bonin Isls., Shikoku, Kiusiu, Loo-choo Isls., Formosa, Corea.
299. **Larus schistisagus** Stejneger *Ō-seguro-kamome.*
1884.—*Larus schistisagus* Stejneger, Auk, p. 201. (Baring Is.)
Slaty-backed Gull.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Formosa.
300. **Larus canus major** Middendorff *Kamome.*
1853.—*Larus canus* L. var. *major* Middendorff, Sibirische Reise, Zool. II, 2, p. 243, Pl. XXIV, fig. 4 (Stanovoi

Range to Ochotsk Sea).

Larus canus (nec L.), Uchida.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Formosa, Corea.

301. **Larus canus brachyrhynchus** Richardson.....*Kō-kamome*.
1832.—*Larus brachyrhynchus* Richardson, Richardson & Swainson's Fauna Bor.—Am. II. p. 422.
Hab. Kuril Isls., Hondo (Sagami).
302. **Larus crassirostris** Vieillot.....*Umi-neko*.
1818.—*Larus crassirostris* Vieillot, Nonv. Dict. d'Hist. Nat. XXI., p. 508. (Nagasaki, Kiusiu).
Black-tailed Gull.
Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Formosa, Corea.
303. **Larus glaucescens** Naumann*Washi-kamome*.
1840.—*Larus glaucescens* Naumann, Vög. Deutschl. X, p. 351. (North America).
Glaucous-winged Gull.
Hab. Hokkaido, Hondo, Corea.
304. **Larus hyperboreus** Gunnerus.....*Shiro-kamome*.
1767.—*Larus hyperboreus* Gunnerus, Leem's, Beskr. Finn. Lapp., pp. 226, 283. (Lapland).
Larus glaucus Fabr., Uchida.
Glaucous Gull.
Hab. Hokkaido, Hondo.
305. **Larus leucopterus** Faber.*Hajiro-kamome*.
1822.—*Larus leucopterus* Faber, Prodromus Isl. Orn., p. 91. (Iceland).
Iceland Gull.
Hab. Hondo.
306. **Larus ridibundus ridibundus** L.*Yuri-kamome*.
1765.—*Larus ridibundus* L., S. N. XII. 1. p. 225. (Europe).

Black-headed-Gull.

Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Formosa, Corea.

307. **Larus ridibundus sibiricus** Buturlin.*Ō-yuri-kamome*.

1911.—*Larus ridibundus sibiricus* Buturlin, Mess. Orn. II, p. 66. (Kolyma Delta and Ussuriland).

Hab. Japan.

308. **Larus saundersi** (Swinhoe).*Zuguro-kamome*.

1871.—*Chroicocephalus saundersi* Swinhoe, P. Z. S. pp. 273, 421, pl. XXII (Near Amoy).

Saunders Gull.

Hab. Saghalin, Hokkaido, Kiusiu, Formosa, Corea.

Rissa Stephens (1826).

309. **Rissa tridactyla pollicaris** Stejneger.*Mitsuyubi-kamome*.

1884.—*Rissa tridactyla pollicaris* Stejneger, Baird, Brewer and Ridgway Water B. N. A. II, p. 202.

Rissa tridactyla (nec L.), Uchida.

Pacific Kittiwake.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku.

Family STERCORARIIDÆ.

Catharacta Brünich (1764).

310. **Catharacta matsudairae** Takatsukasa.*Ō-tōzuku-kamome*.

1922.—*Catharacta matsudairae* Takatsukasa, "Tori," Vol. III, pp. 105, 107 (Sagami Bay).

Catharacta antarctica (nec Lesson), Taka-Tsukasa, "Tori," Vol. II, p. 1.

Hab. Hondo (Uruga and Sagami Bay).

Coprotheres Reichenbach (1850).

311. **Coprotheres pomarinus** (Temminck)*Tōzuku-kamome*.

1815.—*Lestrus pomarinus* Temminck, Man. d'Orn., p. 514.
(Holland and France).

Stercorarius pomarinus (Temminck), Uchida.

Pamatorhine Skua.

Hab. Saghalin, Kurile Isls., Hondo.

Stercorarius Brisson (1760).

312. Stercorarius parasiticus parasiticus (L.)
..... *Kuro-tōzoku-kamome*.

1758.—*Larus parasiticus* L., S. N. X. i. p. 136. (Coast of
Scandinavia).

Stercorarius parasiticus L., Uchida.

Buffon's Skua.

Hab. Saghalin, Kurile Isls, Hondo (Sagami).

313. Stercorarius longicaudus Vieillot.
..... *Shirohara-tōzoku-kamome*.

1819.—*Stercorarius longicaudus* Vieillot, Nouv. Diet. d'Hist.
Nat. (nouv. éd) XXXII, p. 157. (N. Europe, N. Asia and
N. America).

Stercorarius richardsoni (nec Swainson), Uchida.

Arctic Skua.

Hab. Kurile Isls.

Order ALCAE.

Uria Brisson (1760).

314. Uria troile californica (Bryant) *Umi-garasu*.

1861.—*Catarractus californica* Bryant, P. Bost. Soc. N. H.
VIII, p. 142. (Farallone Isls., California).

Alca troile (nec L.), Uchida.

California Murre.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Corea.

315. **Uria lomvia arra** (Pallas.) *Hashibuto-umi-garasu.*

1826.—*Cephus arra* Pallas, Zoogr. Rosso-Asiat. ii. p. 347.

Pallas's Murre.

Hab. Saghalin, Kurile Isls., Hokkaido.

Cephus Pallas (1769).

316. **Cephus carbo** Pallas..... *Keima-furi.*

1827.—*Cephus Carbo* Pallas, Zoogr. Rosso-Asiat. ii. p. 350.
(Kurile Isls.).

Uria carbo (Pallas), Uchida.

Sooty Guillemot.

Hab. Saghalin, Kurile Isls., Hokkaid, Hondo, Corea.

317. **Cephus columba snowi** Stejneger..... *Chishima-umi-bato.*

1897.—*Cephus snowi* Stejneger, Auk, xiv. p. 201. (Kurile Isls.).
Uria snowi (Stejneger) Uchida.

Hab. Kurile Islands, Hokkaido (North end).

318. **Cephus columba columba** Pallas. *Umi-bato.*

1827.—*Cephus Columba* Pallas, Zoogr. Ross-Asiat. ii. p. 348
(Kamtschatka, Bering Strait).

Uria columbia (Pallas), Uchida.

Pigeon Guillemot.

Hab. Saghalin, Kurile Isls., Hokkaido.

Brachyramphus Brandt (1837).

319. **Brachyramphus marmoratus perdix** (Pallas).

..... *Madara-umi-suzumi.*

1827.—*Cephus Perdix* Pallas, Zoogr. Ross-Asiat ii. p. 351, pl.
LXXX (Bering and Ochotsk Seas).

Partridge Auk.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Amami-
Ōshima.

320. **Brachyramphus brevirostris** (Vigors) ... *Kobashi-umi-suzume*.
1828.—*Uria brevirostris* Vigor Zool. Journ. iv. p. 357. (San
Blas, Mexico).
Short-billed Auk.
Hab. Kurile Isls.

Synthliboramphus Brandt (1837).

321. **Synthliboramphus antiquus** (Gmelin) *Umi-suzume*.
1789.—*Alca antiqua* Gmelin, S. N. 1, 2., p. 554. (From W. N.
America to Kamtschatka and Kurile Isls.).
Ancient Auk.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Sulpher Isls.,
Kiusiu, Tsushima, Corea.
322. **Synthliboramphus wumizusume** Temminck.
..... *Kammuri-umi-suzume*.
1835.—*Uria wumizusume* Temminck Al. Col. 579 (Seas of
Japan and Corea.).
Japanese Auk.
Hab. Saghalin, Hokkaido, Hondo, Seven Isls. (Izu), Kiusiu,
Corea.

Ptychoramphus Brandt (1837).

323. **Ptychoramphus aleuticus** (Pallas) *Amerika-umi-suzume*.
1827.—*Uria alcatica* Pallas, Zoogr. Rosso-Asiat. ii. p. 370.
(Eastern parts of Bering Sea).
Hab. Kurile Island.

Phaleris Temminck (1820).

324. **Phaleris psittaculus** (Pallas) *Umi-ōmu*.
1769.—*Alca psittacula* Palls, Special. Zool. i. p. 13, pls. II, V,
fig. 4-6 (Bering Sea).

Parroquet Auk.

Hab. Saghalin, Kurile Islands.

Ciceronia Reichenbach (1852).

325. **Ciceronia pasilla** (Pallas) *Ko-umi-suzume*.

1827.—*Uria pusilla* Pallas Zoogr. Rosso-Asiat. ii. 373. (Kamtschatka).

Simorhynchus pusillus (Pallas), Uchida.

Least Auk.

Hab. Saghalin, Kurile Islands, Hokkaido, Hondo.

Alcella Stone (1907).

326. **Alcella pygmæa** (Gmelin). *Shirahige-umi-suzume*.

1789.—*Alca pygmæa* Gmelin, S. N. I, 2. p. 555. (N. Asia and America).

Simorhynchus pygmæus (Gmelin), Uchida.

Whiskered Auk.

Hab. Saghalin, Kurile Isls., Hondo, Shikoku.

Æthia Merrem (1788).

327. **Æthia cristatella** (Pallas) *Etorofu-umi-suzume*.

1769.—*Alca cristatella* Pallas, Spicil. Zool. v. p. 18, pl. III, (Kuriles).

Simorhynchus cristatellus (Pallas), Uchida.

Crested Auk.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo (Echigo).

Cerorhinca Bonaparte (1828).

328. **Cerorhinca monocerata** (Pallas) *Utō*.

1827.—*Alca monocerata* Pallas, Zoogr. Rosso-Asiat. ii. p.

362. (Coast of Alaska).

Hornbilled Puffin.

Hab. Saghalin, Kurile Isls., Hokkaido, Hndo, Seven Isls.
(Izu), Kiusiu, Tsushima, Corea.

Fratercula Brisson (1760).

- 329. Fratercula corniculata** Naumann *Tsunomedori*.

1821.—*Mormon corniculata* Naumaun Isis. p. 782, pl. 7, fig. 3,
4 (Kamtschatka).

Horned Puffin.

Hab. Saghalin, Kurile Isls., Hando (Awa).

Lunda Pallas (1826).

- 330. Lunda cirrhata** (Pallas) *Etopirika*.

1769.—*Alca cirrhata* Pallass, Spic. Zool. v. p. 7, pl. I, fig. 1-
3. (Between Kamtschatka and America).

Tufted Puffin.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo (Echigo).

Order PTEROCLETES.

Syrrhaptes Illiger (1811).

- 331. Syrrhaptes paradoxus** Pollas *Sakei*.

1773.—*Tetrao paradoxus* Pallas, Reise versch. Prov. Russ.
Reiches, II, p. 712., pl. F. (“Habitat in deserto Tatarico
australiore”).

Pallas' Sand-Grause.

Hab. Corea, Hondo (Sagami ???).

Order COLUMBAE.

Family Columbidae.

Columba L. (1758).

332. ? **Columba livia intermedia** Strickland *Kawarabato*.
1844.—*Columba intermedia* Strickland, Am. & Mag. N. H. p.
39. (India).

Blue Rock Pigeon.

Hab. ? Hondo, ? Loo-choo, ? Formosa.

333. **Columba rupestris rupestris** Pallas *Kōrai-bato*.
1827.—*Columba Oenas ò rupestris* Pallas, Zoogr. Rosso.
Asiat. I. p. 560. ("In Dauria transalpina abundat").
Columba rupestris Bonaparte, Uchida.

Blue Hill-Pigeon.

Hab. Corea.

Alsocomus Tickell (1842).

334. **Alsocomus pulchricollis** (Blyth.) *Taiwan-juzukake-bato*.
1845.—*Columba pulchricollis* Blyth, Jour. As. Soc. Bengal
XIV. p. 866. (Eastern Himalayas).
Ashy Wood-Pigeon.
Hab. Formosa.

Janthoenas Bonaparte (1854).

335. **Janthoenas janthina janthina** (Temminck) *Karasu-bato*.
1830.—*Columba janthina* Temminck, Pl. Col. 503. (Japan).
Carpophaga ianthina (Temm.), Uchida.
Columba janthina janthina Temminck, Hartert.
Japanese Fruit-Pigeon.
Hab. Hondo, Seven Isls. (Izu), Shikoku, Kiusiu, Amami-
Ōshima, Loo-choo, Tsushima.

336. ***Janthoenas janthina jouyi*** Stejneger. ... *Riukiu-karasu-bato*.
 1887.—*Janthoenas jouyi* Stejneger, Am. Naturalist, p. 583.
 (Liu Kiu).
Carpophaga jouyi (Stejn), Uchida.
Columba jouyi (Stejn), Hartert.
 Hab. Loo-choo Isls.
337. ***Janthoenas janthina versicolor* (Kittlitz)**.....
 *Ogasawara-karasu-bato*.
 1832.—*Columba versicolor* Kittlitz, Kupfertaf. i. p. 5, pl. V, fig.
 2 (Bonin Islands).
Carpophaga versicolor (Kittlitz), Uchida.
 Bonin Fruit-Pigeon.
 Hab. Bonin Isls.
338. ***Janthoenas janthina nitens* Stejneger**.....
 *Akagashira-karasu-bato*.
 1887.—*Janthoenas nitens* Stejneger (ex Kittlitz M. S).
 P. U. S. Nat. Mus. X, P. 421 (Bonin Islands).
Carpophaga nitens (Stejn), Uchida.
Columba janthina nitens (Stejn), Hartest.
 Hab. Bonin Isls., Sulpher Isls.

***Streptopelia* Boie (1828).**

339. ***Streptopelia orientalis orientalis* (Latham)** *Kiji-bato*.
 1790.—*Columba orientalis* Latham, Ind. Orn. II. p. 606.
 (China).
 1835.—*Columba gelastis* Temm., Pl. Col. 550. (Japan).
Turtur orientalis (Latham), Uchida.
 Eastern Turtle-dove.
 Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Is. (Izu),
 Shikoku, Kiusiu, Tsushima, Loo-choo Isls., Formosa,
 Botel Tobago, Corea, Quelpart Is.

340. *Streptopelia orientalis stimpsoni* (Stejneger) *Riukiu-kiji-bato*.
 1887.—*Turtur stimpsoni* Stejneger, P. U. S. Nat. Mus. pp. 399, 426 (Riu Kiu Islands).
 Hab. Amami Ōshima. Loo-choo.

341. *Streptopelia decaocto decaocto* (Frivalszky) *Shirako-bato, Juzukake-bato*.
 1838.—*Columba risoria* L. var. *decaocto* Frivalszky, A. M. Társaság Evkönyvei 1834–1836, III. kötet, osztaly 3, pp. 183–184, pl. VIII (Turkestan).
Turtur risorius (nec Temm.). Uchida.
 Indian Ring-dove.
 Hab. Hondo (Saitama, Chiba). Corea.

342. *Streptopelia chinensis chinensis* (Scopoli) *Kanoko-bato*.
 1786.—*Columba chinensis* Scopoli, Del. Flor. & Faun. Insubr., p. 94. (“Tourterelle grise de la Chine”).
Turtur chinensis (Scop.), Uchida.
 Chinese Turtle-dove.
 Hab. Formosa, Botel Tobago Is.

Oenopelia Blanford (1898).

243. *Oenopelia tranquebarica humilis* (Temminck) ... *Beni-bato*.
 1824.—*Columba humilis* Temminck, Pl. Col. 259. (Bengal, Luzon).
Turtur humilis (Temm.). Uchida.
 Dwarf Turtle-dove.
 Hab. Hondo (Saitama, Yokohama), Formosa, Pescadres Isls.

Macropygia Swainson (1837).

344. *Macropygia tenuirostris phoea* McGregor *Onaga-bato*.

1904.—*Macropygia phva* McGregor, Bull. Philipp. Mus. no. 3, p. 9. (Batan Is., Philippines).

Batan Cuckoo-dove.

Hab. Botel Tobago Is. or Kōtōshō.

Chalcoptaphs Gould (1843).

345. **Chalcoptaphs indica** (L.) *Kin-bato*.

1766.—*Columba indica* L., S. N. ed. X, I, p. 284.

Bronze-winged Dove.

Hab. Loo-choo, Formosa.

Sphenurus Swainson (1837).

346. **Sphenurus sieboldii sieboldii** (Temminck)
..... *Azobato, Yamabato*.

1835.—*Columba Sieboldii* Temminck, Pl, Col. 549. (Japan).

Sphenocercus sieboldi (Temm.), Uchida.

Japanese Green Pigeon.

Hab. Hokkaido, Hondo, Shikoku, Kiushiu, Tsushima.

347. **Sphenurus sieboldii sororius** (Swinhoe) *Taiwan-azobato*.
1866.—*Sphenocercus sororius* Swinhoe, Ibis, p. 311.
(Formosa).

Hab. Formosa.

348. **Sphenurus permagnus permagnus** (Stejneger)
..... *Riukiu-azobato*.

1886.—*Treron permagna* Stejneger, Proc. U. S. Nat. Mus. IX.
p. 637. (Okinawa).

Sphenocercus permagnus (Stejneger), Uchida.

Loo-choo Green Pigeon.

Hab. Loo-choo Isls. (Amami-ōshima, Okinawa).

349. **Sphenurus permagnus medioximus** (Bangs) *Chudai-azobato*.

1901.—*Sphenocercus medioximus* (Bangs), Bull. Mus. Comp. Zool., Vol. XXXVI, p. 261 (Ishigaki).

Hab. Loo-cho Isls. (Ishigaki).

350. **Sphenurus formosæ** (Swinhoe) *Taiwan-suaka-awo-bato.*

1863.—*Treron formosae* Swinhoe, Ibis, p. 396. (Formosa).

Sphenocercus formosae (Sw.), Uchida.

Hab. Formosa.

Osmotreron Bonaparte (1854).

351. **Osmotreron bicincta domvillei** Blyth *Muncaka-awo-bato.*

1875.—*Osmotreron domvillei* Blyth, B. Burma, p. 144.

Osmotreron bicincta (nec Jerd.), Uchida.

Eastern Orange-breasted Green Pigeon.

Hab. Formosa. (Straggler).

Order CUCULI.

Family Cuculidae.

Centrops Illiger (1811).

352. **Centropus bengalensis lignator** Swinhoe *Banken.*

1861.—*Cuculus lignator* Swinhoe, Ibis, p. 48. (Formosa and South China).

Centropus javanicus (nec Dum.), Uchida.

Hab. Formosa.

Eudynamys Vigors & Horsfield (1826).

353. **Eudynamis orientalis harterti** Ingram *Oni-kakkō.*

1912.—*Eudynamis orientalis harterti* Ingram, Nov. Zool., Vol. XIX, p. 279. (Hainan).

Eudynamis honorata (nec L.), Kuroda.

Hab. Formosa. (Straggler).

Cuculus L. (1766).

354. **Cuculus canorus telephonus** Heine *Kakkō.*
1863.—*Cuculus telephonus* Heine, Journ. f. Orn. p. 352.
(Japan).
Cuculus canorus (nec L.), Uchida.
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusu, Loo-Choo Isls., Formosa, Corea, Quelpart Is.
355. **Cuculus optatus optatus** Gould *Tsutsudori.*
1845.—*Cuculus optatus* Gould, P. Z. S. p. 18. (Port Essington, Australia).
Cuculus saturatus Hodgs., Uchida.
Himalayan Cuckoo.
Hab. Saghalin, Hokkaido, Hondo, Shikoku, Loo-choo Isls., Corea.
356. **Cuculus optatus kelungensis** Swinhoe *Taiwan-tsutsudori.*
1863.—*Cuculus kelungensis* Swinhoe, Ibis, p. 394. (Kelung, Formosa).
Hab. Formosa.
357. **Cuculus intermedius intermedius** Vahl. *Hototogisu.*
1789.—*Cuculus intermedius* Vahl, Skriv. af Nat. Selskabet. Kjöbenhavn IV, p. 58. ("Tranque-baria," S. India).
Cuculus poliocephalus Lath., Uchida.
Little Cuckoo.
Hab. Hondo, Shikoku, Kiusu, Formosa, Corea.
358. **Cuculus micropterus micropterus** Gould *Suguro-kakkō.*
1837.—*Cuculus micropterus* Gould, P. Z. S., p. 137 (Himalaya).
Hab. Japan (Hokkaido?—Hartert) Corea.

Hierococcyx S. Müller (1839-44).

359. **Hierococcyx fugax nisicolor** (Blyth) *Jiu-ichi.*
1843.—*Cuculus nisicolor* Blyth, Jour. As. Soc. Bengal XII. p. 943. (Nepal).
Hierococcyx fugax (Horsf.), Uchida.
Cuculus fugax nisicolor Blyth, Hartert.
Amoor Cuckoo.
Hab. Hondo, Shikoku, Corea.
360. **Hierococcyx sparverioïdes** (Vigors) *O-jiu-ichi.*
1832.—*Cuculus sparverioides* Vigors, Proc. Committee Zool. Soc. pt. I. p. 173 (Himalaya).
Hab. Japan (Locality unknown).

Order CORACIÆ.

Family Coraciidæ.

Eurystomus Vieillot (1816).

361. **Eurystomus orientalis calonyx** Sharpe *Buppōsō.*
1890.—*Eurystomus calonyx* Sharpe, P. Z. S. p. 551. (Nepal).
Eurystomus orientalis (nec L.), Uchida.
Broad-billed Roller.
Hab. Hondo Kiusiu, Loo-choo Isls., Coaea, Quelpart Island.

Order HALCYONES.

Family Alcedinidæ.

Ceryle Boie (1828).

362. **Ceryle lugubris lugubris** (Temminck)
..... *Yamasemi, Kawachō, Kunokoshōbin.*

1834.—*Alcedo lugubris* Temminck, Nauvean.

Recueil de Planches Coloriees d' Oiseaux, 548 (Japan).

Ceryle guttata (nec Vigor.), Uchida.

Oriental Spotted Kingfisher.

Hab. Hokkaido, Hondo, Kiusiu. ? Corea.

Alcedo L. (1758).

363. **Alcedo atthis japonica** Bonaparte *Kawasemi*.

1754.—*Alcedo japonica* Bp., Conspectus volucr. anis. p. 10 (Japan).

Alcedo ispida bengalensis (nec Gm.), Uchida.

Japanese Common Kingfisher.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Isls, Corea.

364. **Alcedo atthis formosana** Laubmann *Taiwan-kawasemi*.

1920.—*Alcedo atthis atthis formosana* Laubmann.,

Arch. Naturges. Berlin, vol. 84, pt. A, pp. 73, 77. ("Nantu-Distrikt, Formosa").

Formosan Common Kingfisher.

Hab. Formosa.

Halcyon Swainson (1820).

365. **Halcyon smyrnensis fusca** (Boddaert) *Azashōbin*.

1783.—*Alcedo fusca* Boddaert, Table des Planches enlumineez d'Histoire Naturelle de M. D'Aubenton, p. 54 ("Martin-pêcheur de la côte de Malabar").

White-breasted Kingfisher.

Hab. Formosa.

366. **Halcyon miyakoensis** Kuroda *Miyakoshōbin*.

1919.—*Halcyon miyakoensis* Kuroda, "Dōbutsugaku Zasshi," Vol. XXI, pp. 229, 231. (Miyakojima).

Hab. Loo-Choo Isls. (Miyakojima).

367. **Halcyon pileata** (Boddaert) *Yamashobin.*
 1783.—*Alcedo pileata* Boddaert, Table des Planches enluminées d'Histoire Naturelle de M. D'Aub. p. 41. (China).
 Black-capped Kingfisher.
 Hab. Hondo, Corea.

Entomothera Horsfield (1820).

368. **Entomothera coromanda major** (Temminck and Schlegel).
 *Akashobin, Miyamashobin.*
 1842.—*Halcyon coromanda major* Temminck and Schlegel,
 Fanna Japonica, Aves, p. 75 (Japan).
Halcyon coromanda (nec Lath), Uchida.
Halcyon coromanda major T. & S., Hartert.
 Japanese Ruddy Kingfisher.
 Hab. Hokkaido, Hondo, Shikoku, Kiushiu, Corea.
369. **Entomothera coromanda bangsi** Oberholser
 *Rinkin-akashobin.*
 1915.—*Entomothera coromanda bangsi* Oberholser, P. U. S.
 Nat. Mus. Vol. 48, p. 654 (Ishigaki).
Halcyon coromanda rufa (nec Wall.), Uchida.
 Hab. Loo-choo Isls., Formosa.

Order MEROPES.

Family Meropidae.

Cosmaerops Cabanis and Heine (1860).

370. **Cosmaerops ornatus** (Latham.) *Hachi-kui.*
 1801.—*Merops ornatus* Latham, Ind. Orn. Suppl. p. XXXV.
 (Australia).
 Australian Bee-eater.
 Hab. Loo-choo Islands (Straggler) (Ogawa).

Order UPUPÆ.

Family Upupidae.

Upupa L. (1758).

371. **Upupa epops saturata** Lönnberg..... *Yatsugashira*.
1909.—*Upupa epops saturata* Lönnberg, Arkiv. f. Zool. V, p.
29 (Kjachta).
Upupa epops (nec L.), Uchida.
Hab. Hokkaido, Hondo, Kiusiu, Tsushima, Corea.

Order STRIGES.

Family Strigidæ.

Nyctea Stephens (1826).

372. **Nyctea nyctea** (L.) *Shiro-fukuro*.
1758.—*Strix Nyctea* L., Syst. Nat. X, I, p. 93 (Sweden).
Snowy Owl.
Hab. Saghalin, Hokkaido, Hondo (Chiba, Mino), Corea.

Bubo Duméril (1806).

373. **Bubo bubo tenuipes** Clark *Washi-mimizuku*.
1907.—*Bubo tenuipes* Clark, P. U. S. Nat. Mus. XXXII, p.
470 (Fusan, Corea).
Bubo maximus (nec Flem.), Uchida.
Hab. Kiusiu, (Goto Is), South Islands of Satsuma, Corea.
374. **Bubo blakistoni blakistoni** Seeböhm *Shima-fukuro*.
1883.—*Bubo blakistoni* Seeböhm, P. Z. S., p. 466 (Hokkaido,
Japan).
Blakiston's Eagle-Owl.
Hab. Saghalin, Hokkaido.
375. **Bubo blakistoni doerriesi** Seeböhm *Chōsen-shimafukuro*.

1895.—*Bubo doerrisi* Seebold, Bull. B. O. C. V. p. IV.

(Sidemi).

Hab. Corea. (N. E. boundary).

Ketupa Lesson (1831).

376. **Ketupa flavipes** (Hodgson)..... *Uwo-mimizuku.*

1836.—*Cultrungins flavipes* Hodgson, Jour. As. Soc. Bengal,

V. p. 364.

Hab. Formosa.

Otus Pennant (1769).

377. **Otus bakkamoena semitorques** Temminck and Schleger.....
..... *ō-konohazuku.*

1850.—*Otus semitorques* Temminck and Schleger, Fauna Japonica, Aves, p. 25, pl. 8 (Japan).

Scops semitorques (Temm. and Schl.), Uchida.

Feathered-toed Scops Owl.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Quelpart Isl.

378. **Otus bakkamoena ussuricensis** (Buturlin)
..... *Sameiro-ō-konohazuku.*

1910.—*Scops semitorques ussuricensis* Buturlin, Orn. Mitteilungen, p. 119 (Ussuriland).

Hab. Corea.

379. **Otus bakkamoena pryeri** (Gurney) *Riukiu-ōkonohazuku.*

1889.—*Scops pryeri* Gurney, Ibis, p. 302 (Riu-kuu Isls.).

Scops semitorques pryeri Gurn., Uchida.

Pryer's Scops Owl.

Hab. Loo-choo Isls.

380. **Otus bakkamoena glabripes** (Swinhoe). *Taiwan-ōkonohazuku.*

1870.—*Ephialtes glabripes* Swinhoe, Ann. and Mag.

Nat. Hist. ser 4, VI, p. 152 ("South China and Formosa").

Hab. Formosa.

381. **Otus japonicus japonicus** Temminck and Schleger. *Konohazuku.*

1850.—*Otus scops japonicus* Temminck and Schleger, Faun. Jap. Aves, p. 27, pl. 9 (Japan).

Scops japonicus (Temm. and Schleg.), Uchida.

Otus japonicus Temm. and Schleg., Hartert.

Japanese Scops Owl.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

382. **Otus japonicus elegans** (Cassin) *Riukiu-konohazuku.*

1852.—*Ephialtes elegans* Cassin, P. Acad. Phil. p. 185 (Riukiu Isls.).

Scops japonicus elegans (Cass.), Uchida.

Cassin's Scops Owl.

Hab. Amami-Ōshima, Loo-choo Isls., Formosa?

383. **Otus hambroecki** (Swinhoe) *Taiwan-konohazuku.*

1870.—*Ephialtes hambroecki* Swinhoe, Ann. Mag. Nat. Hist. (4) vi. p. 153 (Formosa).

Scops hambroecki (Sw.), Uchida.

Hab. Formosa.

Asio Schaeffer (1779).

384. **Asio otus otus** (L.) *Torafuzuku.*

1758.—*Strix Otus* L., S. N., X, I, p. 92 (Sweden).

Asio otus L., Uchida.

Long-eared Owl.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea.

385. **Asio flammeus flammeus** Pontoppidan. *Komimizuku.*

1763.—*Strix flammea* Pontoppidan, Danske Atlas I. p. 617.
(Denmark).

Asio accipitrinus (Pall.), Uchida.

Short-eared Owl.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu,
Formosa, Corea.

386. **Asio flammeus leucopsis** (Brehm). *Samiro-komimizuku*.

1855.—*Otus leucopsis* Brehm, Vogelfang, p. 413. (“In
Osteuropa bei Sarepta”).

Hab. Japan (Hartert).

Ninox Hodgson (1837).

387. **Ninox scutulata scutulata** (Raffles). *Azobazuku*.

1822.—*Strix scutulata* Raffles, Transact. L. Soc. XIII. p. 280.
(Sumatra).

Ninox scutulata (Raffles), Uchida.

Brown Owlet.

Hab. Hondo, Shikoku, Kiusiu, Loo-choo Isls., Formosa,
Corea.

388. **Ninox scutulata ussuriensis** Buturlin. *Chōsen-azobazuku*.

1910.—*Ninox scutulata ussuriensis* Buturlin, Orn. Mitt. (Mos-
kau) p. 187. (Ussuriland and Corea).

Hab. Corea (Buturlin).

Aegolius Kaup (1829).

389. **Aegolius tengmalmi sibiricus** (Buturlin). *Kinme-fukuro*.

1910.—*Cryptoglaux sibirica* Buturlin, Nasha Okhota, Juni-
Nom., p. 78 (Lake Khanka).

Nyctale tengmalmi (nec Gml.), Uchida.

Hab. Saghalin, Kurile Isls., Hokkaido.

Glaucidium Boie (1826).

390. **Glaucidium passerinum orientale** Taczanowski.
..... *Suzume-fukurō*.
1891.—*Glaucidium passerinum orientale* Taczanowski, Faune.
Orn. Sib. Orient. I, p. 128 (E. Siberia).
Glaucidium passerinum (nec L.), Murata, Report on the
Fauna of the Saghalin Island.
Hab. Saghalin Is.
391. **Glaucidium pardalotum** (Swinhoe).
..... *Hime-fukurō*.
1863.—*Athene pardalota* Swinhoe, Ibis p. 216 (Formosa).
Hab. Formosa.

Surnia Duméril (1806).

392. **Surnia ulula pallasi** Buturlin.
..... *Onaga-fukurō*.
1907.—*Surnia ulula pallasi* Buturlin, Orn. Montber, p. 100.
(Siberia).
Surnia ulula (nec L.), Murata, Rept. on Faun. Saghalin.
Hab. Saghalin, Kurile Isls.

Strix L. (1758).

393. **Strix aluco nivicola** (Blyth).
..... *Chōsenfukurō*.
1845.—*Syrnium nivicolum* Blyth, Journ. As. Soc. Bengal,
XIV, p. 185. ("Himalaya").
1907.—? *Syrnium ma* Clark, Proc. U. S. Nat. Mus., XXXII,
p. 471. ("Fusan, Korea").
Syrnium nivicola (nec Hodggs.), Uchida, (Pt.).
Hab. Corea.
394. **Strix aluco harterti** (La Touche).
..... *Takasago-fukurō*.
1920.—*Syrnium aluco harterti* La Touche, Bull. B. O. C. Vol.
XL, p. 50 (Formosa).

-
- Syrnium nivicola* (nec Hodg.), Uchida (pt.).
Hab. Formosa.
- 395. *Strix nebulosa sakhalinensis* (Butuslin). *Karafuto-fukuro*.**
1907.—*Syrnium cinereum sakhalinense* Buturlin, Journ. f. Ornith., p. 332 (Sakhalin Is.).
Syrnium lapponicum (nec Thunb.), Uchida.
Hab. Sakhalin.
- 396. *Strix uralensis nikolskii* (Buturlin). *Shiboriya-fukuro*.**
1907.—*Syrnium uralense nikolskii* Buturlin, Journ. f. Ornith. p. 333. (South-eastern Siberia).
Hab. Sakhalin, ? Corea.
- 397. *Strix uralensis japonica* (Clark). *Yezo-fukuro*.**
1907.—*Syrnium uralense japonicum* Clark, P. U. S. Nat. Mus., XXXII, p. 471. (Hokkaido).
Syrnium uralensis (nec Pall.), Uchida (pt.).
Hab. Hokkaido, Hondo.
- 398. *Strix uralensis hondoensis* (Clark). *Fukuro*.**
1907.—*Syrnium uralense hondoense* Clark, P. U. S. Nat. Mus. XXXII, p. 472 (Iwaki, Hondo).
Syrnium uralensis (nec Pall.), Uchida (pt.).
Hondo Ural Owl.
Hab. Hondo.
- 399. *Strix uralensis fuscescens* Temminck and Schleger.**
1847.—*Strix fuscescens* Temminck and Schleger, Faun. Jap. Aves, p. 30, pl. 10 (Japan).
Syrnium uralensis fuscescens (Temm. and Schleg.), Uchida.
Hab. Awaji, Shikoku (?), Kiusiu.
- 400. *Strix newarensis caligatus* (Swinhoe). *ō-fukuro*.**
1863.—*Bubo caligatus* Swinhoe, Ibis, p. 218. (Formosa).

Syrnium indrance (nec Sykes), Uchida.

Hab. Formosa.

Tyto Billberg (1828).

401. **Tyto longimembris** Jerdon. *Men-fukurō.*

1893.—*Strix longimembris* Jerdon, Madras Journ. L. Soc. X.
p. 86.

Strix candida Tickell, Uchida.

Hab. Formosa.

Order CAPRIMULGI.

Family Caprimulgidae.

Caprimulgus L. (1758.).

402. **Caprimulgus indicus** jotaka Temminck and Schlegel).

..... *Yotaka, Kasuidori.*
1847.—*Caprimulgus jotaka* Temminck and Schleger, Faun.
Jap., Aves., p. 37, pl. 12 (Japan).

Japanese Goatsucker.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Corea.

403. **Caprimulgus monticola** Franklin.

..... *Taiwan-yotaka, Taiwan-kasuidori.*

1831.—*Caprimulgus monticola* Franklin, P. Z. S., p. 116.

1863.—*Caprimulgus stictomus* Sw., Ibis, p. 250.

Hab. Formosa.

Order CYPSELI.

Family Cypselidae.

Apus Scopoli (1777.).

404. **Apus pacificus** (Latham). *Ama-tsubame.*

1801.—*Hirundo pasifica* Latham, Ind. Orn. Suppl. p. LVIII.
(Australia).

Cypselus pacificus (Lath.), Uchida.

White-rumped Swift.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku,
Formosa, Corea, Quelpart Island.

405. *Apus affinis subfurcatus* (Blyth). *Hime-ama-tsubame*.

1849.—*Cypselus subfurcatus* Blyth, Jour. As. Soc. Bengal
XVIII. p. 807 (Penang).

Hab. Formosa.

***Hirundapus* Hodgson (1836.).**

406. *Hirundaps caudacutus caudacutus* (Latham). *Hariwo-ama-tsubame*.

1801.—*Hirundo caudacuta* Latham, Ind. Orn. Suppl. p. LVII
(Australia).

Chaetura caudacuta (Lath.), Uchida.

Needle-tailed Swift.

Hab. Saghalin, Kurile Islands, Hokkaido, Hondo, Corea.

407. *Hirundapus caudacutus nudipes* Hodgson. *Kurobitac-hariwo-ama-tsubame*.

1836.—*Chaetura nudipes* Hodgson, Jour. As. Soc. Bengal. V.
p. 779 (Nepal).

Hab. Formosa.

Order SCANSORES.

Family Capitonidæ.

***Cyanops* Bonaparte (1854.).**

408. *Cyanops nuchalis* (Gould). *Goshiki-dori*.

1862.—*Megalaema nichalis* Gould, P. Z. S., p. 283 (Formosa).
Hab. Formosa.

Order PICI.

Family Picidae.

Picus L. (1758.).

409. **Picus canus jessoensis** Stejneger. *Yamagera*.
1886.—*Picus canus jessoensis* Stejneger, P. U. S. Nat. Mus.,
IX, p. 106 (Jesso=Hokkaido).
Gecinus canus (nec Gm.), Uchida.
Japanese Grey-headed Green Woodpecker.
Hab. Hokkaido.
410. **Picus canus griseoviridis** (Clark). *Chōsen-yamagera*.
1907.—*Gecinus canus griseoviridis* Clark, P. U. S. Nat. Mus.,
XXXII, p. 473 (“Seoul, Korea”).
Hab. Corea.
411. **Picus canus tancolo** (Gould) *Taiwan-yamagera*.
1862.—*Gecinus tancolo* Gould, P. Z. S. p. 283 (Formosa).
Hab. Formosa.
412. **Picus awokera awokera** Temminck. *Atwogera*.
1826.—*Picus awokera* Temminck, Nouveau Recul de Planches
colorieés d’Oiseaux, p. 585 (Japan).
Gecinus awokera (Temminck), Uchida.
Japanese Green Woodpecker.
Hab. Hondo.
413. **Picus awokera horii** Taka-Tsukasa *Kagoshima-atwogera*.
1918.—*Picus awokera horii* Taka-Tsukasa, “Dōbutsugaku
Zasshi,” Vol. XXX, No. 361, p. 442 (Kagoshima).
Hab. Shikoku (Tosa), Kiusiu (Prefect. Kagoshima, Buzen,
Chikuzen).

- 414.** *Picus awokera takatsukasae* Kuroda. *Fa.-awo-gera.*
1921.—*Picus awokera takatsukasae* Kuroda, Auk, XXXVIII,
p. 576 (Tanegashima).
Hab. Tanegashima.

Sapheopipo Hargitt (1890).

- 415.** *Sapheopipo noguchii* (Seebold). *Noguchi-gera.*
1887.—*Picus noguchii* Seebold, Ibis, p. 178 (Okinawa).
Dryobates noguchii (Seeb.), Uchida.
Pryer's Woodpecker.
Hab. Loo-choo Isls. (Okinawa).

Dryobates Boie (1826).

- 416.** *Dryobates major tscherskii* (Buturlin). *Kita-akagra.*
1910.—*Dendrocopos japonicus tscherskii* Buturlin, Nascha
Okhota, July, p. 53 (Lake Khanka).
Dryobates major major (nec L.), Uchida.
Hab. Saghalin, ? Kurile Isls. (Shikotan).

- 417.** *Dryobates major japonicus* (Seebold). *Yezo-akagra.*
1883.—*Picus japonicus* Seebold, Ibis, p. 24. (Hokkaido).
Japanese Great Spotted Woodpecker.
Hab. ? Kurile Isls., Hokkaido, Hondo (Northern parts), ? Corea.

- 418.** *Dryobates major hondoensis* Kuroda. *Akagra.*
1921.—*Dryobates major hondoensis* Kuroda, Auk, XXXVIII,
p. 577 (Shinano, C. Hondo).
Hab. Hondo (Northern and Central parts), ? Corea.

- 419.** *Dryobates leucotos uralensis* (Malherbe). *Chōsen-akagra.*
1861.—*Picus uralensis* Malh., Monogr. Picidés, I, p. 92, pl.
23, fig. 4, 5 (Ural Mountain).
1907.—*Dryobates leucotos corcensis* Clark, Proc. U. S. Nat.

- Mus., XXXII, p. 472 (Fusan, Corea).
- Dryobates leucotos ussuricus* (Buturlin), Uchida.
- Dryobates leucotos leucotos* (nec Bechstein), Uchida (pt.).
- Ural White-backed Woodpecker.
- Hab. Sakhalin, Corea.
420. ***Dryobates leucotos subciris*** Stejneger. *Yezo-ōakagera*.
 1886.—*Dryobates subciris* Stejneger, Proc. U. S. Nat. Mus., IX, p. 113. (Sapporo, Yesso).
Dryobates leucotos leucotos (nec Bechstein), Uchida (pt.).
 Hab. Hokkaido.
421. ***Dryobates leucotos stejnegeri*** Kuroda. *Ō-akagera*.
 1921.—*Dryobates leucotis stejnegeri* (error) Kuroda, Auk, XXXVIII, p. 579 (Shinano, C. Hondo).
Dryobates leucotos subciris (nec Stejn.), Uchida.
 Japanese White-backed Woodpecker.
 Hab. Hondo (Northern and Central parts).
422. ***Dryobates leucotos takahashii*** Kuroda and Mori
 *Shima-ōakagera*.
 1920.—*Dryobates leucotos takahashii* Kuroda and Mori,
 "Tori," Vol. II, No. 10, pp. 266, 277 (Dagelet Is., Sea
 of Japan).
 Hab. Utsuriotō or Dagelet Is.
423. ***Dryobates leucotos intermedius*** Kuroda. ... *Kansai-ōakagera*.
 1913.—"An intermediate form between *Picus leuconotus sub-*
ciris (Stejneger) and *Picus namiyai* (Stejneger)," Kuroda,
 "Dōbutsugaku Zasshi," p. 333 (Near Drogawa, Prov,
 Yamato, Hondo).
 1921.—*Dryobates leucotis intermedius* Kuroda, Auk,
 XXXVIII, p. 580 (Near Drogawa, Prov. Yamato, Hondo).
 Hab. Hondo (Pacific side of Hondo: Provinces Sagami,
 Suruga, Yamashiro, Yamato, Kii), Shikoku (Iyo).

424. **Dryobates leucotos quelpartensis** Kuroda and Mori.
 Saisiu-ōakagera.
 1918.—*Dryobates leucotos quelpartensis* Kuroda and Mori,
 “Tori,” Vol. II, No. 7, pp. 74, 88 (Quelpart Island).
 Hab. Quelpart Is. or Saisiuto.
425. **Dryobates leucotos insularis** (Gould). ... *Taiwan-ōakagera.*
 1862.—*Picus insularis* Gould, P. Z. S., p. 283 (Formosa).
 Hab. Formosa.
426. **Dryobates lencotos namiyei** Stejneger. ... *Namiye-gera.*
 1886.—*Dryobates namiyei* Stejneger, P. U. S. Nat. Mus., IX,
 p. 16, pl. II (Prov. Yamato, S. western Hondo).
 Stejneger's Woodpecker.
 Hab. Hondo (South-western portions), Shikoku, Kiusiu.
427. **Dryobates leucotos owstoni** (Ogawa). *Owston-gera.*
 1905.—*Picus owstoni* Ogawa, Annot. Z. Jap., V., p. 203, pl. X
 (Amami-ōshima).
 Hab. Amami-ōshima.
428. **Dryobates minor minutillus** (Buturlin). *Ko-akagera.*
 1908.—*Xylocopus minor minutillus* Buturlin, Annuaire Mus.
 Zool. Petersbourg, XIII, p. 246 (Sidemi).
 Dryobates minor (nec L.), Uchida.
 Hab. Saghalin, Hokkaido.

Yungipicus Bonaparte (1854.).

429. **Yungipicus kizuki seebohmi** Hargitt. *Yezo-kogera.*
 1884.—*Lynxipicus seebohmi* Hargitt, Ibis, p. 100 (“Hab. in
 insulis Japonicis ‘Niphon’ et ‘Yezo’ dictis.”—Designated
 by Dr. Stejneger as “Yesso” = Hokkaido),
 Lynxipicus kizuki (nec Temm.), Uchida (pt.).
 Dryobates kizuki seebohmi (Harg.), Hartert (pt.).

Hargitt's Pigmy Woodpecker.

Hab. Sakhalin, Kurile Isls., Hokkaido, ? N. Hondo.

430. ***Yungipicus kizuki nippōn*** Kuroda. *Kogera*.

1922.—*Yungipicus kizuki nippōn* Kuroda. Ibis, p. 88 (Gotemba, Suruga, Hondo).

Yungipicus kisuki (nec Temm.), Uchida (pt.).

Dryobates kizuki sechowmii (nec Harg.), Hartert (pt.).

Hondo Pigmy Woodpecker.

Hab. Hondo (Northern and Central parts), Corea, Quelpart Is.

431. ***Yungipicus kizuki kizuki*** Temminck. *Kiusiu-kogera*.

1836.—*Picus Kizuki* Temminck, Nouveau Recueil de Planches Colorieés d'Oiseaux, IV, livr. 99 (text only) the text erroneously inscribed *Picus atewokera* (Island of Kiusiu).

Yungipicus kisuki Temm., Uchida (pt.).

Dryobates kizuki kizuki (Temminck), Hartert.

Temminck's Pigmy Woodpecker.

Hab. ? Hondo (Southern parts), ? Shikoku, Kiusiu (Northern and ? Southern parts), Yakushima.

432. ***Yungipicus kizuki kotataki*** Kuroda. *Tsushima-kogera*.

1922.—*Yungipicus kizuki kotataki* Kuroda, Ibis, p. 86 (Tsushima).

Hab. Tsushima.

433. ***Yungipicus kizuki matsudairai*** Kuroda. *Miyake-kogera*.

1921.—*Yungipicus kizuki matsudairai* Kuroda, Auk, XXXVIII, p. 576 (Miyakeshima, Seven Isls. of Izu).

Hab. Seven Isls. of Izu (Miyakeshima).

434. ***Yungipicus kizuki amamii*** Kuroda. *Amami-kogera*.

- 1922.—*Yungipicus kizuki amamii* Kuroda, Ibis, p. 88 (Amami-ōshima).
- Hab. Amami-ōshima.
- 435. *Yungipicus kizuki nigrescens* Seeböhm. *Riu-kiu-kogera.***
- 1887.—*Iyngipicus kizuki nigrescens* Seeböhm, Ibis, p. 177. (Naha, Okinawa, Loo-choo group).
- Loo-choo Pigmy Woodpecker.
- Hab. Loo-choo Islands (Okinawa).
- 436. *Yungipicus pygmaeus kaleensis* (Swinhoe) ... *Taiwan-kogera.***
- 1863.—*Picus kaleensis* Swinhoe Ibis, p. 390 (Formosa).
Iyngipicus pygmaeus kaleensis (Swinhoe), Uchida.
Dryobates pygmaeus kaleensis (Swinhoe), Hartert.
- Hab. Formosa.
- 437. *Yungipicus pygmaeus doerriesi* (Hargitt).... *Imuru-kogera.***
- 1881.—*Iyngipicus doerriesi* Hargitt, Ibis, p. 398 (Island of Askold near Vladivostok).
Dryobates pygmaeus doerriesi (Hargitt), Hartert.
- Hab. Corea.
- 438. *Yungipicus wattersi* Salvadori and Giglioli. *Kurohoshi-kogera.***
- 1885.—*Iyngipicus wattersi* Salvadori and Giglioli, Atti della R. Accad. Sc. Tor., XX, p. 825 (Formosa).
- Hab. Formosa.

Picoides Lacépède (1799).

- 439. *Picoides tridactylus sakhalinensis* Buturlin. *Miyubigera.***
- 1907.—*Picoides sakhalinensis* Buturlin, Ornith. Monatb, pp. 10, 11 (Sakhalin and Amur region).
- 1908.—*Picoides (tridactylus) sachalinensis* Lönnberg, J. Coll. Sc. Tokyo, XXIII, Art. 14, p. 41 (Sakhalin Is.).
Picoides tridactylus, (nec L.), Uchida.
- Hab. Saghalin.

Dryocopus Boie (1826).

440. **Dryocopus martius silvifragus** Riley. *Kumagera*.
1915.—*Dryocopus martius silvifragus* Riley, Proc. Biol. Soc.
Wash., XXVIII, p. 162. (Hokkaido).
Picus martius (nec L.), Uchida.
Dryocopus martius martius (nec L.), Hartert (pt.).
Japanese Great Black Woodpecker.
Hab. Saghalin, Hokkaiko.
441. **Dryocopus martius morii** Kuroda *Chōsen-kumagera*.
1921.—*Dryoscopus* (error) *martius morii* Kuroda, Auk,
XXXVIII, p. 575 (Central Corea).
Hab. Corea.

Thripotax Cabanis and Heine (1863).

442. **Triponax richardsi** (Tristram) *Kitataki, Amanojakuma*.
1879.—*Dryocopus richardsi* Tristram, P. Z. S., p. 386, pl.
XXXI ♀ (Tsushima).
1887.—*Thripotax kalinowskii* Taczanowski, P. Z. S., p. 607
(Corea).
Picus richardsi (Tristram), Uchida.
Tristram's Woodpecker.
Hab. Tsushima, Corea.

Jynx L. (1758).

443. **Jynx torquilla japonica** Bonaparte *Arisui*.
1850.—*Jynx japonica* Bonaparte, Conspl. Gen. Av. 1, p. 112
("Ex Japonica").
Jynx torquilla (nec L.), Uchida.
Japanese Wryneck.
Hab. Saghalin, Hondo, Shikoku, Kiusiu.

444. **Jynx torquilla hokkaidi** Kuroda *Yezo-arisui.*
1921.—*Jynx torquilla hokkaidi* Kuroda, Auk, XXXVIII, p.
582 (Prov. Kitami, Hokkaido).
Hab. Hokkaido.

Order PASSERES.

Family PITTIIDÆ.

Pitta Vieillot (1816).

445. **Pitta nymphæ** Temminck and Schleger. *Yairochō, Akadannu.*
1850.—*Pitta nymphæ* Temminck and Schleger, Faun. Jap.,
Aves, Suppl. t. A.
1864.—*Pitta orcas* Swinhoe, Ibis, p. 428 (Formosa).
Fairy Pitta.
Hab. Hondo (Shinano, Fuji-yama, Oki Is.), Kiushū (Buzen),
Tsushima, Formosa, Corea, Quelpart Isl.

Family ALAUDIDÆ.

Alauda L. (1758).

446. **Alauda arvensis pekinensis** Swinhoe *Ō-hibari.*
1863.—*Alauda pekinensis* Swinhoe, Proc. Zool. Soc. Lond., p.
69 (Peking).
1884.—*Alauda blakistoni* Stejneger, Proc. Biol. Soc. Washing-
ton, II, p. 98 (Kamtschatka).
Blakiston's Skylark.
Hab. Saghalin, Kurile Is., Hokkaido, Hondo, Shikoku, Corea,
Quelpart Is.
447. **Alauda arvensis intermedia** Swinhoe *Chiu-hibari.*
1863.—*Alauda intermedia* Swinhoe, Proc. Zool. Soc. Lond.,
p. 89 (Shanghai).

- Hab. Hondo, Kiusiu, Corea, Quelpart Is.
- 448. *Alauda arvensis japonica* Temminck and Schlegel ... *Hibari*.**
 1848.—*Alauda japonica* Temminck and Schlegel, Siebold's
 Fauna Japon., Aves, p. 87, pl. 47 (Japan).
 Japanese Skylark.
 Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Is.,
 Shikoku, Kiusiu, Tanegashima, Yakushima, Amami-
 Ōshima.
- 449. *Alauda arvensis dulcivox* Brooks *Ko-hibari*.**
 1873.—*Alauda dulcivox* Brooks, Stray Feathers, I, p. 484
 (Alpine Region of North India ?).
 Alauda arvensis cinerea Ehrencke, Hartert.
 Eastern Skylark.
 Hab. Corea.
- 450. *Alauda gulgula coelivox* Swinhoe *Taiwan-ko-hibari*.**
 1850.—*Alauda coelivox* Swinhoe, Zoologist, p. 6724; Ibis,
 1860, pp. 52, 132 (Amoy).
 Hab. Formosa.
- 451. *Alauda gulgula sala* Swinhoe *Taiwan-hibari*.**
 1870.—*Alauda sala* Swinhoe, Ibis, p. 355 (Hainan).
 1871.—*Alauda wattersi* Swinhoe, Proc. Z. S. London, p. 389
 (S. Formosa and Pescadores).
 Hab. Formosa, Pescadores Is.
- Galerida Boie (1828).**
- 452. *Galerida cristata coreensis* Taczanowski ... *Kammuri-hibari*.**
 1887.—*Galerita cristata coreensis* Taczanowski, Proc. Zool.
 Soc. Lond., p. 603 (Seoul in Corea).
 Corydus cristatus (nec Linn.), Takatsukasa, Uchida's Brd.
 Jap. II, p. 210.

Corean Crested-lark.

Hab. Corea.

Chionophilos Brehm (1832).

453. **Chionophilos alpestris euroa** (Thayer and Bangs).....

..... *Ilama-hibari*.

1914.—*Otocorvus alpestris euroa* Thayer and Bangs, Proc. N.

Engl. Zoöl. Club, V, p. 43 (Kolyma, Eastern Siberia).

Otocorvus alpestris (nec Linn.), Uchida.

Hab. Saghalin, Kurile Is., Hokkaido, Hondo (Shinano-straggler!), Tsushima (straggler!).

Family MOTACILLIDÆ.

Budytes Cuvier (1817).

454. **Budytes flavus taivanus** Swinhoe *Tsumenaga-sekirei*.

1863.—*Budytes taivanus* Swinhoe, Proc. Zool. Soc., p. 334
(Formosa).

Motacilla flava taivanus (Swinhoe), Hartert.

Eastern Yellow-Wagtail.

Hab. Saghalin, Kurile Is., Hondo (Suruga, Musashi), Loo-choo Isls. (Okinawa), Formosa.

455. **Budytes flavus simillimus** (Hartert).

..... *Mamijiro-tsumenaga-sekirei*.

1905.—*Motacilla flava simillima* Hartert, Vog. pal., I, p. 289
(Kamtschatka, Corea, China, etc.).

Eastern Blue-headed Wagtail.

Hab. N. Kurile Is. (Paramushir), Corea.

456. **Budytes flavus macronyx** Stresemann.....

..... *Kita-tsumenaga-sekirei*.

1920.—*Budytes fluvius macronyx* Stresemann, Avif. Macedon., p. 76 (Wladivostok).

Motacilla flava borealis (nec Sund.), Hartert (pt.)
Stresemann's Grey-headed Wagtail.

Hab. Corea.

Calobates Kaup (1829).

457. **Calobates cinereus melanope** (Pallas). *Kisckirei*.

1776.—*Motacilla Melanope* Pallas, Reise d. versch. Prov. Russ. Reichs, III, p. 696 (Dauria).

Motacilla boarula melanope Pall., Hartert.
Eastern Grey Wagtail.

Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Yaku-shima, Tanegashima, Amami-Ōshima, Kikaigashima, Okinawashima, Corea, Dagelet Is., Tsushima, Quelpart Is., Formosa, Pescadores Is., Botel Tobago or Kōtōshō.

Motacilla L. (1758).

458. **Motacilla alba leucopsis** Gould *Hōjirō-sekirei*.

1837.—*Motacilla leucopsis* Gould, Proc. Zool. Soc. Lond., p. 78 (India).

White-faced Wagtail.

Hab. Corea, Dagelet Is., Tsushima, Formosa, Pescadores Is., Botel Tobago or Kōtōshō.

459. **Motacilla alba lugens** Kittlitz *Haku-sekirei*.

1833.—*Motacilla lugens* Kittlitz, Kupfertafeln zur Naturg. d. Vög., 2, p. 16, pl. XXI, fig. 1 (Kamtschatka).

Kamtschatkan Wagtail, Japanese Pied-Wagtail.

Hab. Saghalin, Kurile Is., Hokkaido, Hondo, Seven Is. (Ōshima), Shikoku, Yakushima, Tanegashima, Amami-Ōshima, Tsushima, Corea, Quelpart Is., Ishigakishima,

Pescadores Is. (Taisho).

460. **Motacilla alba ocularis** Swinhoe *Taiwan-huku-sekirei*.

1860.—*Motacilla ocularis* Swinhoe, Ibis, p. 55 (Amoy).

Streak-eyed Wagtail.

Hab. Corea, Formosa.

461. **Motacilla alba grandis** Sharpe. *Seguro-sekirei*.

1885.—*Motacilla grandis* Sharpe, Cat. B. Brit. Mus. X, p. 492

(Japan).

Motacilla alba japonica (nec Sw.), Uchida.

Japanese Wagtail.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, S. Corea. (S. Keishō Distr. and S. Chūsei Distr.).

Dendronanthus Blyth (1844).

462. **Dendronanthus indicus** (Gmelin) *Iwanti-sekirei*.

1788.—*Motacilla indica* Gmelin, Syst. Nat. I, p. 662 (India).

Forest Wagtail or Indian Wagtail.

Hab. Hondo (Suruga-straggler), Bonin Is. (straggler), Corea.

Anthus Bechstein (1807).

463. **Anthus trivialis hodgsoni** Richmond *Binzui, Kihiburi*.

1864.—*Anthus maculatus* Jerdon, Brd. India, iii, p. 873 (India)

Preoccupied.

1907.—*Anthus hodgsoni* Richmond, Carnegie Inst. Washington, Publication, No. 54, p. 493 (China).

Anthus trivialis maculatus Jerdon, Hartert.

Anthus maculatus Hodgson, Uchida.

Eastern Tree-pipit.

Hab. Saghalin, Kirule Is., Hokkaido, Hondo, Shikoku, Kiusiu, Tusu-hima, Corea, Ishigakishima.

464. **Anthus trivialis yunnanensis** Uchida and Kuroda *Kobashi-binzui, Kobashi-kihiburi*.

1916.—*Anthus maculatus yunnanensis* Uchida and Kuroda,
Annot. Zoot. Jap. ix, p. 134 (Yunnan and Formosa).

Anthus maculatus brevirostris (nec Hodgson), Uchida
and Kuroda.

Short-billed Tree-pipit.

Hab. Formosa, Pescadores Is. (Taisho).

465. ***Anthus spinoletta borealis* Hesse** *Karafuto-tahibari*.

1915.—*Anthus barvalis* Hesse, Journ. f. Orn., p. 386 (Sakhalin).

1916.—*Anthus spinoletta reuteri* Munsterhjelm, Nyt. Mag.

Naturw. Kristiania, p. 165 (Sakhalin).

Hab. Sakhalin Is.

466. ***Anthus spinoletta japonicus* Temminck and Schlegel.** *Tahibari*.

1847.—*Anthus spratensis japonicus* Temminck and Schlegel,
Siebold's Fauna Japon., Aves, p. 59, pl. 24 (Japan).

Japanese Water-Pipit, Japanese Alpine-Pipit.

Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Tanegashima, Tsushima, Corea, Quelpart Is.

467. ***Anthus cervinus* (Pallas)** *Muneaka-tahibari*.

1827.—*Motacilla Cervina* Pallas, Zoogr. Rosso-Asiat, I, p.
511 (Siberia).

Red-throated Pipit.

Hab. Saghalin, Kurile Is., Corea, Quelpart Is., South Islands
of Satsuma, Amami-Ōshima, Loo-choo Isls., Formosa.

468. ***Anthus roseatus* Blyth** *Chōsen-tahibari*.

1847.—*Anthus roseatus* "Hodgson," Blyth, Journ. Asiat. Soc.
Bengal, XVI, p. 437 (Nepal).

Anthus rosaceus Hodgs., Takatsukasa, Uchida's Brd. Jap.
II, p. 211.

Hodgson's Pipit.

Hab. Corea.

- 469.** *Anthus gustavi* Swinhoe *Sejiro-tahiburi*.
 1863.—*Anthus gustavi* Swinhoe, Proc. Zool. Soc. Lond., p. 90
 (Amoy).
 Petchora Pipit.
 Hab. Corea.
- 470.** *Anthus richardi richardi* Vieillot *Momijiro-tahiburi*.
 1818.—*Inthus Richardi* Vieillot, Nouv. Dict. d'Hist. Nat.
 XXVI, p. 491 (France).
Anthus richardi Vieill., Uchida.
 Richard's Pipit.
 Hab. Formosa.
- 471.** *Anthus richardi striolatus* Blyth *Ko-momijiro-tahiburi*.
 1847.—*Inthus striolatus* Blyth, Journ. Asiatic. Soc. Bengal,
 XVI, p. 435 (Darjeeling).
 Blyth's Pipit.
 Hab. Corea.

Family BRACHYPODID.E.

Microscelis Gray (1840).

- 472.** *Microscelis amaurotis amaurotis* (Temminck) *Hiyodori*.
 1830.—*Turdus amaurotis* Temminck, Pl. Col. II, pl. 497
 (Japan).
Hypsipetes amaurotis amaurotis (Temm.), Hartert.
 Brown-eared Bulbul.
 Hab. Honshū, Shikoku, Kiusiu, Tanegashima. Yakushima,
 Tsushima, Corea, Quelpart Is., Seven Isls.
- 473.** *Microscelis amaurotis hensoni* (Stejneger) *Ezo-hiyodori*.
 1893.—*Hypsipetes amaurotis hensoni* Stejneger, Proc. U. S.
 Nat. Mus. XV, p. 347 (Hakodate, Jesso).
 Hab. Hokkaido, Hondo, Tsushima, Loo-choo Isls, Corea.
- 474.** *Microscelis amaurotis pryeri* (Stejneger) ... *Riukiu-hiyodori*.

- 1887.—*Hypsipetes pryeri* Stejneger, Proc. U. S. Nat. Mus. IX,
 " 1886," p. 642 (Okinawa).
Hypsipetes amaurotis pryeri Stejn., Hartert.
 Loo-choo Bulbul.
 Hab. Amami-Ōshima, Okinawashima, Iheyashima, Jamamishima.
- 475. Microscelis amaurotis stejnegeri (Hartert) .Ishigaki-hiyodori.**
 1907.—*Hypsipetes amaurotis stejnegeri* Hartert, Vög. pal., I,
 p. 464 (Ishigaki).
 Hab. S. Loo-choo Isls. (Ishigakishima, Iriomoteshima, Miyakoshima).
- 476. Microscelis amaurotis harterti KurodaTaiwan-hiyodori.**
 1922.—*Microscelis amaurotis harterti* Kuroda, Annot. Zool.
 Japon., Vol. X, Art. 4, p. 116 (Botel Tobago).
 Hab. Formosa (Kōshun Forests, Botel Tobago or Kōtōshō,
 Kwashōtō, Kizantō).
- 477. Microscelis amaurotis ogawae (Hartert)Amami-hiyodori.**
 1907.—*Hypsipetes amaurotis ogawae* Hartert, Vög. pal., I, p.
 465 (Amami-Ōshima).
 Hab. Amami-Ōshima.
- 478. Microscelis amaurotis squamiceps (Kittlitz)**
 *Ogasawara-hiyodori.*
 1830.—*Oriolus squamiceps* Kittlitz, Mém. Acad. Imp. St.
 Petersb., I, p. 241, pl. XVI (Bonin Islands).
Hypsipetes amaurotis squamiceps (Kittl.), Hartert.
 Bonin-island Bulbul.
 Hab. Bonin Islands.
- 479. Microscelis amaurotis magnirostris (Hartert)**
 *Hashibuto-hiyodori.*
 1905.—*Hypsipetes amaurotis magnirostris* Hartert, Bull. B. O.
 C. XV, p. 46 (S. Dionisio, Volcano Isls.).

Sulphur Island Bulbul.

Hab. Sulphur Isls. or Volcano Isls.

Haringtonia Mathews and Iredale (1917).

480. **Haringtonia psaroides nigerrimus** (Gould) ... *Kuro-hijodori*.
1862.—*Hyipsipetes nigerrimus* Gould, P. Z. S., p. 282 (Formosa).

Hab. Formosa.

Pycnonotus Boie (1826).

481. **Pycnonotus sinensis formosae** Harttert *Shirogashira*.
1910.—*Pycnonotus sinensis formosae* Harttert, Nov. Zool.,
XVII, pp. 229–230 (Formosa).

Hab. Formosa.

482. **Pycnonotus sinensis taivanus** Styan *Kurogashira*.
1893.—*Pycnonotus taivanus* Styan, Ibis, p. 470 (Formosa).
Hab. Formosa.

Spizixus Blyth (1845).

483. **Spizixus cinereicapillus** Swinh. *Kayanobori*.
1871.—*Spizixus cinereicapillus* Swinhoe, Proc. Zool. Soc.
London, p. 370 (Formosa).
Hab. Formosa.

Apalopteron Bonaparte (1854).

484. **Apalopteron familiare** (Kittlitz) *Meguro*.
1830.—*Ixos familiaris* Kittlitz, Mém. Acad. St. Pétersb. des
Savans Etrang. I, p. 235, pl. XIII (Bonin Islands).
Apalopteron familiare (Kittl.), Uchida.

Hab. Bonin Islands.

Family MUSCICAPIDÆ.

Terpsiphone Gloger (1827).

485. **Terpsiphone atrocaudata atrocaudata** Eyton *Sankō-chō*.

- 1835.—*Muscipeta princeps* Temm., Pl. Col. III, pl. 584 (North Japan and Corea). (Preoccupied!).
- 1839.—*Muscipeta atrocandata* Eyton, P. Z. S., p. 102 (Japan).
- 1847.—*Muscipeta principalis* Temm. and Schl., Faun. Jap., Aves, p. 47, pl. 17 E. (Japan).
- 1910.—*Terpsiphone orwestoni* "Jony," Stejneger, Proc. U. S. Nat. Mus., XXXVII, p. 654 (Fujiyama, Hondo).
Terpsiphone princeps princeps (Temm.), Uchida.
Terpsiphone princeps atrocaudata (Eyton), Uchida.
Tchitrea princeps princeps (Temm.), Hartert.
- Japanese Paradise Flycatcher.
Hab. Hondo, Kiusiu, Tsushima, Corea, Quelpart Is.
- 486. *Terpsiphone atrocaudata illex* Bangs *Riukiu-sankō-chō*.**
1901.—*Terpsiphone illex* Bangs, Bull. Mus. Comp. Zoöl. Harvard XXXVI, p. 264 (Ishigaki).
Tchitrea princeps illex (Bangs), Hartert.
Hab. Loo-choo Isls., Formosa.
- 487. *Terpsiphone incei* (Gould) *Kawari-sankō-chō*.**
1852.—*Muscipeta Incei* Gould, B. Asia II, pl. 19 (Shanghai).
Tchitrea incei (Gould), Hartert.
Hab. Corea.
- 488. *Terpsiphone nigra* McGregor *Kuro-sankō-chō*.**
1907.—*Terpsiphone nigra* McGregor, Phil. Journ. Sci. 2, sec. A, p. 340.
Hab. Kōtōshō or Botel Tobago.

Hemichelidon Hodgson (1845).

- 489. *Hemichelidon sibirica sibirica* (Gmelin) *Same-bitaki*.**
1788.—*Muscicapa sibirica* Gmelin, Syst. Nat. I, p. 939 (Eastern Siberia).

Muscicapa sibirica sibirica Gm., Hartert.

Siberian Flycatcher.

Hab. Sakhalin, Hokkaido, Hondo, Shikoku, Formosa, Corea.

490. **Hemichelidon griseisticta griseisticta** Swinhoe ... *Yizobitaki*.

1861.—*Hemichelidon griseisticta* Swinhoe, Ibis, p. 330
(Amoy and Taku).

Muscicapa griseisticta (Sw.), Hartert.

Hab. Saghalin, Kurile Is., Formosa, Corea.

491. **Hemichelidon griseisticta habereri** (Parrot) ... *Chishima-bitaki*.

1907.—*Muscicapa griseisticta habereri* Parrot, Ornith. Monatsber. Berlin, XV, pp. 168—170. (Itrop, Kurile Isls.).

Hab. Kurile Islands (Itrop).

492. **Hemichelidon cinereiceps** Sharpe *Miyamahitaki*.

1887.—*Hemichelidon cinereiceps* Sharpe, Ibis, p. 441 (Kina Balu).

1845.—*Hemichelidon ferruginea* Hodgson, Proc. Zool. Soc. Lond., p. 32. (Preoccupied).

Muscicapa ferruginea (Hodgs.), Hartert.

Hab. Formosa.

Alseonax Cabanis (1850).

493. **Alseonax latirostris** Raffles) *Kosame-bitaki*.

1821.—*Muscicapa latirostris* Raffles, Trans. L. Soc. Lond. XIII, 2, p. 312 (Sumatra).

Brown Flycatcher.

Hab. Saghalin, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Is, Formosa, Corea, Saisiutō Is. or Quelpart Is.

Muscicapula Blyth (1843).

494. **Muscicapula hypertyhra** (Blyth) ... *Muneakahitaki*.

1842.—*Muscicapa hypertyhra* Blyth, Journ. As. Soc. Beng.,

XI, p. 885.

Hab. Formosa.

Siphia Hodgson (1837).

495. **Siphia parva albicilla** (Pallas) *Ojirohitaki*.

1827.—*Muscicapa Albicilla* Pallas, Zoogr. Rosso-Asiat. I, P. 462 (Daurian Region).

Muscicapa parva albicilla Pall., Hartert.

Hab. Corea.

Cyornis Blyth (1842).

496. **Cyornis vivida** Swinhoe *Chabara-ōruri*.

1864.—*Cyornis vivida* Swinhoe, Ibis, p. 363 (Formosa).

Muscicapa vivida (Sw.), Uchida.

Hab. Formosa.

Hypothymis Boie (1826).

497. **Hypothymis azurea oberholseri** Stresemann. *Kuroycrihitaki*.

1913.—*Hypothymis azurea oberholseri* Stresemann, Nov. Zool, XX, p. 295 (Formosa).

Hypothymis azurea (nec Bodd.), Uchida.

Hab. Formosa.

Zanthopygia Blyth (1847).

498. **Zanthopygia narcissina narcissina** Temminck. *Kibitaki*.

1835.—*Muscicapa narcissina* Temm., Pl. Col. 577, fig. 1 (Japan).

Muscicapa narcissina narcissina Temm., Hartert.

Narcissus Flycatcher.

Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu.

499. **Zanthopygia narcissina jakuschima** (Hartert) *Yakushima-kibitaki*.

- 1907.—*Muscicapa narcissina jakuschima* Hartert, Vög. pal. Faun., I, p. 491 (Yakushima).
 Hab. Yakushima, Amami-Ōshima ?, Okinawa-Shima ?.
- 500. *Zanthopygia narcissina owstoni* (Bangs) ... *Kiukiu-kibitaki*.**
 1901.—*Zanthopygia owstoni* Bangs, Bull. Mus. Comp. Zoöl. Harvard, XXXVI, p. 265 (Ishigaki).
Muscicapa narcissina owstoni (Bangs), Hartert.
 Hab. S.-Loochoo Is. (Ishigaki, Iriomote).
- 501. *Zanthopygia narcissina zanthopygia* (Hay). *Mamijiro-kibitaki*.**
 1845.—*Muscicapa Zanthopygia* Hay, Madras Journal XIII, 2, p. 162 (Malacca).
Muscicapa narcissina zanthopygia Hay, Hartert.
Muscicapa narcissina tricolor Hartlaub, Uchida.
 Hab. Hondo (Shinano-straggler !), Corea, Saishiuto or Quelpart Is.

Poliomyias Sharpe (1897).

- 502. *Poliomyias mugimaki* (Temminck) *Mugimaki*.**
 1835.—*Muscicapa Mugimaki* Temm., Pl. Col. 577, fig. 2 (Japan).
 Mugimaki Flycatcher.
 Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Formasa, Corea, Saisiuto or Quelpart Is.

Cyanoptila Blyth (1847).

- 503. *Cyanoptila cyanomelana* (Temminck) *Oruri*.**
 1828.—*Muscicapa cyanomelana* Temm. Pl. Col. 470 (Japan-♂).
 1847.—*Muscicapa gularis* Temm. and Schl., Faun. Jap. Aves, p. 43, pl. XVI (Japan-♀).
Muscicapa cyanomelana Temm., Uchida.
 Japanese Blue Flycatcher.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Formosa, Corea, Saisintō or Quelpart Is.

Abrornis Hodgson (1844).

504. **Abrornis albogularis fulvifacies** Swinhoc *Koshijirohitaki*.
1870.—*Abrornis fulvifacies* Sw., Proc. Zool. Soc. Lond. p.
132 (Szechwan, China).
1912.—*Abrornis albogularis formosana* Laubmann, Ornith.
Monatsber Berlin, XX, p. 174 (Formosa—juv.!).
Cryptolopha fulvifacies (Sw.), Uchida.
Hab. Formosa.

Oreocinchla Gould (1837).

505. **Oreocinchla dauma aurea** Holandre *Tora-tsugumi*.
1825.—*Turdus aureus* Hol., Faune dép. Moselle, p. 60 (Metz).
Turdus dauma aureus Hol., Hartert.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Is.,
Formosa, Corea, Bōkotō=Pescadores.
506. **Oreocinchla dauma amami** (Hartert) *Ôtora-tsugumi*.
1922.—*Turdus dauma amami* Hartert, Vög. pal. Faun., III,
p. 2155. Nom. emend.
1905.—*Geocichla major* Ogawa, Ann. Zool. Jap. V. p. 178
(Amami-ōshima). (Preoccupied).
Turdus dauma major (Ogawa), Uchida.
Hab. Amami-ōshima.

Cichloselys Bonaparte (1854).

507. **Cichloselys sibiricus davisoni** (Hume) *Mamijiro*.
1877.—*Turdus davisoni* Hume, Stray Feathers, V. p. 63
(Tenasserim, Birmah).
Turdus sibiricus davisoni (Hume), Hartert.

Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Corea.

508. **Cichloselys sibiricus sibiricus** (Pallas) *Siberia-mamijiro.*

1776.—*Turdus sibiricus* Pall., Reise d. versch. Prov. Ruso.
Reichs III, p. 694 (Dauria).

Turdus sibiricus sibiricus Pall., Hartert.

Hab. Corea.

Merula Leach (1816).

509. **Merula cardis** (Temminck) *Kuro-tsugumi.*

1830.—*Turdus cardis* Temm., Pl. Col. 518 (Japan).

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Corea.

510. **Merula eunomus** (Temminck) *Tsugumi.*

1830.—*Turdus eunomus* Temminck Pl. Col. 514 (Japan).

Turdus fuscatus Pallas, Hartert.

Hab. Saghalin Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Is., Formosa, Corea, Saisiutō or Quelpart Is.

511. **Merula naumanni** (Temminck) *Hachijō-tsugumi.*

1820.—*Turdus naumannii* Temminck, Man. d' Orn. I, p. 170
(".....en Silésie et en Autriche ;.....").

Hab. Hokkaido, Hondo, Shikoku, Tsushima, Loo-choo Is., Formosa, Corea, Saisiutō or Quelpart Is.

512. **Merula ruficollis atrogularis** (Temminck). *Nodoguro-tsugumi.*

1820.—*Turdus atrogularis* Temminck, Man. d'Orn. I, p. 169
("Habitat : rarement en Autriche et en Silésie.....").

Turdus atrigularis Temm., Uchida "Tori," III, p. 9, 1921.

Hab. Yayeyama Isls. (Ishigaki Is.-straggler !)

513. **Merula hortulorum** (Scalder) *Kara-akahara.*

1863.—*Turdus hortulorum* Scaler, Ibis, p. 196 (Amoy, S. China).

Hab. Hondo, Corea.

514. **Merula obscura** (Gmelin) *Mamichajinai.*

- 1789.—*Turdus obscurus* Gmelin, Syst. Nat. I, 2, p. 816 (Lake Baikal).
- Hab. Hokkaido, Hondo, Kiushiu, Loo-choo Is., Formosa, Corea, Saisiutō=Quelpart Is.
- 515. *Merula pallida* (Gmelin)** *Shirohara*.
- 1789.—*Turdus pallidus* Gmelin, Syst. Nat. I, 815 (Lake Baikal).
- Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Is., Formosa, Corea, Saisiutō or Quelpart Is.
- 516. *Merula chrysolaus* (Temminck)** *Akahara*.
- 1831.—*Turdus chrysolaus* Temminck, Pl. Col. 537 (Japan).
- Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo Is., Formosa, Corea, Bōkotō=Pescadores.
- 517. *Merula celaenops celaenops* (Stejneger)** *Akakokko*.
- 1887.—*Turdus celaenops* Stejn., Science, X, p. 108 (Miyakeshima, Seven Isls., Izu.)
- Hab. Hondo (straggler), Seven Islands of Izu.
- 518. *Merula celaenops yakushimensis* Ogawa.....**
- *Yakushima-akakokko*.
- 1905.—*Merula celaenops yakushimensis* Ogawa, Ann. Zool, Jap. V, p. 180 (Yakushima).
- Turdus celaenops yakushimensis* (Ogawa), Hartert.
- Hab. Yakushima.
- 519. *Merula albiceps* (Swinhoe)** *Taiwan-tsugumi*.
- 1864.—*Turdus albiceps* Swinhoe, Ibis, p. 363 (Formosa).
- Hab. Formosa.

Ægithocichla Sharpe (1903).

- 520. *Ægithocichla terrestris* (Kittlitz)** *Ogasawara-gabichō*.
- 1830.—*Turdus terrestris*, Kittl. Mem. Acad. St. Pétersb. I, p. 245, Pl. XVII (Bonin-Islands).
- Hab. Bonin Islands. (Extinct species).

Monticola Boie (1822).

521. **Monticola solitaria magna** (La Touche) *Iso-hiyodori*.
1920.—*Petrophila solitaria magna* La Touche, Bull. B. O. C.,
XL, p. 97 (Japan).
Hab. Kurile Is., Hokkaido, Hondo, Shikoku, Kiushu, Loo-
choo Is., Bonin Is., Corea, Saishutō=Quelpart Is.
522. **Monticola solitaria latouchei** Kuroda *Tsushima-isohyo*.
1922.—*Monticola solitaria latouchei* Kuroda, Ibis, p. 92 (Tsu-
shima).
Hab. Tsushima and Iki Islands.
523. **Monticola solitaria philippensis** (P. L. S. Müller)
1776.—*Turdus philippensis* P. L. S. Müll., Natursystem, An-
hang, p. 142.
Monticola solitarius (P. L. S. Müll.), Uchida.
Hab. Formosa.
524. **Monticola gularis** (Swinhoe) *Hime-isohyo*.
1862.—*Oreocetes gularis* Sw., Proc. Zool. Soc. Lond. p. 318,
Ibis, 1863, p. 93, pl. III (Peking).
Hab. Corea.

Myiophonus Temminck (1823).

525. **Myiophonus insularis** Gould *Kurichō*.
1862.—*Myiophonus insularis* Gould, P. Z. S., p. 280 (Formosa).
Hab. Formosa.

Saxicola Bechstein (1802).

526. **Saxicola torquata stejnegeri** (Parrot) *Nobitaki*.
1908.—*Pratincola rubicola stejnegeri* Parrot, Verh. orn. Ges.
Bayern VIII, p. 124 (Etrop and Hokkaido).
Pratincola maura (nec Pall.), Uchida.
Hab. Saghalin, Kurile Is., Hondo, Kiusiu, Loo-choo Is., For-

mosa, Corea.

Ianthia Blyth (1847).

527. **Ianthia cyanura** (Pallas) *Ruribitaki.*
1773.—*Motacilla Cyanurus* Pall., Reise d. versch. Prov. Russ.
Reichs II, p. 709 (Yenesei).
Tersiger cyanurus (Pall.), Hartert.
Siberian Blue-tail.
Hab. Saghalin, Kurile Is., Hokkaido, Hondo, Shikoku, Kiusiu,
Loo-choo Is., Formosa, Corea, Saisiutō=Quelpart Is.
528. **Ianthia johnstoniae** Ogilvie-Grant *Arisanhitaki.*
1906.—*Ianthia johnstoniae* O.-Grant, Bull. B. O. C., XVI, p.
118 (Formosa).
Hab. Formosa.
529. **Ianthia goodfellowi** Ogilvie-Grant *Kikuchihitaki.*
1912.—*Ianthia goodfellowi* O.-Grant, Ibis p. 649, (Formosa).
Hab. Formosa.
530. **Ianthia indica formosana** (Hartert) *Takasago-hitaki.*
1909.—*Tarsiger indica formosanus* Hartert, Bull. B. O. C.,
XXV, p. 32 (Formosa).
Hab. Formosa.

Chaimarronis Hodgson (1844).

531. **Chaimarronis fuliginosus affinis** (Ogilvie-Grant) *Kawa-bitaki.*
1906.—*Xanthopygia affinis* O.-Grant, Bull. B. O. C., XVI, p.
118 (Formosa).
Chaimarrornis fuliginosa affinis (Grant), Hartert.
Hab. Formosa.

Phoenicurus Forster (1817).

532. **Phoenicurus auroreus auroreus** (Pallas) *Jō-bitaki.*
1776.—*Motacilla aurorea* Pall., Reise d. versch. Prov. d.

Russ. Reichs III, 695 ("In salicetis circa Selengam et collaterales fluvios, usque ad Sinarum fines").

Ruticilla aurorea (Gm.), Uchida.

Hab. Kurile Isls. Hokkaido, Hondo, Kiusiu, Formosa, Corea,
Saisiutō=Quelpart Is.

- 533. *Phoenicurus ochruros rufiventris* (Vieillot) ...*Kuro-jō-bitaki*.**
1818.—*Oenanthe rufiventris* Vieill., Nouv. Dict. d' Hist. Nat.
Nouv. Ed., XXI. p. 431 (India).

Hab. Corea.

Cyano-sylvia Brehm (1828).

- 534. *Cyano-sylvia svecica robusta* (Buturlin) ...*Ogawa-komadori*.**
1907.—*Cyanecula svecica robusta* But., Psovedia i Rusheinaia
Okhota, March, no. 6 and Orn. Monatsber. May, p. 79
(Kolyma Delta).
Erithacus cyaneculus caeruleoculus (nec Pallas), Kuroda,
Zool. Mag. XXVIII. p. 508.
Hab. Hondo (Suruga-straggler !), Corea.

Erithacus Cuvier (1801).

- 535. *Erithacus akahige* (Temminck)*Komadori*.**
1824.—*Sylvia akahige* Temm., Pl. Col. 571 ("Riu-kuu Isls.",
errore).
Luscinia akahige (Temm.), Hartert.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Seven
Isls. of Idzu, Yakushima.

- 536. *Erithacus sibilans* (Swinhoe)*Shimagoma*.**
1863.—*Larvivora sibilans* Sw., Proc. Zool. Soc. Lond. p. 292
(Near Macao in China).

Luscinia sibilans (Sw.), Hartert.

Hab. Saghalin, Hondo ?, Corea.

Calliope Gould (1837).

537. **Calliope calliope calliope** (Pallas) *Nogoma.*
1776.—*Motacilla Calliope* Pall., Reise d. versch. Prov. Russ.
Reichs III, 697 (“..... Jenisea usque ad Lenam”).
Luscinia calliope (Pall.), Hartert (pt.).
Erithacus calliope (Pall.), Uchida.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusu, Formosa, Bōkotō or Pescadores, Corea.

Larvivora Hodgson (1837).

538. **Larvivora cyanea** (Pallas) *Koruri.*
1776.—*Motacilla Cyane* Pall., Reise d. versch. Prov. Russ.
Reichs III, p. 697 (Dauria).
Luscinia cyane (Pall.), Hartert.
Erithacus cyaneus (Pall.), Uchida.

Hab. Saghalin, Hokkaido, Hondo, Corea.

Icterus Stejneger (1887).

539. **Icturus komadori komadori** (Temminck) *Akahige.*
1824.—*Sylvia komadori* Temm. Pl. Col. 570 (“Korea,”
errore).
Luscinia komadori komadori (Temm.), Hartert.
Erithacus komadori (Temm.), Uchida.
Hab. Yayeyama Islands, Amami-Ōshima, Tokunoshima.
540. **Icturus komadori namiyei** Stejneger *Hontō-akahige.*
1886.—*Icturus namiyei* Stejn., Proc. U. S. Nat. Mus. IX. p.
645 (Okinawa).
Luscinia komadori namiyei (Stejn), Hartert.
Erithacus komadori namiyei (Stejn.), Uchida.
Hab. Loo-choo (Okinawa).

Brachypteryx Horsfield (1820).

541. **Brachypteryx goodfellowi** Ogilvie-Grant.....
..... *Kobane-chimedori, Kikuchi-chimedori.*
1912.—*Brachypteryx goodfellowi* Grant, Bull. B. O. C. XXIX,
p. 108 (Formosa).
Brachypteryx cruralis (nec Hodgson), Kuwada.
Hab. Formosa.

Notodera Lesson (1827).

542. **Notodera leucura montium** (Swinhoe) *Kon-hitaki.*
1864.—*Myiomela montium* Swinh., Ibis, p. 362 (Formosa).
Hab. Formosa.

Microcichla Sharpe (1883).

543. **Microcichla scouleri fortis** Hartert *Shirokuro-hitaki.*
1910.—*Microcichla scouleri fortis* Hartert, Vög. pal., I, p.
761. (Formosa).
Microcichla scouleri (nec Vigor.), Uchida.
Hab. Formosa.

Acanthopneuste Blasius (1858).

544. **Acanthopneuste tenellipes** (Swinhoe) *Ezo-mushikui.*
1860.—*Phylloscopus tenellipes* Sw., Ibis p. 53 (Near Amoy).
Pale-legged Willow-Warbler.
Hab. Saghalin, Hokkaido, Hondo (Shinano, Tokyo), Kiusiu
(Chikuzen) Corea.
545. **Acanthopneuste nitidus plumbeitarsus** (Swinhoe)
..... *Yanagi-mushikui.*
1861.—*Phylloscopus plumbeitarsus* Swinhoe, Ibis, p. 330 (Near
Peking).

Phylloscopus nitidus plumbicatarsus Sw., Hartert.

Hab. Corea.

546. **Acanthopneuste borealis borealis** (Blasius) *Komushikui.*

1858.—*Phyllopeuste borealis* Blas., Naumannia, p. 313 (Near Ochotsk Sea).

Phylloscopus borealis borealis (Blas.), Hartert.

Phylloscopus borealis (Blas.), Uchida.

Arctic Willow-Warbler.

Hab. Kurile Isls., Hokkaido, Hondo, Kiusiu, Formosa, Corea, Saisiutō or Quelpart Is.

547. **Acanthopneuste borealis xanthodryas** (Swinhoe)..... *Meboso.*

1863.—*Phylloscopus xanthodryas* Sw., P. Z. S., p. 296 (Amoy).

Swinhoe's Willow-Warbler.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Formosa, Corea, Saisiutō or Quelpart Is.

548. **Acanthopneuste occipitalis coronata** (Temminck and Schlegel *Sendai-mushikui.*

1847.—*Ficedula coronata* T. and S., Fauna Jap., Aves, p. 48, pl. 18 (Japan).

Phylloscopus occipitalis coronata (T. and S.), Hartert.

Phylloscopus coronatus (T.), Uchida.

Temminck's Crowned Willow-Warbler.

Hab. Hokkaido, Hondo, Shikoku, Formosa, Corea, Saisiutō = Quelpart Is.

549. **Acanthopneuste ijimæ Stejneger** *Ijima-miboso.*

1892.—*Acanthopneuste ijimæ* Stejneger, Proc. U. S. Nat.

Mus. XV, p. 372 (Seven Is.)

Phylloscopus ijimae (Stejn.), Hartert.

Hab. Seven Islands of Idzu.

Reguloides Blyth (1847).

550. **Reguloides proregulus proregulus** (Pallas).....
.....*Karafuto-mushikui*.
1827.—*Motacilla Proregulus* Pall., Zoogr. Rosso-Asiat. I, p. 499 (Dauria).
Phylloscopus proregulus proregulus (Pall.), Hartert.
Phylloscopus proregulus Pall., Uchida.
Hab. Saghalin, Corea.
551. **Reguloides superciliatus superciliatus** (Gmelin).....
.....*Kimayu-mushikui*.
1788.—*Motacilla superciliata* Gm., Syst. Nat. I, p. 975 (Russia).
Phylloscopus superciliatus superciliatus (Gm.), Hartert.
Phylloscopus superciliatus (Gm.), Uchida.
Hab. Formosa, Corea.

Herbivocula Swinhoe (1871).

552. **Herbivocula schwarzii** (Radde).....*Karafuto-mujisekka*.
1863.—*Sylvia (Phyllopcnuste) schwarzii* Radde, Reise Süden V. Ost-Sib. II, p. 260, pl. IX (Tarei-Nor and Bureja Mountains).
Lusciniola schwarzii (Radde), Uchida.
Radde's Bush-Warbler.
Hab. Sakhalin, Corea.
553. **Herbivocula fuscata** (Blyth)*Mujisekka*.
1842.—*Phyllopcnuste fuscata* Blyth, Journ. As. Soc. Bengal. XI, p. 113 (Near Calcutta).
Lusciniola fuscata (Blyth), Uchida.
Phylloscopus fuscata (Blyth), Hartert.
Hab. Formosa.

Horornis Hodgson (1845).

554. **Horornis cantans cantans** (Temm. and Schleg.) *Uguisu*.
1847.—*Salicaria cantans* T. and S., Fauna Jap., Aves, p. 51,
pl. 19 (Japan).
Horeites cantans cantans (T. and S.), Hartert.
Horeites cantans (T. and S.), Uchida.
Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Loo-
choo, Formosa, Saisiutō.
555. **Horornis cantans ijimae** Kuroda *Ijima-uguisu*.
1922.—*Horornis cantans ijimae* Kuroda, Annot. Zool. Japan.
Vol. X, Art. 4, p. 117 (Miyakeshima).
Hab. Seven Islands of Izu.
556. **Horornis cantans canturians** (Swinhoe) *Taiwan-uguisu*.
1860.—*Arundinax canturians* Sw., Ibis, p. 52 (Amoy and
Shanghai).
Arundinax minutus Sw., Ibis, 1860, p. 52 (Amoy).
Horeites cantans canturians (Sw.), Hartert.
Hab. Formosa.
557. **Horornis cantans borealis** (Campbell) *Chōsen-uguisu*.
1892.—*Cettia minuta borealis* Campb., Ibis p. 235 (Chemul-
po in Corea).
Horeites cantans borealis (Campb.), Hartert.
Hab. Corea, Formosa (Hartert).
558. **Horornis cantans diphone** (Kittlitz) *Ogasawara-uguisu*.
1830.—*Sylvia diphone* kittl., Mem. Acad. Imp. Sc. Petersb. p.
237 and pl. (Bonin Isls.).
Horeites cantans diphone (Kittl.), Hartert.
Hab. Ogasawara-shima or Bonin Isls., Volcano Isls. ?
559. ? **Horornis flavolivacea** (Blyth) *Kibara-uguisu*.
1845.—*Nornis flavolivacea* Blyth, Journ. As. Soc. Bengal,
XIV, p. 590 (Nepal).

Cettia flavolivacea (Hodgs.)?, Uchida, "Dōbutsugaku Zasshi," XXII, p. 264 (1910).

Hab. Sakhalin (Uchida).

560. **Horornis acanthizoides concolor** (Ogilvie-Grant)
..... *Miyama-uguisu.*

1912.—*Horcites acanthizoides concolor* O.-Grant, Bull. B. O. C. XXIX, p. 107 (Formosa).

Hab. Formosa.

561. **Horornis fortipes robustipes** (Swinhoe) ...*Taiwan-ko-uguisu.*
1866.—*Horcites robustipes* Sw., P. Z. S., p. 39^s (Formosa).
Hab. Formosa.

Urosphena Swinhoe (1877).

562. **Urosphena squameiceps** (Swinhoe)*Yabasame, Shiosazai.*
1863.—*Tribura squameiceps* Sw., Proc. Zool. Soc. Lond., p. 292 (Canton).
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Loo-choo, Formosa,
Corea.

Locustella Kaup (1829).

563. **Locustella fasciolata** (Gray)*Yezo-senniu.*
1860.—*Microcephalus fasciolatus* Gray, Proc. Zool. Soc. Lond.
p. 349 (Batjan).
Hab. Saghalin, Hokkaido, Loo-choo, Corea.
564. **Locustella ochotensis ochotensis** (Middendorff).*Shima-senniu.*
1853.—*Sylvia (Locustella) ochotensis* Middendorff, Sibir. Reise II, 2, p. 185, pl. XVI, fig. 7 (Udskoj Ostrog).
Locustella ochotensis (Midd.). Hartert (pt.)
Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Formosa.
565. **Locustella ochotensis pleskei** Taczanowski *Ukiyama-senniu.*
1889.—*Locustella pleskei* Tacz., Proc. Zool. Soc. London, p.

620 (Corea).

- 1893.—*Locustella hondensis* Stejn., Proc. U. S. Nat. Mus., XVI, p. 957 (Hondo, Japan).

Hab. Hondo (Shinano), Kiusiu (Chikuzen), Corea (Chemulpo).

566. **Locustella certhiola** (Pallas) *Shibertia-senniu*.

- 1827.—*Motacilla Certhiola* Pallas, Zoogr. Ross-Asiat. I. p. 509 (Baikal Region)

Hab. Corea.

567. **Locustella lanceolata** (Temminck) *Makino-senniu*.

- 1840.—*Sylvia lanceolata* Temm., Man. d'Ornith. ed. II, IV, p. 614 (Russia).

Hab. Saghalin, Kurile Isls., Hokkaido, Corea.

Bradypterus Swainson (1837).

568. **Bradypterus luteoventris** (Hodgson) *Taiwan-ōsekka*.

- 1845.—*Tribura luteoventris* Hodgson, Proc. Zool. Soc. London, p. 30 (Nepal).

Lusciniola luteoventris (Hodgs.), Hartert.

Hab. Formosa.

569. **Bradypterus pryeri pryeri** (Seeböhm) *ō-sekka*.

- 1884.—*Megalurus pryeri* Seeböhm, Ibis, p. 40. (Yokohama, Japan).

Hab. Hondo (Yokohama, Tokyo, Owari).

Phragmaticola Jerdon (1845).

570. **Phragmaticola aëdon** (Pallas) *Harshibuto-ōyoshikiri*.

- 1776.—*Muscicapa Aëdon* Pallas, Reise d. versch. Prov. Russ. Reichs III, p. 695 (Dauria).

Lusciniola aëdon (Pall.), Cat. of B. Ri-ōke Museum, p. 59, 1918.

Hab. Corea.

Acrocephalus Naumann (1811).

571. **Acrocephalus arundinaceus orientalis** (Temminck and Schlegel) *Ō-yoshikiri.*
1847.—*Salicaria turdina orientalis* T. and S., Fauna Jap., Aves, p. 50. pl. XXB (Japan, Borneo, Makassar and Sumatra).
Acrocephalus orientalis (T. and S.), Uchida.
Hab. Hokkaido, Hondo, Formosa, Corea.
572. **Acrocephalus bistrigiceps** Swinhoe *Koyoshikiri.*
1860.—*Acrocephalus bistrigiceps* Swinhoe, Ibis, p. 51 (Amoy).
Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Corea.

Suya Hodgson (1806).

573. **Suya crinigera** Hodgson, subsp *Hauchiwa-dori.*
1836.—*Suya crinigera* Hodgson, Asiat. Research XIX, p. 183.
Hab. Formosa.

Burnesia Jerd (1863).

574. **Burnesia sonitans** (Swinhoe) *Awo-hauchiwa-dori.*
1858.—*Prinia sonitans* Swinhoe, Zoologist, p. 6229.
Hab. Formosa.

Prinia Horsfield (1821).

575. **Prina inornata formosa** Harington *Mami-hauchiwadori.*
1913.—*Prinia inornata formosa* Harington, Bull. B. O. C., XXXI, p. 111. (Formosa).
Prinia extensicauda (nec Swinhoe), Uchida.
Hab. Formosa.

Sutoria Nicholson (1851).

576. **Sutoria sutoria** (Forster) *Onaga-hauchiwadori.*

1781.—*Motacilla sutoria* Forster, Ind. Zool. p. 17, pl. 8.

Sutoria sutoria (Forst.), Sharpe, Cat. B. Br. Mus., VII, p. 215, 1883.

Long-tailed Warbler.

Hab. Formosa.

Cisticola Kaup (1829).

577. **Cisticola cisticola brunneiceps** (Temminck and Schleger) *Sekka*.

1850.—*Salicaria (Cisticola) brunneiceps* (T. and S.,) Fauna Jap. Aves. p. 134, pl. 20 (Japan).

Cisticola cisticola (nec Sharpe), Uchida

Hab. Hondo, Shikoku, Kiusiu, Loo-choo Isls.

578. **Cisticola cisticola tintinnabulans** (Swinhoe) *Kara-sekka*.

1859.—*Calamanthella tintinnabulans* Sw., Journ. A. S. N. China Branch, II, 1859; cf. Sw., Ibis, 1860, pp. 51, 131 (Amony, Shanghai, Formosa).

Cisticola cisticola (nec Temm), Uchida.

Hab. Formosa.

579. **Cisticola exilis volitans** Swinhoe *Taiwan-sekka*.

1859.—*Cisticola volitans* Swinhoe, Journ. A. S. N. China Branch, II, 1859; id, Ibis, 1863, p. 304; id. P. Z. S., p. 352 (Formosa).

Hab. Formosa.

Rhopophilus Giglioli and Salvadori (1870).

580. **Rhopophilus pekinensis pekinensis** (Swinhoe) *Karachimedori*.

1868.—*Drymoeca pekinensis* Sw., Ibs, p. 62 (Peking).

Hab. Corea.

Alcippe Blyth (1844).

581. **Alcippe nipalensis morrisonia** Swinhoe *Mijirochimedori*.

1863.—*Alcippe morrisonia* Swinhoe, Ibis, p. 296 (Formosa).
Hab. Formosa.

Proparus Hodgson (1844).

582. **Proparus formosanus** Ogilvie-Grant *Arisan-chimedoris*.
1906.—*Proparus formosanus* O.-Grant, Bull. B. O. C. XVI, p.
120 (Formosa).
Hab. Formosa.

Liocichla Swinhoe (1877).

583. **Liocichla steerii** Swinhoe *Yobudori*.
1877.—*Liocichla steerii* Swinhoe, Ibis, p. 474, pl. XIV
(Formosa).
Hab. Formosa.

Actinodura Gould (1836).

584. **Actinodura morrisoniana** Ogilvie-Grant *Shimadori*.
1906.—*Actinodura morrisoniana* O.-Grant, Bull. B. O. C. XVI,
p. 119 (Formosa).
Hab. Formosa.

Yuhina Hodgson (1836).

585. **Yuhina brunneiceps** Ogilvie-Grant *Kammuri-chimedoris*.
1906.—*Yuhina brunneiceps* O.-Grant, Bull. B. O. C. XVI, p.
121 (Formosa).
Hab. Formosa.

Stachyridopsis Sharpe (1883).

586. **Stachyridopsis ruficeps praecognitus** (Swinhoe)
..... *Dzuaka-chimedoris*.

1866.—*Stachyris praccognitus* Swinhoe, Ibis, p. 310
(Formosa).

Stachyrhodopsis praccognitus (Sw.), Uchida.

Hab. Formosa.

Schiœniparus Hume (1874).

587. **Schiœniparus brunneus brunneus** (Gould) *Chimedori*.

1862.—*Alcippe brunnea* Gould, P. Z. S. p. 280 (Formosa).

Hab. Formosa.

Lioptila Blyth (1847).

588. **Lioptila auricularis** (Swinhoe) *Mimijiro-chimedori*.

1864.—*Kittacincla auricularis* Swinhoe, Ibis, p. 361 (Formosa).

Malacias auricularis (Sw.), Uchida.

Hab. Formosa.

Herpornis Hodgson (1844).

589. **Herpornis xantholeuca tyrannulus** Swinhoe. *Awo-chimedori*.

1870.—*Herpornis tyrannulus* Swinhoe, Ibis, p. 347 (Hainan).

Hab. Formosa.

Dryonastes Sharpe (1883).

590. **Dryonastes poecilorhynchus** (Gould) *Take-dori*.

1862.—*Garrulax poecilorhyncha* Gould, P. Z. S. p. 281
(Formosa).

Hab. Formosa.

Trochalopteron Hodgson (1844).

591. **Trochalopteron taewanum** (Swinhoe) *Hoibii*.

1859.—*Garrulax taewanus* Swinhoe, Journ. As. Soc. N.
China, II, p. 228 (Formosa).

Hab. Formosa.

592. **Trochalopteron morrisonianum** Ogilvie-Grant. *Kimbaw-huibii*.
1906.—*Trochalopterum morrisonianum* O.-Grant, Bull. B. O.
C. XVI, p. 120 (Formosa).

Hab. Formosa.

Garrulax Lesson (1821).

593. **Garrulax ruficeps** Gould *Chagashira*.
1862.—*Garrulax ruficeps* Gould, P. Z. S. p. 281 (Formosa).
Hab. Formosa.

Pomatorhinus Horsfield (1820).

594. **Pomatorhinus musicus** Swinhoe *Hinemaruhashi*.
1859.—*Pomatorhinus musicus* Swinhoe, Jour. As. Soc. N.
China, II, p. 228 (Formosa).
Hab. Formosa.
595. **Pomatorhinus erythrocnemis** Gould *Maruhashi*.
1862.—*Pomatorhinus erythrocnemis* Gould, P. Z. S. p. 281
(Formosa).
Hab. Formosa.

Family PRUNELLIDÆ.

Prunella Vieillot (1816).

596. **Prunella collaris erythrropygius** (Swinhoe) *Iwa-hibari*.
1870.—*Accentor erythrropygius* Swinhoe, P. Z. S. p. 124, pl.
IX (N. China).
Hab. Hondo, Corea.
597. **Prunella collaris nipalensis** (Blyth) *Miyama-iwahibari*.
1843.—*Accentor Nipalensis* Blyth, Journ. A. Soc. Bengal XII,
p. 958 (Nepal).
Hab. Formosa.

Tharrhaleus Kaup (1829).

598. **Tharrhaleus rubidus** (Temminck and Schleger). *Kayakuguri*.
1850.—*Accentor modularis rubidus* Temminck and Schleger,
Faun. Jap. Aves, p. 69, pl. 39 (Japan).
Prunella rubidus (Temm. and Schleg), Hartert.
Accentor rubidus (Temm. and Schleg), Uchida.
Hab. Kurile Isls., Hokkaido, Hondo, Kiusiu.
599. **Tharrhaleus montanellus** (Pallas) *Yamahibari*.
1776.—*Motacilla montanella* Pallas, Reise versch. Prov.
Russ. Reichs II, I, p. 695 (Dauria).
Accentor montanella (Pallas), Uchida.
Prunella montanella (Pallas), Hartert.
Hab. Hokkaido, Hondo, Corea.

Family TROGLODYTIIDÆ.

Troglodytes Vieillot (1807).

600. **Troglodytes troglodytes daurica** Dybowski and Taczanowski.
..... *Karafuto-misosazai*.
1884.—*Troglodytes dauricus* Dybowski and Taczanowski, Bull
Soc. Zool. France, p. 155 (Dauria).
Troglodytes fumigatus dauricus Dyb. and Tacz., Uchida.
Hab. Saghalin.
601. **Troglodytes troglodytes kurilensis** Stejneger
..... *Chishima-misosazai*.
1888.—*Troglodytes fumigatus kurilensis* Stejneger, U. S. Nat.
Mus., XI, p. 548 (N. Kuriles).
Hab. Kurile Isls.
602. **Troglodytes troglodytes fumigata** Temminck *Misosazai*.
1835.—*Troglodytes fumigatus* Temminck, Man. d'Orn. II. ed.,
III, p. 161 (Japan).

Hab. Hokkaido, Hondo, Shikoku, ? Kiushiu.

603. **Troglodytes troglodytes utanoi** Kuroda. *Tsushima-misosazai*.
1922.—*Troglodytes troglodytes utanoi* Kuroda, Ibis, p. 96
(Tsushima).

Hab. Iki Is., Tsushima.

604. **Troglodytes troglodytes ogawae** Hartert ... *Ogawa-misosazai*.
1910.—*Troglodytes troglodytes ogawae* Hartert, Vög. Pal. I,
p. 784 (Yakushima).

Troglodytes fumigatus ogawae Hartert, Uchida.

Hab. Tanegashima, Yakushima.

605. **Troglodytes troglodytes taivana** Hartert ... *Arisan-misosazai*.
1910.—*Troglodytes troglodytes taivanus* Hartert, Vög. Pal. I,
p. 776 (Formosa).

Hab. Formosa.

606. **Troglodytes troglodytes peninsulæ** (Clark)
..... *Chōsen-misosazai*.
1907.—*Olbiorchilus fumigatus peninsulae* Clark, P. U. S. Nat.
Mus., XXXII, p. 474 (Fusan, Corea).
Troglodytes fumigatus peninsulae (Clark), Uchida.

Hab. Corea.

Note. **Troglodytes troglodytes** subsp., Kuroda and Mori, "Tori," vol. II, No. 7,
p. 81 (Quelpart Isl.).

Pnoepyga Hodgson (1845).

607. **Pnoepyga albiventris formosana** Ingram. *Takasago-misosazai*.
1909.—*Pnoepyga formosana* Ingram, Bull. B. O. C. XXIII, p.
97 (Formosa).

Hab. Formosa.

Cinclus Borkhausen (1797).

608. **Cinclus pallasii pallasii** Temminck *Kawagurasu*.

1820.—*Cinclus Pallasii* Temminck, Man. d'Orn. II, ed. I, p. 177 ("Krim?," errore!).

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Yakushima, Corea.

609. ***Cinclus pallasii marila* Swinhoe**.....*Takasago-kawagarasu*.

1860.—*Cinclus marila* Swinhoe, Ibis, p. 187 (Formosa).

Cinclus pallasii (nec Temm.), Uchida.

Hab. Formosa.

Family HIRUNDINIDÆ.

Hirundo L. (1758).

610. ***Hirundo rustica gutturalis* (Scopoli)**.....*Tsubame, Tsubakuro*.

1765.—*Hirundo gutturalis* Scopoli, Del Flor. and Faun. Insubr. II. p. 96 (Philippines).

Chelidon rustica gutturalis (Scop.), Hartert.

Eastern Chimney-Swallow.

Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Loochoo, Formosa, Corea, Quelpart Is, Bonin Isls.,

611. ***Hirundo rustica tytleri* (Jerdon)***Akahara-tsubame*.

1864.—*Hirundo Tytleri* Jerdon, B. India III, p. 870 (India).

Chelidon rustica tytleri (Jerd.), Hartert.

Tytler's Chimmey-Swallow.

Hab. Kiusiu, Corea.

612. ***Hirundo daurica nipalensis* Hodgson***Koshiaka-tsubame*.

1837.—*Hirundo Nipalensis* Hodgson, Jour. As. Soc. Bengal. V, "1836" p. 380 (C. Nepal).

Chelidon daurica nipalensis (Hodgs.), Hartert

Hodgson's Mosque-Swallow.

Hab. Hondo, Shikoku, Kiusiu, Corea.

613. **Hirundo daurica striolata** (Temminck and Schlegler)
 *Ō-koshiakatsubame*.
 1847.—*Hirundo striolata* Temm. and Schleg., Faun. Jap.
 Aves, p. 33 (Java).
Chelidon daurica striolata (Temm. and Schleg.), Hartert.
 Javan Mosque-Swallow.
- Hab. Formosa.
614. **Hirundo javanica namiyei** (Stejneger) *Riu-kiu-tsubame*.
 1866.—*Hirundo namiyei* Stejneger, P. U. S. Nat. Mus. XI, p.
 646 (Riu-Kiu Isls.)
Chelidon javanica namiyei (Stejn.), Hartert.
 Liu-Kiu Swallow.
 Hab. Loo-choo Isls.
- Delichon Moore (1855).**
615. **Delichon urbica whiteleyi** (Swinhoe) *Siberia-iwatsubame*.
 1862.—*Chelidon whiteleyi* Sw., Proc. Zool. Soc. London, p.
 320 (Peking).
 Hab. Sakhalin (Hartert).
616. **Delichon urbica dasypus** (Bonaparte) *Iwa-tsubame*.
 1850.—*Chelidon dasypus* Bonaparte, Conspr. Av. I, p. 343
 (Borneo).
 1862.—*Chelidon blakistoni* Swinh., P. Z. S., p. 320 (Hakodate,
 Japan).
Chelidon urbica dasypus Bp., Uchida.
Hirundo urbica dasypus (Bp.), Hartert.
 Blakiston's Martin.
 Hab. Kurile Isls., Hokkaido, Hondo.
617. **Delichon urbica nigrimentalis** (Hartert) .. *Hime-iwa-tsubame*.
 1910.—*Hirundo urbica nigrimentalis* Hartert, Vög. pal., p.

810 ("Kuatun," Fukien, S. W. China).

Chelidon urbica nigromentalis (Hart.), Kuroda.

Hab. Formosa.

Riparia Forster (1817).

618. **Riparia riparia ijimæ** (Lönnberg)

..... *Shōdō-tsubame*, *Sunamuguri-tsubame*.

1908.—*Clivicola riparia ijimæ* Lönnberg, Jour. Coll. Sc. Tokyo, XXIII. pt. 14, p. 38 (Sakhalin).

Cotile riparia (nec L.), Uchida.

Riparia riparia ijimæ (Lönnb.), Hartert.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo.

619. **Riparia paludicola chinensis** (J. E. Gray)

..... *Chōsen-shōdō-tsubame*.

1830-32.—*Hirundo chinensis* J. E. Gray, in Hardw. Ill. Ind. Zool. pl. 35, Fig. 3.

Cotile sinensis Tacz., Uchida.

Indian Sand-Martin.

Hab. Corea, Formosa.

Family CAMPEPHAGIDÆ.

Graucalus Cuvier (1817).

620. **Graucalus macei rex-pineti** Swinhoe *Oni-sanshokui*.

1863.—*Graucalus rex-pineti* Swinhoe, Ibis, p. 265 (Formosa).

Hab. Formosa.

Volvocivora Hodgson (1837).

621. **Volvocivora lugubris asakurai** (Kuroda). *Asakura-sanshokui*.

1920.—*Campyphaga lugubris asakurai* Kuroda, "Dōbutsugaku Zasshi" (=Tokio Zool. Mag.) XXXII, pp. 244 and 247 (Horisha, Formosa).

Hab. Formosa (Horisha, Nanto Distr.).

Pericrocotus Boie (1826).

622. **Pericrocotus cinereus cinereus** Lafresnaye *Sanshokui*.

1845.—*Pericrocotus cinereus* Lafresnaye, Rev. Zool., p. 94 (Luzon).

1887.—*Pericrocotus japonicus* Stejn., Proc. U. S. Nat. Mus., IX, "1886," p. 649 (Hondo, Japan).

1907.—*Pericrocotus cinereus intermedius* Clark, op. cit., XXXII, p. 474 ("Seoul, Korea").

Pericrocotus cinereus Lafr., Hartert.

Ashy Minivet or Siberian Minivet.

Hab. Hondo, Seven Isls. (Nijima), Shikoku, Kiusiu, Tsushima, Corea, Formosa, Pescadores Isls. (Taisho).

623. **Pericrocotus tegimæ** Stejn. *Riukiu-sanshokui*.

1887.—*Pericrocotus tegimæ* Stejneger, Proc. U. S. Nat. Mus., IX, "1886," p. 648 (Okinawa, Riu-Kiu Isls.).

Loo-choo Minivet.

Hab. S. Kiusiu (Satsuma), Tanegashima, Yakushima, Amami Ōshima, Riu-Kiu (Okinawa, Ishigakshima, Irimoteshima).

624. **Pericrotus solaris griseigularis** Gould *Beni-sanshokui*.

1862.—*Pericrotus griseigularis* Gould, Proc. Zool. Soc. Lond., p. 282 (Formosa).

Hab. Formosa.

Family DICRURIDÆ.

Captia Hodgson (1837).

625. **Captia brauniana** Swinhoe.....*Hime-ōchū*.

1863.—*Captia brauniana* Swinhoe, Ibis, p. 269 (Formosa).

Hab. Formosa.

Buchanga Hodgson (1837).

426. **Buchanga atra cathœca** (Swinhoe)*Ōchū*.

1871.—*Dicrurus cathacus* Swinhoe, Proc. Zool. Soc. Lond., p. 377 (China, Hainan and Formosa).

Buchanga atra (nec Hermann), Uchida.

Chinese Black-Drongo.

Hab. Formosa.

Family AMPELIDÆ.

Bombycilla Vieillot (1807).

627. **Bombycilla garrula centralasiae** Poljakov*Ki-renjaku*.

1915.—*Bombycilla garrulus centralasiae* Poljakov, Messager Ornith., VI, p. 137 (Turkestan, C. Asia).

Bombycilla garrulus garrulus (nec L.), Hartert (part.).

Ampelis garrulus (nec Linn.), Uchida.

Eastern Waxwing.

Hab. Saghalin, Hokkaido, Hondo, Seven Isls. (Miyakeshima), Kiusiu, Corea.

628. **Bombycilla japonica** (Siebold)*Hi-renjaku*.

1824.—*Bombyciora japonica* Siebold, De hist. Nat. Japon. statu etc., p. 13 (".....in prov. Tyko et Tsikusen habitat") (Tokyo and Chikuzen).

1827.—*Bombycilla phoenicoptera* Temminck, Pl. Col. II, 450 (Japan).

Ampelis japonicus Seebold (errore!), Uchida.

Japanese Waxwing.

Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Okinawa,
Corea, Formosa.

Family LANIIDÆ.

Lanius L. (1758).

- 629. *Lanius tigrinus* Drapiez** ... *Cligo-mozu, Tora-mozu.*
1828.—*Lanius tigrinus* Drapiez, Dist. Class. Hist. Nat. XIII,
p. 523 (Java).
Thick-billed Shrike.
Hab. Hondo, Corea, Tsushima, South Islands of Satsuma.
- 630. *Lanius sphenocercus sphenocercus* Cabanis** ... *Ō-kara-mozu.*
1873.—*Lanius sphenocercus* Cabanis, Journ. f. Ornith., p. 73
(Canton).
Lanius sphenocercus Cab., Uchida.
Long-tailed Grey-Shrike.
Hab. Hondo (Kōbe-straggler!). Tsushima, Corea.
- 631. *Lanius excubitor mollis* Eversmann** ... *Ō-mozu.*
1853.—*Lanius mollis* Eversmann, Bull. Soc. Imp. Nat.
Moscou, XXVI, p. 498 (Southern Altai).
Mongolian Grey-Shrike.
Hab. Hokkaido, Hondo.
- 632. *Lanius excubitor bianchii* Hartert** ... *Karafuto-ōmozu.*
1907.—*Lanius excubitor bianchii* Hartert, Vög. pal. I, p. 424
(Sakhalin).
Hab. Saghalin, Hokkaido (Iburi-straggler!).
- 633. *Lanius schach schach* L.** ... *Takasago-mozu.*
1758.—*Lanius schach* L., Syst. Nat., X, I, p. 94 (China).

- 1863.—*Lanius schach* var. *formosæ* Swinh., Ibis, p. 270
 (Formosa).
- Hab. Formosa.
- 634. *Lanius cristatus superciliosus* Latham.....*Aka-mozu*.**
- 1801.—*Lanius Superciliosus* Latham, Ind. Ornith. Suppl., p.
 XX (Batavia, Java).
- Japanese Red-tailed-Shrike.
- Hab. Saghalin, Hokkaido, Hondo, Kiusiu.
- 635. *Lanius cristatus lusionensis* L.*Shima-mozu, Riukiu-mozu*.**
- 1766.—*Lanius lusionensis* L., Syst. Nat. ed. XII, I, p. 135
 (Luzon).
- Philippine Red-tailed-Shrike.
- Hab. Corea, Riukiu, Formosa, Pescadores Isls.
- 636. *Lanius bucephalus* Temminck and Schlegel*Mozu*.**
- 1844.—*Lanius bucephalus* Temminck and Schlegel, Siebold's
 Fauna Japon. Aves, p. 39, pl. 14 (Japan).
- Bull-headed Shrike.
- Hab. Saghalin, Hokkaido, Hondo, Seven Isls. (Ōshima and
 Hachijōshima), Shikoku, Kiusiu, Tanegashima, Yaku-
 shima, Amami-Ōshima, Riukiu, Corea, Tsushima, Quel-
 part Is.

Family SITTIDÆ.

Sitta L. (1758).

- 637. *Sitta europaea albifrons* Taczanowski ...*Shirobitac-gojūkara*.**
- 1882.—*Sitta albifrons* Taczanowski, Bull. Soc. Zool. France
 VII, p. 385 (Kamtschatka).
- Kamtschatkan Nuthatch.
- Hab. Northern Kurile Isls.
- 638. *Sitta europaea sakhalinensis* Buturlin ...*Shirohara-gojūkara*.**

- 1916.—*Sitta europaea sakhalinensis* Buturlin, Trav. Soc. Nat. Petrograd, XLIV, p. 170 (Sakhalin).
- Sitta europaea uralensis* (nec Gloger), Hartert (part.).
- Sitta europaea uralensis* (nec Licht.), Uchida.
- Saghalin Nuthatch.
- Hab. Saghalin.
- 639. *Sitta europaea clara* Stejneger** *Yesso-gojukara*.
- 1886.—*Sitta amurensis clara* Stejn., Proc. U. S. Nat. Mus., IX, pp. 390, 392 ("Yesso").
- Sitta europaea uralensis* (nec Gloger), Hartert (part.).
- Hab. Hokkaido, ? Kurile Isls.
- 640. *Sitta europaea amurensis* Swinhoe** *Amuru-gojukara*.
- 1871.—*Sitta amurensis* Swinhoe, Proc. Zool. Soc. Lond., p. 350 (Amurland).
- Amoor Nuthatch.
- Hab. Corea.
- 641. *Sitta europaea hondoensis* Buturlin** ... *Gojukara, Kimawari*.
- 1916.—*Sitta europaea hondoensis* Buturlin, Trav. Soc. Nat. Petrograd, XLIV, p. 171 (Hondo, Japan).
- Sitta europaea amurensis* (nec Sw.), Hartert (part.).
- Hondo Nuthatch.
- Hab. Hondo, Shikoku.
- Note.—*Sitta europaea* subsp. nov. ?, Hartert, Vög. pal. I, p. 331 (1905).
Hab. Kiusiu (Hiuga).
- 642. *Sitta europaea taivana* Buturlin** *Chabara-gojukara*.
- 1911.—*Sitta formosana* Buturlin, Nasa Ochota St. Petersburg V, 8, p. 51 (Formosa). Preoccupied.
- 1915.—*Sitta europaea taivana* Buturlin, Mess. Ornith, Moskva, VI, pp. 311-312 (Formosa). Nom. emend.
- Sitta europaea sinensis* (nec Verr.), Uchida.
- Hab. Formosa.

643. **Sitta europaea bedfordi** Ogilvie-Grant.....*Akahara-gojūkara.*
 1909.—*Sitta bedfordi* Ogilvie-Grant, Bul. B. O. C., XXIII, p.
 591 (Quelpart Island).
 Quelpart Island Nuthatch.
 Hab. Quelpart Is.
644. **Sitta canadensis corea** Ogilvie-Grant.....*Chōsen-gojūkara.*
 1906.—*Sitta corea* Ogilvie-Grant, Bull. B. O. C., XVI, p. 87
 (Séoul, Corea).
 Corean Grey-Nuthatch.
 Hab. Corea (Keiki Distr.).

Family REGULIDÆ.

Regulus Vieillot (1807).

645. **Regulus regulus japonensis** Blakiston*Kikuitadaki.*
 1862.—*Regulus japonensis* Blakiston, Ibis, 1862, p. 320 (Hako-
 date, Yezo).
 Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiusiu, Tanega-
 shima, Tsushima, Corea, Formosa.
646. **Regulus ignicapillus goodfellowi** Ogilvie-Grant.....
*Niitaka-kikuitadaki.*
 1906.—*Regulus goodfellowi* Ogilvie-Grant, Bull. B. O. C.,
 XVI, p. 122 (Mt. Morrison, Formosa).
 Hab. Formosa.

Family PARIDÆ.

Parus L. (1758).

647. **Parus major minor** Temminck and Schlegel*Shijūkara.*
 1848.—*Parus minor* Temminck and Schlegel, Fauna Japon.
 Aves, p. 70, pl. 33 (Japan).

Japanese Titmouse.

Hab. Saghalin, Hokkaido, Hondo, Seven Isls. (Miyakejima and Hachijōjima), Shikoku, ? Corea.

648. *Parus major dageletensis* Kuroda and Mori.
..... *Utsuryō-shijūkara.*

1920.—*Parus major dageletensis* Kuroda and Mori, “Tori,” ii,
No. 10, pp. 270 and 278 (Dagelet Island or Utsuryōtō).

Dagelet Island Titmouse.

Hab. Dagelet Is. in the Sea of Japan.

649. *Parus major quelpartensis* Kuroda *Shima-shijūkara.*
1917.—*Parus major quelpartensis* Kuroda, “Tori,” i, No. 5,
p. 3, pl. VI (Quelpart Island).

Quelpart Island Titmouse.

Hab. Quelpart Is., Tsushima, Iki, Northern Kiusu.

650. *Parus major kagoshimæ* Taka-Tsukasa. *Kagoshima-shijūkara.*
1919.—*Parus major kagoshimæ* Taka-Tsukasa, “Dōbutsugaku
Zasshi” (Tokio Zool. Mag.), XXXI, p. 55 (Prefect.
Kagoshima in Southern Kiushiu).

Taka-Tsukasa’s Titmouse.

Hab. Kiusu (Kagoshima).

651. *Parus major okinawæ* Hartert
..... *Riukiu-shijūkara, Okinawagara.*

1905.—*Parus major okinawæ* Hartert, Vög. pal., p. 346 (Oki-
nawa).

Hab. Amami-Ōshima, Tokunoshima, Okinawashima.

652. *Parus nigriloris* Hellmayr. *Ishigakigara.*
1900.—*Parus nigriloris* Hellmayr, Ornith. Monastber., VIII,
p. 139 (Ishigaki).

1901.—*Parus stejnegeri* Bangs, Bull. Mus. Comp. Zoöl.
Harvard, XXXVI, p. 267 (Ishigaki).

Hab. S. Riu-ku (Ishigakishima, Iriomoteshima).

653. **Parus monticolus insperatus** Swinhoe *Kibara-shijūkara*.
1866.—*Parus insperatus* Swinhoe, Ibis, pp. 308, 402 (Formosa).
Hab. Mountains of Formosa.

Machlolophus Cabanis (1851).

654. **Machlolophus holsti** (Seeböhm) *Taiwan-shijukara*.
1894.—*Parus holsti* Seeböhm, Bull. B. O. C. IV, p. 7
(Formosa).
Hab. Mountains of Formosa.

Sittiparus Selys-Longchamps (1884).

655. **Sittiparus varius varius** (Temminck and Schlegel)
..... *Yamagara*.
1848.—*Parus varius* Temminck and Schlegel, Fauna Japon.
Aves, p. 71, pl. 35 (Hondo, Japan).
1890.—*Parus sieboldi* Seeböhm, B. Jap. Empire, p. 85 (new
name for *P. varius*).
1862.—*Parus rubidus* Blakiston, Ibis, p. 321 (err. typ.).
1918.—*Parus varius hakodatensis* Momiyama, “Dōbutugaku
Zasshi,” XXX, p. 345 (Hakodate). Colour variety!
Parus varius varius Temm. and Schl., Hartert. (pt.).

Varied Titmouse.

Hab. Kurile Isls., Hokkaido, Hondo.

656. **Sittiparus varius namiyei** (Kuroda) *Namiye-yamagara*.
1918.—*Parus varius namiyei* Kuroda, “Dōbutugaku Zasshi”
(Tokio Zool. Mag.), XXX, pp. 316 and 322 (Niijima,
Seven islands).

Namiye's Varied Titmouse.

Hab. N. Seven Isls. (Niijima).

657. **Sittiparus varius ijimae** Kuroda *Tsushima-yamagara*.
1922.—*Sittiparus varius ijimae* Kuroda, Ibis, p. 98 (Tsushima).

- Hab. Shikoku, N. Kiusiu, Tsushima.
- 658. *Sittiparus varius sunsunpi* (Kuroda) ... *Tanegashima-yamagara.***
- 1919.—*Parus varius sunsunpi* Kuroda, “Dōbutsugaku Zasshi” (Tokio Zool. Mag.), XXXI, pp. 230 and 232 (Tanegashima).
- Hab. Tanegashima.
- 659. *Sittiparus varius yakushimensis* (Kuroda) ... *Yakushima-yamagara.***
- 1919.—*Parus varius yakushimensis* Kuroda, “Dōbutsugaku Zasshi” (Tokio Zool. Mag.) XXXI, pp. 230 and 232 (Yakushima).
- Yakushima Varied Titmouse.
- Hab. Yakushima, Amami Ohshima (straggler!).
- 660. *Sittiparus varius saisiuensis* (Kuroda and Mori) ... *Saisiu-yamagara.***
- 1920.—*Parus varius saisiuensis* Kuroda and Mori, “Tori” II, No. 10, pp. 272 and 279, pl. V. fig. 5 (Quelpart Island).
Quelpart Is. Varied Titmouse.
- Hab. Quelpart Is. and Corea.
- 661. *Sittiparus varius utsurioensis* (Kuroda and Mori) ... *Shima-yamagara.***
- 1920.—*Parus varius utsurioensis* Kuroda and Mori, “Tori” II, No. 10, pp. 271 and 280, pl. V. fig. 2 (Dagelet Island or Utsuryōtō).
Dagelet Is. Varied Titmouse.
- Hab. Dagelet Is. in the Sea of Japan.
- 662. *Sittiparus varius owstoni* (Ijima) ... *Owston-yamagara.***
- 1893.—*Parus owstoni* Ijima, “Dōbutsugaku Zasshi” (Tokio Zool. Jap.) V, p. 445 (Miyakeshima, Seven Islands).
Parus varius owstoni Ijima, Hartert.
- Hab. Seven Isls. (Miyakeshima, Hachijōshima).

- 663. *Sittiparus varius amamii* Kuroda. *Amami-yamagara*.**
- 1905.—*Parus varius*, subsp. nov. ?, Hartert, Vög. pal., I,
p. 355 (Okinawashima).
- 1922.—*Sittiparus varius amamii* Kuroda, Annot. Zool.
Japon., Vol. X, Art. 4, p. 117 (Amami-Ōshima).
- Parus castaneoventris* (nec Gould), Stejn., Proc. U. S. Nat.
Mus., 1886, p. 650.
- Hab. Amami-Ōshima, Riukiu Isls. (Okinawashima).
- 664. *Sittiparus varius castaneoventris* (Gould) *Taiwan-yamagara*.**
- 1862.—*Parus castaneoventris* Gould, Proc. Zool. Soc. Lond.
p. 280 (Formosa).
- Parus varius castaneoventris* Gould, Hartert.
- Formosan Titmouse.
- Hab. Mountains of Formosa.
- Periparus Selys-Longchamps (1884).**
- 665. *Periparus ater amurensis* (Buturlin) *Kita-higara*.**
- 1907.—*Parus ater amurensis* Buturlin, Ornis, Monatsb., p. 80
(Amur and Ussuri).
- Parus ater insularis* (nec Hellm.) Uchida.
- Parus ater pekingensis* (nec David.) Lönnberg, Journ Col.
Sci. Jap. Nuiv. Tokyo, 1908, XXIII, p. 20.
- Amur Coal Tit.
- Hab. Saghalin, Corea.
- 666. *Periparus ater insularis* (Hellmayr) *Higara*.**
- 1902.—*Parus ater insularis* Hellmayr, Ornith. Jahrb. XIII, p.
36 (Suruga, Hondo, Japan).
- Japanese Coal-Tit.
- Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiushu, Yaku-
shima, Riukiu Isls.
- 667. *Periparus ater teraokai* Kuroda *Tsushima-higara*.**

1922.—*Periparus ater teraokai* Kuroda, Ibis, p. 100 (Tsushima).

Tsushima Coal-Tit.

Hab. Tsushima.

668. *Periparus ater ptilosus* (Ogilvie-Grant) *Taiwan-higara*.

1912.—*Parus ater ptilosus* Ogilvie-Grant, Bull. B. O. C., XXIX, p. 108 (Formosa).

Formosan Coal-Tit.

Hab. Mountains of Formosa.

Pœcile Kaup (1829).

669. *Pœcile atricapilla sachalinensis* (Lönnberg) *Ezo-kogara*.

1908.—*Parus atricapillus sachalinensis* Lönnberg, Jour. Col. Sci. Imp. Univ. Tokyo, XXIII, p. 20 (Saghalin).

Parus atricapillus baicalensis (nec Swinh.), Hartert. (part.).

Lönnberg's Marsh-Tit.

Hab. Saghalin, Kurile Isls., Hokkaido.

670. *Pœcile atricapilla restrictus* (Hellmayr) *Kogara*.

1879.—? *Parus palustris japonicus* Seebold, Ibis, p. 32. (Japan.) (part?).

1900.—*Parus borealis restrictus* Hellmayr, Orn. Jahrb., p. 215 ("Nipon").

Parus atricapillus restrictus Hellmayr, Hartert.

Japanese Marsh-Tit.

Hab. Hondo.

671. *Pœcile palustris hensoni* (Stejneger)

..... *Henson-kogara*, *Hensongara*.

1892.—*Parus hensoni* Stejneger, Proc. U. S. Nat. Mus. XV, p.

342 (Hakodate).

1892.—*Parus sebohni* Stejn., Proc. U. S. Nat. Mus., XV, p. 343 (Sapporo).

Parus palustris hensonii Stejn., Hartert.

Hab. S. Kurile Isls., Hokkaido.

672. *Pœcile palustris crassirostris* (Taczanowski) *Iashibuto-kogura*.

1885.—*Pœcilia palustris crassirostris* Taczanowski, Bull. S. Zool. France, x, p. 470 (Sidemi).

Parus palustris crassirostris (Tacz.), Hartert.

Hab. Saghalin, Corea.

673. *Pœcile palustris hellmayri* (Bianchi) *Chōsen-kogura*.

1902.—*Parus palustris hellmayri* Bianchi, Ann. Mus. Zool. Acad. Imp. Sci. Petersb., VII, p. 236 (Peking).

Hab. Corea.

Ægithalos Hermann (1804).

674. *Ægithalos caudatus caudatus* (L.) *Shima-enaga*.

1758.—*Parus caudatus* L., Syst. Nat. ed. X, p. 190 (Sweden).

1897.—*Aegithalus caudatus japonica* Prazák, J. f. O., p. 291 (Northern Japan).

1907.—*Acredula trivirgata magna* Clark, Proc. U. S. Nat. Mus., XXXII, p. 475 ("Seoul, Korea").

Acredula caudata (L.), Uchida.

White-headed Long-tailed Tit.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo (Mutsu, Shimo-tsuke-straggler !), Corea.

675. *Ægithalos caudatus trivirgatus* (Temminck and Schlegel) ... *Enaga*.

1848.—*Parus (Megisturus) trivirgatus* Temminck and Schlegel, Fauna Japon., Aves, p. 71, pl. 34 (Japan).

Acredula caudata trivirgata (Temm. and Schl.), Uchida.
Japanese Long-tailed Tit.
Hab. Hondo, Shikoku, Kiushiu, Tsushima, Quelpart Is.

Ægithaliscus Cabanis (1851).

676. **Ægithaliscus concinnus concinnus** (Gould)
..... *Zuaka-enaga*, *Zuakagara*.
1855.—*Psaltria concinna* Gould, B. Asia, II, 65 (China).
..... *Egithalus concinnus* (Gould), Uchida.
Hab. Mountains of Formosa.

Remiz Jarocki (1821).

677. **Remiz pendulinus consobrinus** (Swinhoe).
..... *Suinho-gara*.
1870.—*Egithalus consobrinus* Swinhoe, Proc. Zool. Soc.
London, p. 133 (Jangtse-kiang, China).
1907.—*Remiz consobrinus suffusus* Clark, Proc. U. S. Nat.
Mus., XXXII, p. 474 (Fusan, Corea).
1907.—*R[emiz] c[onsobrinus] japonicus* Clark, Proc. U. S.
Nat. Mus., XXXII, p. 475 (Japan).
Anthoscopus pendulinus consobrinus (Swinh.), Hartert.
Chinese Penduline Tit or Swinhoe's Penduline Tit.
Hab. Corea, N. Kiushiu (Vicinity of Nagasaki-straggler),
Hondo (Chiba-straggler).

Suthora Hodgson (1837).

678. **Suthora webbiana webbiana** Gray *Daruma-enaga*.
1852.—*Suthora Webbiana* Gray, Proc. Zool. Soc. Lond., p. 70,
pl. 49 (Shanghai).
1892.—*Suthora fulvicauda* Campbell, Ibis, p. 237 (Chemulpo,
Corea, juv !

- 1892.—*Suthora longicauda* Campbell, Ibis, p. 237. (Near Seoul, Corea, ♀ ad. !).
 Chinese Suthora or Chinese Crow-Tit.
 Hab. Corea.
- 679. Suthora webbiana bulomachus** Swinhoe
 *Takasago-daruma-enaga.*
 1866. *Suthora bulomachus* Swinhoe, Ibis, p. 300, pl. 9 (Formosa).
 Hab. Formosa.
- 680. Suthora gularis morrisoniana** Ogilvie-Grant
 *Kibane-daruma-enaga.*
 1906.—*Suthora morrisoniana* Ogilvie-Grant, Bull. B. O. C.,
 XVI, p. 119 (Mt. Morrison, Formosa).
 Hab. Mountains of Formosa.

Panurus Koch (1816).

- 681. Panurus biarmicus russicus** (Brehm) *Higegara.*
 1831.—*Mystacinus Russicus* Brehm, Handb. Naturg. Vög.
 Dentschl., p. 472 (Russia).
 Hab. N. Hondo (Prefect. Yamagata-straggler !)

Family ORIOLIDÆ.

Oriolus L. (1766).

- 682. Oriolus indicus indicus** Jerdon *Korai-uguisu.*
 1847.—*Oriolus indicus* Jerdon, Ill. Ind. Ornith. text to pl. 15
 (Malabar, Travancore, etc.).
 Black-naped Oriole.
 Hab. Corea, Formosa.

Psaropholus Jardine and Selby (1837).

- 683. Psaropholus trailli ardens** Swinhoe *Higoromo.*

1862.—*Psaropholus ardens* Swinh., Ibis, p. 363 (Formosa).

Oriolus ardens (Sw.), Uchida.

Hab. Formosa.

Family PLOCEIDÆ.

Munia Hodgson (1836).

684. **Munia atricapilla formosana** Swinhoe *Taiwan-kinpara*.

1865.—*Munia formosana* Swinhoe, Ibis, p. 366 (Formosa).

Formosan Black-headed Mannikin.

Hab. Formosa.

685. **Munia punctulata topela** Swinhoe. *Shima-kinpara*.

1863.—*Munia topela* Swinhoe, Ibis, p. 380 (Formosa).

Topela-Finch.

Hab. Formosa.

Uroloncha Cabanis (1850).

686. **Uroloncha acuticauda squamicollis** Sharpe

Koshijiro-kinpara.

1890.—*Uroloncha squamicollis* Sharpe, Cat. Brd. Brit. Mus.

XIII, p. 359 ("China; Formosa and Hainan").

Uroloncha acuticauda (nec Hodgs.). Uchida.

Chinese Sharp-tailed Finch.

Hab. Formosa.

Family CORVIDÆ.

Corvus Linnæus (1758).

687. **Corvus corax kamtschaticus** Dybowski *Watarigurasu*.

1883.—*Corvus corax kamtschaticus* Dyb., Bull. Soc. Zool.

France, pp. 362, 363 (Kamtschatka).

Corvus corax (nec L.), Uchida.

Kamtschatkan Raven.

Hab. Saghalin, Kurile Isls., Hokkaido.

688. **Corvus coronoides japonensis** Bonaparte *Hashibutogarasu*.

1850.—*Corvus japonensis* Bonaparte, Conspl. Av., I, p. 386
(Japan. Type locality was designated by Stresemann as
“Yesso”).

Corvus macrorhynchos japonensis Bp., Hartert.

Corvus macrorhynchos japonensis Bp., Uchida.

Japanese Jungle-Crow.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Seven Isls.,
Bonin Isls., Shikoku, Kiusiu, Tanegashima, Yakushima,
? Dagelet Is., ? Quelpart Is.

689. **Corvus coronoides mandshuricus** Buturlin

Chōsen-hasibutogarasu.

1913.—*Corvus coronoides mandshuricus* Buturlin, Mess. Orn.
IV, No. 1, p. 40 (Ussuriland).

Manchurian Jungle Crow.

Hab. Corea.

690. **Corvus coronoides connectens** Stresemann

Riukiu-hasibutogarasu.

1916.—*Corvus coronoides connectens* Stresem., Verh. Orn.
Ges. Bayern, XII, 4, p. 281 (“Miyako-shima”).

Corvus macrorhynchos levillanti (nec Lsss.), Uchida.

Loo-choo Jungle-Crow.

Hab. ? Tsushima, Amami-Ōshima, Okinawashima, Miyako-
shima, Ishigakishima.

691. **Corvus coronoides osai** Ogawa *Osagarasu*.

1905.—*Corvus macrorhynchos osai* Ogawa, Annot. Zool. Jap.,
v, p. 196 (Kobama-shima, S. Riukiu).

Ogawa's Jungle-Crow.

Hab. Okinawashima, Ishigakishima, Iriomoteshima, Kobama, Aragusuku.

692. **Corvus coronoides colonorum** Swinhoe
..... *Taiwan-hashibutogarasu.*

1864.—*Corvus colonorum* Swinh., Ibis, p. 427 (Formosa).

Formosan Jungle Crow.

Hab. Formosa.

693. **Corvus corone interpositus** Laubmann ... *Hashiboso-garasu.*
1917.—*Corvus corone interpositus* Laubmann, Verh. Orn. Ges.

Bayern, XIII, 2, p. 201 (Hondo, Japan).

Corvus corone orientalis (nec Everms.), Hartert (Part.).

Japanese Carrion-Crow.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusu, Tsushima, Quelpart Is., Corea.

Trypanocorax Sandevall (1872).

694. **Trypanocorax frugilegus pastinator** (Gould). *Miyama-garasu.*

1845.—*Corvus pastinator* Gould, Proc. Zool. Soc. Lond., p. 1
(Chusan, China).

Corvus frugilegus pastinator Gould, Hartert.

Eastern Rook.

Hab. Hondo, Shikoku, Kiusu, Tsushima, Corea, Quelpart Is., Formosa.

Colœus Kaup (1829).

695. **Colœus dauricus dauricus** (Pallas)
..... *Kokumaru-garasu, Tōgarasu.*

1776.—*Corvus dauricus* Pallas, Reise Russ Reichs, III, p. 697
(Baikal Region).

1859.—*Corvus neglectus* Schlegel, Bijdr. Dierk. Amsterdam,
Afl. 8, Corvus, p. 16 (Japan). (Dark phase!).

Coloeus dauricus (Pall.), Hartert.

Coloeus neglectus (Schleg.), Hartert.

Daurian Jackdow.

Hab. Hokkaido, Hondo, Kiusiu, Corea.

Note.—**Coloeus neglectus** is a dark form of typical *dauricus*. (See Hartert, Vög.
pal, III, 1921, p. 2025).

Nucifraga Vieillot (1816).

696. **Nucifraga caryocatactes japonicus** Hartert
..... *Hoshigarasu, Takegarasu.*

1897.—*Nucifraga caryocatactes japonicus* Hartert, Nov.
Zool., p. 134 (Japan).

Japanese Nutcracker.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo.

697. **Nucifraga caryocatactes macrorhynchos** Brehm
..... *Hashinaga-hoshigarasu.*

1823.—*Nucifraga macrorhynchos* Brehm, Lehrb. Naturg.
europ. Vögel I, p. 103 ("Gebirgswälder mittl. nördl. Eur.
u. Asien").

Nucifraga caryocatactes macrorhynchus Brehm, Taka-
tsukasa in Uchida's Brd. Jap.

Long-billed Nutcracker.

Hab. Corea.

698. **Nucifraga caryocatactes owstoni** Ingram *Taiwan-hoshigarasu.*

1909.—*Nucifraga owstoni* Ingram, Bull. B. O. C., XXV, p.
86 (Formosa).

Formosan Nutcracker.

Hab. Mountains of Formosa.

Pica Vieillot (1816).

699. **Pica pica sericea** Gould *Kasasagi.*
1845.—*Pica sericea* Gould, Proc. Zool. Soc. Lond., p. 2
(Amoy, China).
1848.—*Pica varia japonica* Schlegel, Fauna Japon., Aves, p.
81.
Hab. Kiusiu (Prov. Chikuzen, Chikugo, Hizen, Higo), ? Tsu-
shima, Corea, Formosa.
700. **Pica pica bactriana** Bonaparte *Chishima-kasasagi.*
1850.—*Pica bactriana* Bonaparte, Conspl. Av. I, p. 383 (E.
Persia)
Pica pica kamtschatica Stejn., Uchida.
Hab. Kurile Islands.

Cyanopica Bonaparte (1850).

701. **Cyanopica cyanus interposita** Hartert *Koma-onaga.*
1917.—*Cyanopica cyanus interposita* Hartert, Nov. Zool.
XXIV, p. 493 (Tsin-ling Mt., Corea and Peking).
Cyanopica cyanus cyanus (nec Pall.), Hartert (part.).
North Chinese Azure-winged Magpie.
Hab. Corea.
702. **Cyanopica cyanus japonica** Parrot *Onaga.*
1905.—*Cyanopica cyanus japonica* Parrot, Orn. Monatsber., p.
26 (Japan).
Cyanopica cyanus cyanus (nec Pall.), Hartert. (part.).
Cyanopica cyanus (nec Pall.) Uchida.
Japanese Azure-winged Magpie.
Hab. Hondo, Kiusiu (Prov. Hizen).

Urocissa Cabanis (1850).

703. **Urocissa cærulea** Gould *Yamamusume.*

1862.—*Urocissa cœrulea* Gould, Proc. Zool. Soc., p. 282 (Formosa).

Formosan Blue-pie.

Hab. Formosa.

Dendrocitta Gould (1833).

704. **Dendrocitta formosæ formosæ** Swinhoe. ...*Taiwan-onagadori*.

1863.—*Dendrocitta sinensis*, var. *formosæ* Swinhoe, Proc. Zool. Soc., p. 387 (Formosa).

Dendrocitta formosæ Sw., Uchida.

Formosan Tree-pie.

Hab. Formosa.

Garrulus Vieillot (1816).

705. **Garrulus glandarius japonicus** Schlegel ...*Kakcsu, Kashidori*.

1814.—*Garrulus glandarius japonicus* Schlegel, Fauna Jap. Aves, p. 83, pl. 43 (Japan).

Garrulus japonicus Temm. and Schleg., Uchida.

Japanese Jay.

Hab. Hondo, Shikoku, Kiusiu, Yakushima.

706. **Garrulus glandarius namiyei** Kuroda*Tsushima-kakesu*.

1922.—*Garrulus glandarius namiyei* Kuroda, Ibis, p. 102 (Tsushima).

Hab. Tsushima.

707. **Garrulus glandarius brandtii** Eversmann ...*Miyama-kakesu*.

1842.—*Garrulus brandti* Eversmann, Add. Pall, Zoogr. fasc. III, p. 8 (Altai).

1908.—*Garrulus glandarius taczanowskii* Lönnberg, Journ. Coll. Sc. Tokyo, XXIII, Art. 14, p. 7 (Saghalin).

Brandt's Jay.

Hab. Saghalin, Kurile Isls., Hokkaido, Corea.

708. **Garrulus bispecularis taivanus** Gould ... *Takasago-kakesu*.
1862.—*Garrulus taivanus* Gould, Proc. Zool. Soc., p. 282
(Formosa).
1863.—*Garrulus insularis* Swinh., Proc. Zool. Soc., p. 304
(Formosa).
Hab. Formosa.

Lalocitta Reichenow (1906).

709. **Lalocitta lidthi** (Bonaparte) *Kuri-kakesu*.
1850.—*Garrulus lidthi* Bonaparte, Conspl. Av., I, p. 376
("Hab. Japan": Amami-Ōshima).
1921.—*Garrulus lidthi samuel* Floerike, Mitt. a. d. Vogel-
welt, 19, Nos. 3 and 4, p. 107. (Nomen nudum).
Lidthi's Jay.
Hab. Amami-Ōshima.

Perisoreus Bonaparte (1838).

710. **Perisoreus infaustus sakhaliensis** Buturlin *Akawo-kakesu*.
1916.—*Perisoreus infaustus sakhalensis* Buturlin, Mess.
Ornith., Moscow, pp. 40, 43 (Sakhalin).
Perisoreus infaustus sibiricus Bodd., Hartert (part.) (See
Vög. pal., III, p. 2034).
Hab. Saghalin.

Family STURNIDÆ.

Spodiopsar Sharpe (1889).

711. **Spodiopsar cineraceus** (Temminck) *Mukudori*.
1832.—*Sturnus cineraccus* Temminck, Pl. Col., p. 556 (Japan).
Grey Starling.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Amami-Ōshima,
Riukiu, Corea, Formosa.

Sturnia Lesson (1837).

712. **Sturnia violacea** (Boddaert) *Kō-mukudori*.
1783.—*Motacilla violacea* Boddaert, Tab. Pl. Enl., p. 11
(Philippines).
1850.—*Lamprotornis pyrrhogonyx* Temm. and Schl., Faun.
Japon., Aves, p. 86 (Japan).
Red-cheeked Myna.
Hab. Saghal'n, Kurile Isls., Hokkaiko, Hondo, Shikoku, Kiu-
siu, Riukiu, Formosa.
713. **Sturnia sinensis** (Gmelin) *Kara-mukudori*.
1788.—*Oriolus sinensis* Gmelin, Syst. Nat., I, p. 394 (China).
Chinese Myna.
Hab. Hondo (near Tokyo-straggler), Formos*t*.

Æthiopsar Sharpe (1889).

714. **Æthiopsar cristatellus formosanus** Hartert *Kāren*.
1912.—*Ethiopsar cristatellus formosanus* Hartert, Bull. B.
O. C., XXI, p. 14 (Formosa).
Ethiopsar cristatellus (nec L.), Uchida.
Formosan Crested-Myna.
Hab. Formosa.

Family ZOSTEROPIDÆ.

Zosterops Vigors and Horsfield (1826).

715. **Zosterops palpebrosa japonicus** Temminck and Schlegel.....
..... *Mejirō*.

1848.—*Zosterops japonicus* Temminck and Schlegel, Fauna Japon., Aves, p. 57 (Japan).

Japanese White-eye.

Hab. Hokkaido, Hondo, Northern Seven Is. (Oshima and Nijimi), Shikoku, Kiusiu.

716. *Zosterops palpebrosa stejnegeri* Seebold *Shichito-mejiro*.

1891.—*Zosterops stejnegeri* Seebold, Ibis, 1891, p. 273 (“Fatsizio or Hatchi-now-sima” = Hachijōshima).

Seven-island White-eye.

Hab. Seven Is., Bonin Isls.

717. *Zosterops palpebrosa alani* Hartert *Iwojima-mejiro*.

1905.—*Zosterops palpebrosa alani* Hartert, Bull. B. O. C., XV, p. 45 (“S. Dionisio, Volcano Islands”).

1911.—*Zosterops palpebrosa iwojimaeensis* Uchida, “Dōbutsugaku Zasshi,” XXIII, p. 537 (“Habitat: Minami Iwojima of the sulphur Islands”).

Sulphur-Island White-eye.

Hab. S. Sulphur Is. or S. Dionisio. Volcano Islands.

718. *Zosterops palpebrosa ijimae* Kuroda *Ijima-mejiro*.

1917.—*Zosterops palpebrosa ijimae* Kuroda, “Tori” I, No. 5, p. 4, pl. VI, fig. 3; fig. 2 (Tsushima).

Ijima’s White-eye.

Hab. Northern coasts of Kiusiu, Iki, Tsushima, Quelpart Is., S. Corea (S. Keishō Distr. and Zenra Distr.), Dagelet Is.

719. *Zosterops palpebrosa insularis* Ogawa *Shiman-mejiro*.

1905.—*Zosterops japonica insularis* Ogawa, Ann. Zool. Japan, v, p. 186 (Tanegashima and Yakushima).

Ogawa’s White-eye.

Hab. Tanegashima, Yakushima.

720. *Zosterops palpebrosa loochooensis* Tristram . . . *Rinkiu-mejiro*.

1889.—*Zosterops simplex* var. *loochoensis* Tristram, Ibis,
p. 229 (Loochoo Islands).

Loochoo White-eye.

Hab. Amami-Oshima, Tokunoshima, Kikaigashima, Okino-
Erabushima, Okinawashima, Ishigakshima, Iriomote-
shima.

721. *Zosterops palpebrosa peguensis* Baker..... *Hime-mejiro*.

1922.—*Zosterops palpebrosa peguensis* Baker, Ibis, p. 144
("Moulmein").

Zosterops palpebrosa simplex (nec Sw.), Uchida.

Hab. Formosa.

722. *Zosterops palpebrosa batanis* McGregor *Kikuchi-mejiro*.

1907.—*Zosterops palpebrosa batanis* McGregor, Phil. Journ.
Sci., 2, Sec. A, p. 343 (Batan Is.).

Hab. Botel Tobago or Kōtōshō, Kashōtō.

723. *Zosterops erythropleurus* Swinhoe. *Chōsen-mejiro*, *Ko-mejiro*.

1863.—*Zosterops erythropleurus* Swinhoe, Ibis, p. 294 (N.
China).

Chinese White-eye or Red-flanked White-eye.

Hab. Corea (N. Heian Distr.—straggler!).

Family DICÆIDÆ.

Dicæum Cuvier (1817).

724. *Dicæum formosum* Ogilvie-Grant..... *Hanadori*.

1912.—*Dicæum formosum* Ogilvie-Grant, Bull. B. O. C., p.
109 (Formosa).

Formosan Flower-pecker.

Hab. Mountains of Formosa.

- 725.** *Dicæum minullum uchidai* Kuroda. *Azæo-hanadori.*
 1920.—*Dicæum minullum uchidai* Kuroda, "Tori," p. 230
 (Horisha in Formosa).
 1912.—*Dicæum sp.*, Uchida, Annot. Zool. Japan. VIII, p. 203
 (Formosa).
 Hab. Formosa.

Family CERTHIIDÆ.

Certhia L. (1758).

- 726.** *Certhia familiaris familiaris* L. *Kito-kibashiri.*
 1758.—*Certhia familiaris* L., Syst. Nat. ed. X, p. 118
 (Sweden).
 Tree-creepers.
 Hab. Saghalin, Kurile Isls., Hokkaido, Corea.
727. *Certhia familiaris japonica* Hartert *Kibashiri.*
 1897.—*Certhia familiaris japonica* Hartert, Nov. Zool., p.
 138 (Hondo, Japan).
 Japanese Tree-creepers.
 Hab. Hokkaido (Sapporo), Hondo, Shikoku.

Family FRINGILLIDÆ.

Coccothraustes Pallas (1811).

- 728.** *Coccothraustes coccothraustes japonicus* Temm. and Schleg.
 *Shime.*
 1850.—*Coccothraustes vulgaris japonicus* Temminck and Schle-
 gel, Fauna Japon., Aves, pl. 51 (Japan).
 Japanese Hawfinch.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Corea, Quelpart Is.,
Riukiu, Formosa.

729. ? **Coccothraustes coccothraustes verticalis** Tugarinow and
Buturlin *Korafuto-shime*.

1911.—*Coccothraustes coccothrustes verticalis* Tugar. and But.,
Mater. Vög. Jeness. Gouv., in Mitt. Krasnojarsker Sektion
Russ. Geogr. Ges. 1911 (Baikal and Sakhalin).

Hab. Sakhalin Is.

Eophona Gould (1851).

730. **Eophona personata personata** (Temminck and Schlegel)
..... *Ikaru*.

1850.—*Coccothraustes personatus* Temminck and Schlegel,
Fauna Japon., Aves, p. 91, pl. 52.
Eophona personata Gould, Uchida.

Japanese Grosbeak, Japanese Masked Hawfinch.

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Quelpart Is., Formosa.

731. **Eophona personata maguirestris** Hartert ... *Hashibuto-ikaru*.
1896.—*Eophona personata magnirostris* Hartert, Bull. B. O.
C., V, p. XXXVIII (Siberia).

Large-billed Masked Hawfinch.

Hab. Corea (straggler).

732. **Eophona migratoria migratoria** Hartert *Ko-ikaru*.
1903.—*Eophona melanura migratoria* Hartert, Vög. Pal., p.
59 (" Sidimi ").

Migratory Black-tailed Hawfinch.

Hab. Hondo, Kiusiu, Corea, Quelpart Is., Formosa.

Chloris Cuvier (1800).

733. **Chloris sinica kawarahiba** (Temminck). *O-kawarahiba.*
1835.—*Fringilla kawarahiba* Temminck, Pl. Col., 588, fig. 1.
(Japan).
Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu.
734. **Chloris sinica clarki** Kuroda and Mori
..... *Matsushima-kawarahiba.*
1920.—*Chloris sinica clarki* Kuroda and Mori, “Tori,” II, No.
10, pp. 273 and 281 (Dagelet Is.).
Hab. Dagelet Is. or Matsushima in the Sea of Japan.
735. **Chloris sinica ussuriensis** Hartert *Chōsen-kawarahiba.*
1903.—*Chloris sinica ussuriensis* Hartert, Vög. pal., I, p. 64
(“Sidemi-Mündung, Ostsibirien”).
Hab. Sakhalin, Corea.
736. **Chloris sinica minor** (Temm. and Schleg.) *Ko-kawarahiba.*
1850.—*Fringilla kawarahiba minor* Temminck and Schlegel,
Fauna Japon., Aves., p. 89, pl. 49 (Japan).
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Quelpart Is., Tsu-
shima, ? Formosa.
737. **Chloris sinica kittlitzii** (Seebohm). *Ogasawara-kawarahiba.*
1890.—*Fringilla kittlitzii* Seebohm, Ibis, p. 101 (Bonin
Islands).
Hab. Bonin Isls. (Nakondo-shima, Haha-shima), Sulphur Isls.

Spinus Koch (1816).

738. **Spinus spinus** (L.). *Mahitwa.*
1758.—*Fringilla Spinus* L., Syst. Nat., ed. X., p. 181
(Sweden).
Acanthis spinus (L.), Hartert.
Siskin.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Tsusnima, Corea, Riukiu.

Acanthis Bechstein (1803).

739. **Acanthis linaria holboellii** (Brehm). *Beni-hiwa*.
1831.—*Linaria Holboellii* Brehm, Handb. Naturg., Vog. Deutchl., p. 280 (Germany).
Acanthis flammea holboellii (Brehm), Hartert.
Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Corea.
740. **Acanthis hornemannii exilipes** (Coues). *Kō-beni-hiwa*.
1861.—*Aegithus exilipes* Coues, Proc. Acad. Nat. Sci. Phil., p. 385 (Fort Simpson, Arctic America).
Hab. Saghalin, Hondo (straggler).

Pyrrhula Pallas (1811).

741. **Pyrrhula pyrrhula kamtschatica** Taczanowski
..... *Benibara-uso*.
1882.—*Pyrrhula pyrrhula kamtschatica* Taczanowski, Bul. Soc. Zool. France, p. 395.
Kamtschatkan Bullfinch.
Hab. N. Kurile Isls., Hondo (Prov. Toyama—straggler).
742. **Pyrrhula pyrrhula kurilensis** Sharpe *Chishima-uso*.
1886.—*Pyrrhula kurilensis* Sharpe, Zoologist, p. 485 (Kurile Islands).
Hab. Kurile Islands.
743. **Pyrrhula phrrhula griseiventris** Lafresnaye *Uso*.
1841.—*Pyrrhula griseiventris* Lafresnaye, Rev. Zool., p. 241 (Japan).
1850.—*Pyrrhula orientalis* Temm. and Schl., Fauna Japon., Aves, p. 91, pl. 53 (Japan).

1882.—*Pyrrhula rosacea* Seebohm, Ibis, p. 371 (Yokohama).
Hab. Saghalin, Hokkaido, Hondo, Shikoku, Kiushiu, Tsushima,
Corea.

744. **Pyrrhula owstoni** Rosthschild and Hartert *Takrisa-ōuso*.

1907.—*Pyrrhula owstoni* Rosthschild and Hartert, Bull. B. O.
C., XXI, p. 9 (Formosa).

Hab. Mountains of Formosa.

745. ? **Pyrrhula nipalensis uchidai** Kuroda *Uchida-uso*.

1916.—*Pyrrhula uchidai* Kuroda, "Dōbutsugaku Zasshi"
(Tokio Zool. Mag.), XXVIII, p. 265 (Shishaban, Akō
Distr. in Formosa); id, Annal. Zool. Jap., IX, p. 295
(1917).

Kuroda's Bullfinch.

Hab. Mountains of Formosa.

746. **Pyrrhula erythaca arizanica** Ogilvie-Grant *Arisan-uso*.

1912.—*Pyrrhula arizanica* Ogilvie-Grant, Ibis, p. 644 (Mt.
Arizan, Formosa).

Hab. Mountains of Formosa.

Uragus Keys and Blasius (1840).

747. **Uragus sibiricus ussuriensis** Buturlin *Oniga-beni-mashiko*.

1915.—*Uragus sibiricus ussuriensis* Buturlin, Mess. Orn. p.
128 (Ussuriland).

Uragus sibirica sibirica (nec Pallas), Kuroda.

Hab. Corea.

748. **Uragus sibiricus sanguinolentus** (Temminck and Schlegel)
..... *Beni-mashiko*.

1850.—*Pyrrhula sanguinolenta* Temm. and Schl., Fauna
Japon., Aves, p. 92, pl. 54 (Japan).

Uragus sibirica sanguinolenta (Temm. and Schl.), Hartert.

Uragus sanguinolenta (Temm. and Schl.), Uchida.
Hab. Saghalin, Kurile Isls., Hokkaido, Hendo, Shikoku.

Carpodacus Kaup (1829).

749. **Carpodacus roseus** (Pallas) *Ō-mashiko*.
1776.—*Fringilla rosea* Pall., Reise d. versch, Prov. d. Russ. Reiches III, p. 699 (“.....in salicetis ad Udam et Selingam occurrit ”).
Carpodacus rosea (Pall.), Hartert.
Pallas's Rose-finches.
Hab. Saghalin, Hokkaido, Hondo, Kiusiu, Corea.
750. **Carpodacus vinaceus formosanus** Ogilvie-Grant
..... *Takasago-mashiko*.
1911.—*Carpodacus formosanus* Ogilvie-Grant, Bull. B. O. C., XXVII, p. 51 (Formosa).
Formosan Rose-finches.
Hab. Mountains of Formosa.
751. **Carpodacus erythrinus grebnitskii** Stejneger ... *Aka-mashiko*.
1885.—*Carpodacus erythrina grebnitskii* Stejneger, Orn. Expl. Command. Is. and Kamtschatka, p. 265 (Kamtschatka and ? Siberia).
Carpodacus erythrinus (nec Pall.), Uchida.
Kamtschatkan Scarlet-finches.
Hab. Saghalin.

Chaunoproctus Bonaparte (1850).

752. **Chaunoproctus ferreirostris** (Vigors) ... *Ogasawara-mashiko*.
1828.—*Coccothraustes ferreirostris* Vigors, Zool. Journ., p. 354 (Bonin Islands).

Hab. Bonin Islands (extinct species).

Pinicola Viciot (1807).

753. **Pinicola enucleator sakhalensis** Buturlin
..... *Karafuto-ginzan-mashiko.*

1915.—*Pinicola enucleator sakhalensis* Buturlin, Mess. Orn.,
pp. 129, 130 (Sakhalin).

Pinicola enucleator (nec L.), Uchida (pt.)

Sakhalin Island Pine-Grosbeak.

Hab. Sakhalin.

754. **Pinicola enucleator urupensis** Buturlin*Ginzan-mashiko.*

1915.—*Pinicola enucleator urupensis* Buturlin, Mess. Orn., p.
239 ("Urup," Kurile Isls.).

Pinicola enucleator (nec L.), Uchida (pt.).

Hab. Kurile Isls., Hokkaido.

Loxia L. (1758).

755. **Loxia curvirostra caucasica** Buturlin*Isuka.*

1907.—*Loxia curvirostra caucasica* Buturlin, Orn. Monatsber.,
p. 9 (Caucasus).

Loxia curvirostra curvirostra (nec L.), Hartert (pt.).

Eastern Crossbill.

Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Sulphur Is.,
Corea.

756. **Loxia curvirostra japonica** Ridway*Shirohara-isuka.*

1870.—*Loxia albiventris* Swinhoe, Proc. Zool. Soc. Lond., p.
437 (Peking). Preoccupied.

1885.—*Loxia curvirostra japonica* Ridway, Proc. Biol. Soc.
Washington, II, p. 101 (Japan).

Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Corea.

757. **Loxia leucoptera elegans** Homeyer *Naki-isuka.*
1879.—*Loxia elegans* Homeyer, Journ. f. Orn., p. 180 (Amur).
Loxia bifasciata (nec Brehm), Uchida.
Eastern Two-barred Crossbill.
Hab. Hondo (Awomori, Shinano).

Fringilla Brehm (1828).

758. **Montifringilla brunneinucha** (Brandt). *Hagi-mashiko.*
1842.—*Fringilla (Linaria) brunneinucha* Brandt, Bull. Sc. Acad. St. Petersb., X, p. 252 (Kurile Islands).
Japanese Ground-Linnet.
Hab. Kurile Isls., Hokkaido, Hondo.

Fringilla L. (1758).

759. **Fringilla montifringilla** L. *Aotori.*
1758.—*Fringilla Montifringilla* L., Syst. Nat. ed. X, p. 179
(Sweden).
1909.—*Fringilla montifringilla subcuncolata* Kleinschmidt,
Falco, V, p. 14 (Japan).
Brambling.
Hab. Saghalin, Hokkaido, Hondo, Bonin Isls., Corea, Kiusiu,
Riukiu Isls.

Passer Koch (1816).

760. ? **Passer montanus kaibatoi** Munsterhjelm. *Karafuto-suzume.*
1916.—*Passer montanus kaibatoi* Munsterhjelm, Nyt. Mag.
Natur v. Kristiania, p. 170 (Sakhalin).
Passer montana saturatus (nec Stejn), Hartert (part.).
Passer montanus (nec Linn.), Uchida.
Saghalin Tree-Sparrow.
Hab. Saghalin.

- 761.** ♀ *Passer montanus dybowskii* Domaniewski...*Chōsen-suzume*.
 1915.—*Passer montanus dybowskii* Domaniewski, Comp. rend.
 Soc. Sci. Varsovie, VIII, fasc. 7, pp. 562, 566 (Ussuri and
 Corea).
Passer montanus saturatus (nec Stejn.), Hartert (part.).
 Hab. Corea.
- 762.** *Passer montanus saturatus* Stejneger.....*Suzume*.
 1885.—*Passer montanus saturatus* Stejneger, Proc. U. S. Nat.
 Mus., VIII, p. 19 (Riukiu Islands).
 1910.—*Passer montanus orientalis* Clark, Proc. U. S. Nat.
 Mus., XXXVIII, p. 69 (Hakodate and Fusan, Corea).
Passer montana (nec Linn.), Uchida.
 Japanese Tree-Sparrow.
 Hab. Kurile Isls., Hokkaido, Hondo, Seven Isls., Shikoku, Kiuisiu,
 Tsushima, Quelpart Is., Tanegashima, Yakushima,
 Amami-Ōshima, Riukiu Isls.
- 763.** *Passer montanus taiwanensis* Hartert*Taiwan-suzume*.
 1904.—*Passer montana taiwanensis* Hartert, Vög. pal., I, p.
 161 ("Daihoku," Formosa).
Passer montana (nec Linn.), Uchida.
 Formosan Tree-Sparrow,
 Hab. Formosa, Percadores Isls.
- 764.** *Passer rutilans rutilans* (Temminck).....*Niunai-suzume*.
 1829.—*Fringilla rutilans* Temminck, Pl. Col., III, p. 488
 (Japan).
 1850.—*Passer russatus* Temm. and Schl., Faun. Japan., Aves,
 p. 90, pl. 50 (Japan).
Passer rutilans (Temm.), Uchida.
 Russet Sparrow.
 Hab. Sakhalin, Hokkaido, Hondo, Shikoku, Kiuisiu, Corea,
 Dagelet Is., Quelpart Is., Amami-Ōshima, Formosa.

Emberiza L. (1758).

765. **Emberiza leucocephalos** S. G. Gmelin *Shiraga-hōjirō.*
1771.—*Emberiza leucocephalos* S. G. Gmelin, Nov. Comm.
Acad. Sci. Imp. Petrop., XV, p. 480, pl. 23, fig. 3 (Astra-
chan).
Pine Bunting.
Hab. Saghalin, Hokkaido, Hondo (Shinano—straggler), Corea.
766. **Emberiza rutila** Pallas *Shima-nojiko.*
1776.—*Emberiza rutila* Pallas, Reise d. versch. Prov. d. Russ.
Reichs, III, p. 698 (Mongolia).
Chestnut Bunting, Ruddy Bunting.
Hab. Corea, Hokkaido (straggler), Hondo (Shinano—strag-
gler).
767. **Emberiza aureola** Pallas *Shima-awoji.*
1773.—*Emberiza aureola* Pallas, Reise d. versch. Prov. d.
Russ. Reichs, II, p. 711 (Siberia).
Yellow-breasted Bunting.
Hab. Saghalin, Kurile Isls., Hokkaido, Formosa.
768. **Emberiza elegans** Temminck *Miyama-hōjirō.*
1835.—*Emberiza elegans* Temminck, Pl. Col., 583 (Japan).
Yellow-throated Bunting.
Hab. Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, Corea,
Formosa.
769. **Emberiza spodocephala** spodocephala Pallas. ... *Kara-awoji.*
1776.—*Emberiza spodocephala* Pallas, Reise d. versch. Prov.
d. Russ. Reichs, III, p. 698 (Daurian Region).
Black-faced Bunting.
Hab. Hondo (Tokyo—straggler !), Corea, S. Riukiu (Ishigaki-
shima), Formosa.
770. **Emberiza spodocephala personata** Temminck *Awoji.*

1835.—*Emberiza personata* Temminck, Pl. Col., 580 (N. Japan.).

Japanese Bunting.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, Yakushima, Tanegashima, Amami-Ōshima, Riukiu (Okinawa group).

771. **Emberiza spodocephala**, subsp. ? *Kita-awaji*.

1891.—*Emberiza personata* (nec Temm.), Tacz., Faune Siberie orient. I, p. 567 ("Sacchalin").

1904.—*Emberiza spodocephala* subsp.?, Hartert, Vög. pal. I, p. 178 (Sacchalin).

Emberiza spodocephala personata (nec Temm.). Uchida (part.).

Hab. Saghalin.

772. **Emberiza sulphurata** Temminck and Schlegel *Nojiko*.

1848.—*Emberiza sulphurata* Temm. and Schl., Fauna Japon., Aves, p. 100, pl. 60 (Japan).

Japanese Yellow Bunting.

Hab. Hokkaido, Hondo, Kiusiu, Tsushima, Corea, Formosa.

773. **Emberiza cioides ciopsis** Bonaparte *Hōjirō*.

1850.—*Emberiza ciopsis* Bonaparte, Conspl. Av., I, p. 466 (Japan).

Emberiza cioides ciopsis Bp., Hartert. (pt.).

Japanese Meadow Bunting.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, ? Formosa.

774. **Emberiza cioides ijimae** Stejneger *Ijima-hōjirō*.

1894.—*Emberiza cioides ijime* Stejneger, Proc. U. S. Nat. Mus. XVI, p. 638 (Tsushima).

Emberiza cioides ciopsis (nec Bp.) Harter. (part.).

Emberiza cioides castaneiceps (nec Moore), Uchida.

- Hab. Dagelet Is., Quelpart Is., Tsushima, Iki, Kiusiu, Yaku-shima, Tanegashima, Seven Islands.
- 775. *Emberiza cioides castaneiceps* Moore** *Chōsen-hōjirō*.
 1855.—*Emberiza cioides costaniceps* Moore, Proc. Zool. Soc. Lond., p. 215 (China).
 Siberian Meadow Bunting.
 Hab. Corea.
- 776. *Emberiza fucata fucata* Pallas.** *Hō-aka*.
 1770.—*Emberiza fucata* Pallas, Reise d. versch. Prov. d. Russ. Reichs, III, p. 608 ("Ad Ononem et Ingodam in ripis").
 Grey-headed Bunting.
 Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Tanegashima, Amami-Ōshima, Corea, Quelpart Is., Formosa.
- 777. *Emberiza pusilla* Pallas** *Ko-hōaka*.
 1776.—*Emberiza pusilla* Pallas, Reise d. versch. Prov. d. Russ. Reichs, III, p. 697 (Daurian Region).
 Little Bunting.
 Hab. Corea, Quelpart Is., Hondo.
- 778. *Emberiza rustica* Pallas** *Kashiradata*.
 1776.—*Emberiza rustica* Pallas, Reise d. versch. Prov. d. Russ. Reichs, III, p. 698 ("In salicetis Davuriae").
 Rustic Bunting.
 Hab. Saghalin, Kurile Isls., Hokkaido, Hondo, Shikoku, Kiusiu, Tsushima, Corea.
- 779. *Emberiza tristrami* Swinhoe** *Shirohara-hōjirō*.
 1870.—*Emberiza tristrami* Swinhoe, Proc. Zool. Soc. Lond., p. 441 (Amoy).
 Tristram's Bunting.
 Hab. Corea.
- 780. *Emberiza chrysophrys* Pallas** *Kimayu-hōjirō*.

1776.—*Emberiza chrysophrys* Pallas, Reise d. versch. Prov. d. Russ. Reichs, III, p. 698 (Daurian Region).

Yellow-browed Bunting.

Hab. Corea (S. Zenra Distr.).

781. **Emberiza variabilis** Temminck. *Kuroji*.

1835.—*Emberiza variabilis* Temminck, Pl. Col., 583 (North Japan).

Japanese Grey-Bunting.

Hab. Kurile Isls., Hokkaido, Hondo, Shikoku, Kiushiu, Yakushima, Amami-Ōshima.

Cynchramus Boie (1826).

782. **Cynchramus yessoensis** yessoensis (Swinhoe) *Ko-jurin*.

1863.—*Emberiza minor* (nec Midd.), Blakiston, Ibis, p. 99 (“Hakodate, Yesso”).

1874.—*Schanicola yessoensis* Swinhoe, Ibis, p. 161 (Hakodate).
Emberiza yessoensis (Swinh.), Hartert.

Japanese Reed-Bunting.

Hab. Kurile Isls., Hokkaido, Hondo.

783. **Cynchramus yessoensis continentalis** (Witherby)
..... *Chōsen-ko-jurin*.

1913.—*Emberiza yessoensis continentalis* Witherby, Bull. B. O. C., XXXI, p. 74 (Nanking, China).

Chinese Reed-Bunting.

Hab. Corea.

784. **Cynchramus pallasi** Cabanis *Sibiria-jurin*.

1851.—*Cynchramus pallasi* Cabanis, Mus. Hein. I, p. 130 (Transbaikalia).

Emberiza pallasi (Cab.), Hartert.

Emberiza passerina (nec Pall.), Takatsukasa, Uchida Brd.

Jap., II.

Hab. Corea.

785. ♀ **Cynchramus schoeniclus pallidior** (Hartert). . *Chōsen-ō-jurin*.

1904.—*Emberiza schoeniclus pallidior* Hartert, Vög. pal., I, p. 197 (“Aiderli in Turkestan”).

Hab. Corea.

786. **Cynchramus schoeniclus pyrrhulinus** (Swinhoe) *ō-jurin*.

1876.—*Emberiza pyrrhulinus* Swinhoe, Ibis, p. 333, pl. VIII, fig. 2 (Hakodate, Japan).

Emberiza schoeniclus pyrrhulinus Swinh., Hartert.

Finch-billed Reed-Bunting.

Hab. Hokkaido, Hondo, Shikoku.

Calcarius Bechstein (1803).

787. **Calcarius lapponicus coloratus** Ridgway... *Tsumenaga-hōjiro*.

1898.—*Calcarius lapponicus coloratus* Ridgway, Auk, XV, p. 320 (“Copper Island, Kamtschatka”).

Calcarius lapponica (nec Linn.), Uchida.

Hab. Saghulin, Kurile Isls., Hokkaido, Hondo (Shinano and Suruga—straggler!), Corea.

Plectrophenax Stejneger (1882).

788. **Plectrophenax nivalis townsendi** Ridgway. ... *Yuki-hōjiro*.

1901.—*Plectrophenax nivalis townsendi* Ridgway, B. North and Middle America, I, p. 147 (Kommander Is., Pribilof. Is., etc.).

Passerina nivalis nivalis (nec L.), Harter. (part).

Passerina nivalis (nec Linn.), Uchida.

Hab. Saghulin, Kurile Isls., Hokkaido, Hondo (Uzen—straggler), Corea.

Alphabetical Index to Generic Names.

A

Abrornis	124
Acanthis	174
Acanthopneuste	131
Accipiter	42
Acrocephalus	137
Actitis	69
Ægithaliscus	159
Ægithalos	158
Ægithocichla	126
Aegolius	99
Ægyptius	35
Æthia	85
Æthiopsar	168
Aix	30
Alauda	111
Alcedo	94
Alcella	85
Alcippe	138
Alseonax	121
Alsocomus	87
Amaurornis	56
Anas	28
Anous	78
Anser	32
Anteliotringa	65
Anthropoides	59
Anthus	115
Apalopteron	119
Apus	102
Aquila	35
Arboricola	51

Archibuteo	39
Ardea	13
Ardeola	16
Arenaria	63
Areoturnix	46
Arquatella	65
Asio	98
Astur	41

B

Bambusicola	50
Bombycilla	148
Botaurus	18
Brachypteryx	131
Brachyramphus	83
Bradypterus	136
Branta	30
Bubo	95
Bubulcus	16
Bucephala	24
Buchanga	148
Budytes	113
Bulweria	7
Burnesia	137
Butastur	37
Buteo	39
Butorides	16

C

Calcarius	184
Calliope	130
Calobates	114
Canutus	64
Caprimulgus	102

<i>Chaptia</i>	148	<i>Crocethia</i>	65
<i>Carpodacus</i>	176	<i>Cuculus</i>	92
<i>Casarca</i>	29	<i>Cyanopica</i>	165
<i>Casmerodus</i>	14	<i>Cyanops</i>	103
<i>Catharacta</i>	81	<i>Cyanoptila</i>	123
<i>Centrops</i>	91	<i>Cyano-sylvia</i>	129
<i>Cephus</i>	83	<i>Cygnopsis</i>	33
<i>Cerchneis</i>	45	<i>Cygnus</i>	34
<i>Cerorhinca</i>	85	<i>Cynchramus</i>	183
<i>Certhia</i>	171	<i>Cyornis</i>	122
<i>Ceryle</i>	93						
<i>Chaimarrornis</i>	128						
<i>Chalcophaps</i>	90	D					
<i>Charadrius</i>	60	<i>Dafila</i>	25
<i>Chaulelasmus</i>	28	<i>Delichon</i>	145
<i>Chaunoproctus</i>	176	<i>Demigretta</i> (corr.)	<i>Demigretta</i>	
<i>Chen</i>	31			15
<i>Chionophilos</i>	113	<i>Dendrocitta</i>	166
<i>Chlidonia</i>	75	<i>Dendrocygna</i>	29
<i>Chloris</i>	173	<i>Dendronanthus</i>	115
<i>Ciceronia</i>	85	<i>Dicæum</i>	170
<i>Cichloselys</i>	124	<i>Diomedea</i>	8
<i>Ciconia</i>	19	<i>Ditelmatias</i>	72
<i>Cinclus</i>	143	<i>Dryobates</i>	105
<i>Circus</i>	40	<i>Dryocopus</i>	110
<i>Cirrepidesmus</i>	61	<i>Dupetor</i>	19
<i>Cisticola</i>	138	<i>Dytes</i>	2
<i>Clangula</i>	23						
<i>Coccothraustes</i>	171	E					
<i>Colœus</i>	163	<i>Egretta</i>	13
<i>Columba</i>	87	<i>Emberiza</i>	180
<i>Colymbus</i>	1	<i>Entomothera</i>	95
<i>Coprotheres</i>	81	<i>Eophona</i>	172
<i>Corvus</i>	161	<i>Erionetta</i>	22
<i>Cosmaerops</i>	95	<i>Erithacus</i>	129
<i>Coturnix</i>	50	<i>Erolia</i>	63
					<i>Erythropus</i>	44
					<i>Erythroscelus</i>	66

Eudromias	61	Halcyon	94						
Eudynamys	91	Haliæetus	37						
Eunetta	26	Hemichelidon	120						
Eupoda	61	Hemigarzetta	15						
Eurynorhynchus	66	Herbivocula	133						
Eurystomus	93	Heteropygia	64						
Excalfactoria	50	Heteroscelus	68						
F															
Falcipennis	51	Hierococcyx	93						
Falco	43	Hierofalco	44						
Fratercula	86	Hierophasis	50						
Fregata	12	Himantopus	70						
Fringilla	178	Hirundapus	103						
Fulica	57	Hirundo	144						
Fuligula	24	Histrionicus	22						
Fulmarus	8	Horornis	134						
G															
Galerida	112	Hydrophasianus	74						
Gallicrex	56	Hydroprogne	76						
Gallinago	72	Hypotænidia	53						
Gallinula	56	Hypothymis	122						
Garrulus	166	I										
Gelochelidon	75	Iauthia	128						
Glareola	59	ICoturus	130						
Glaucidium	100	IIornis	67						
Glottis	67	Ixbrychus	18						
Gorsachius	17	J										
Graphophasianus	48	Janthoenas	87						
Graucalus	146	Jynx	110						
Gygis	79	K										
Gygisterna	77	Ketupa	97						
Gypaetus	35	L										
H										Lagopus	52
Hæmatopus	74	Lalocitta	167						
Lanius		Lanius	149						

Larus	...	79
Larvivora	...	130
Leptophæthon	...	10
Leucoblepharon	...	31
Limicola	...	66
Limnobiaenus	...	55
Limosa	...	70
Lobipes	...	69
Locustella	...	135
Lophaithyia	...	3
Loxia	...	177
Lunda	...	86
Lymnocryptes	...	73

M

Machlolophus	...	154
Macropygia	...	89
Macrorhamphus	...	66
Mareca	...	26
Marila	...	24
Megalornis	...	57
Melanitta	...	23
Melanonyx	...	31
Melanopelargus	...	19
Melanosterna	...	77
Mergellus	...	22
Mergus	...	21
Merula	...	125
Mesoscolopax	...	72
Micranous	...	78
Microcichla	...	131
Microsarcops	...	62
Microscelis	...	117
Milvus	...	38
Monticola	...	127

Montifringilla	...	178
Motacilla	...	114
Munia	...	161
Muscicapula	...	121
Myiophonus	...	127

N

Nannocnus	...	18
Neocalaphasis	...	49
Neospilura	...	73
Nettion	...	27
Ninox	...	99
Nipponia	...	19
Notodera	...	131
Nucifraga	...	164
Numenius	...	71
Nyctea	...	96
Nycticorax	...	17
Nyroca	...	24

O

Oceanodroma	...	3
Oenopelia	...	89
Oidemia	...	23
Olor	...	34
Onychoprion	...	77
Oreocinchla	...	124
Oriolus	...	160
Ortygops	...	54
Osmotreron	...	91
Otis	...	59
Otodytes	...	2
Otus	...	97

P

<i>Pagoa</i> ...	61
<i>Pandion</i> ...	45
<i>Panurus</i> ...	160
<i>Parasula</i> ...	11
<i>Parus</i> ...	152
<i>Passer</i> ...	178
<i>Pedetaithya</i> ...	3
<i>Pelecanus</i> ...	12
<i>Pelidona</i> ...	63
<i>Pericrocotus</i> ...	147
<i>Periparus</i> ...	156
<i>Perisoreus</i> ...	167
<i>Pernis</i> ...	38
<i>Phaeopus</i> ...	71
<i>Phalacrocorax</i> ...	11
<i>Phalaropus</i> ...	69
<i>Phaleris</i> ...	84
<i>Phasianus</i> ...	47
<i>Philomachus</i> ...	65
<i>Phoenicurus</i> ...	128
<i>Phragamaticola</i> ...	136
<i>Pica</i> ...	165
<i>Picoides</i> ...	109
<i>Picus</i> ...	104
<i>Pinicola</i> ...	177
<i>Piscatrix</i> ...	10
<i>Pisobia</i> ...	63
<i>Pitta</i> ...	111
<i>Platalea</i> ...	20
<i>Plectrophenax</i> ...	184
<i>Pluvialis</i> ...	62
<i>Pnoepyga</i> ...	143
<i>Podiceps</i> ...	2
<i>Pœcile</i> ...	157

<i>Poliolimnas</i> ...	54
<i>Poliomyias</i> ...	123
<i>Polionetta</i> ...	28
<i>Polysticta</i> ...	22
<i>Porzana</i> ...	54
<i>Prinia</i> ...	137
<i>Procelsterna</i> ...	78
<i>Prunella</i> ...	141
<i>Pseudogeranus</i> ...	58
<i>Pseudoglottis</i> ...	67
<i>Pseudotadorna</i> ...	28
<i>Pterodroma</i> ...	7
<i>Ptychoramphus</i> ...	84
<i>Puffinus</i> ...	5
<i>Pycnonotus</i> ...	119
<i>Pyrrherodias</i> ...	13
<i>Pyrrhula</i> ...	174

Q

<i>Querquedula</i> ...	26
------------------------	----

R

<i>Rallina</i> ...	53
<i>Rallus</i> ...	53
<i>Recurvirostra</i> ...	70
<i>Reguloides</i> ...	133
<i>Regulus</i> ...	152
<i>Remiz</i> ...	159
<i>Rhopophilus</i> ...	138
<i>Rhyacophilus</i> ...	68
<i>Riparia</i> ...	146
<i>Rissa</i> ...	81
<i>Rostratula</i> ...	74

S

<i>Sapheopipo</i> ...	105
-----------------------	-----

Sarcogeranus	58
Saxicola	127
Sœophaëthon	9
Scolopax	72
Siphia	122
Sitta	150
Sittiparus	154
Spatula	25
Sphenurus	90
Spilornis	36
Spilura	73
Spinus	173
Spizaëtus	36
Spizixus	119
Spodiopsar	167
Squatarola	62
Stercorarius	82
Sterna	76
Sternula	77
Streptopelia	88
Sturnia	168
Subspilura	73
Sula	10
Surnia	100
Suthora	159
Sutoria	137
Suya	137
Synthliboramphus	84
Syrrhaptes	86
T					
Tadorna	29
Terekia	69
Terpsiphone	119
Tetrao	52
Tetrastes	51
Thalasseus	76
Thallasoaëtus	37
Tharrhaleus	142
Threskiornis	20
Thriponeax	110
Tringa	68
Troglodytes	142
Tryngites	65
Trypanocorax	163
Turnix	46
Tyto	102
U					
Upupa	96
Uragus	175
Uria	82
Urile	11
Urocissa	165
Uroloncha	161
Urosphena	135
V					
Vanellus	62
X					
Xema	79
Y					
Yungipicus	107
Z					
Zosterops	168
Zanthopygia	122

Alphabetical Index to Japanese Names.

A

<i>Abi.</i>	1
<i>Ahōdori.</i>	8	
<i>Ajisashi.</i>	76		
<i>Akaashi-chōzenbō.</i>	44				
<i>Akaashi-katsuodori.</i>	10				
<i>Akaashi-muzunagidori.</i>	6				
<i>Aka-ashi-shigi.</i>	67				
<i>Akadanna.</i>	111				
<i>Akacri-kaitsuburi.</i>	3				
<i>Akagashira-karasu-bato.</i>	88				
<i>Akagashira-sagi.</i>	16				
<i>Akagera.</i>	105				
<i>Akahajiro.</i>	25				
<i>Akahara.</i>	126				
<i>Akahara-daka.</i>	41				
<i>Akahara-gojūkara.</i>	152				
<i>Akahara-tsubame.</i>	144				
<i>Akahige.</i>	130				
<i>Akakokko</i>	126				
<i>Aka-mashiko.</i>	176				
<i>Aka-mozu.</i>	150				
<i>Akao-nettaichō.</i>	9				
<i>Akashōbin.</i>	95				
<i>Aka-tsukushigamo.</i>	29				
<i>Akawo-kakesu.</i>	167				
<i>Aka-yamadori.</i>	49				
<i>Akayerci-hircashi-shigi.</i>	69				
<i>Amami-hiyodori.</i>	118				
<i>Amami-kogera.</i>	108				
<i>Amami-yamagara.</i>	156				
<i>Amami-yama-shigi.</i>	72				

<i>Almanojakuma.</i>	110
<i>Amasagi.</i>	16
<i>Ame-tsubame.</i>	102
<i>Amerika-hidori.</i>	26
<i>Amerika-kogamo.</i>	27
<i>Amerika-umi-suzume.</i>	84
<i>Amūru-gojūkara.</i>	151
<i>Amūru-kogera.</i>	109
<i>Anadori.</i>	7
<i>Ancha-zuru.</i>	59
<i>Arisan-hitaki.</i>	128
<i>Arisan-misosazai.</i>	143
<i>Arisan-uso.</i>	175
<i>Arisui.</i>	110
<i>Asakura-sanshoku.</i>	146
<i>Atori.</i>	178
<i>Awo-ashi-shigi.</i>	67
<i>Azobato.</i>	90
<i>Azochimedori.</i>	140
<i>Azobazuku.</i>	99
<i>Azogera.</i>	104
<i>Awo-hanadori.</i>	171
<i>Awo-hauchiwadōri.</i>	137
<i>Awoji.</i>	180
<i>Awosagi.</i>	13
<i>Awo-shigi.</i>	73
<i>Awoshōbin.</i>	94
<i>Awotsura-katsuwodori.</i>	11

B

<i>Ban.</i>	56
<i>Banken.</i>	91
<i>Beni-ajisashi.</i>	76

<i>Benibara-uso.</i>	174
<i>Beni-bato.</i>	89
<i>Beni-hiwa.</i>	174
<i>Beni-mashiko.</i>	175
<i>Beni-sanshokni.</i>	147
<i>Binzui.</i>	115
<i>Birōdokinkuro.</i>	23
<i>Buppōsō.</i>	93

C

<i>Chabara-gojukara.</i>	151
<i>Chabara-ōruri.</i>	122
<i>Chigo-hayabusa.</i>	44
<i>Chigo-mozu.</i>	149
<i>Chimedori.</i>	140
<i>Chishima-bitaki.</i>	121
<i>Chishima-kasasagi.</i>	165
<i>Chishima-misosazai.</i>	142
<i>Chishima-raichō.</i>	52
<i>Chishima-shigi.</i>	65
<i>Chishima-ugarasu.</i>	11
<i>Chishima-umi-bato.</i>	83
<i>Chishima-uso.</i>	174
<i>Chiu-hibari.</i>	111

<i>Chōgenbō.</i>	45
<i>Chōsen-awobazuku.</i>	99
<i>Chōsen-chōgenbō.</i>	45
<i>Chōsenfukuro.</i>	100
<i>Chōsen-gejūkara.</i>	152
<i>Chōsen-hashibutogarasu.</i>	162
<i>Chōsen-hōjiro.</i>	182
<i>Chōsen-karvarahīwa.</i>	173
<i>Chōsen-kogara.</i>	158
<i>Chōsen-ko-jurin.</i>	183
<i>Chōsen-kumagera.</i>	110
<i>Chōsen-mejiro.</i>	170

<i>Chōsen-mifuuuzura.</i>	46
<i>Chōsen-misosazai.</i>	143
<i>Chōsen-ōakagera.</i>	105
<i>Chōsen-ō-jurin.</i>	184
<i>Chōsen-ōwashi.</i>	38
<i>Chōsen-shimafukurō.</i>	96
<i>Chōsen-shōdō-tsubame.</i>	146
<i>Chōsen-suzume.</i>	179
<i>Chōsen-tahibari.</i>	116
<i>Chōsen-uguisu.</i>	134
<i>Chōsen-yamagera.</i>	104
<i>Chōsen-yezoraichō.</i>	51
<i>Chūdai-awo-bato.</i>	90
<i>Chu-jishigi.</i>	73
<i>Chū-sagi.</i>	71
<i>Chu-shaku-sihgi.</i>	14

D

<i>Dai-sagi.</i>	14
<i>Dai-shaku-shigi.</i>	71	
<i>Daisen.</i>	62	
<i>Daruma-enaga.</i>	159	

E

<i>Enaga.</i>	158
<i>Eriguro-ajisashi.</i>	77	
<i>Erimaki-shigi.</i>	65	
<i>Essai.</i>	42
<i>Etopirika.</i>	86
<i>Etorofu-umi-suzume.</i>	85
<i>Ezo-hiyodori.</i>	117
<i>Ezo-kogara.</i>	157
<i>Ezo-mushikui.</i>	131

F

<i>Fukurō.</i>	102
----------------	-----	-----	-----	-----	-----

<i>Furnma-kamome.</i>	8	<i>Hariwo-ama-tsubame.</i>	...	103
				<i>Hashibiro-gamo.</i>	...	25
				<i>Hashiboso-shirochidori.</i>	...	60
				<i>Hashiboso-garasu.</i>	...	163
				<i>Hashiboso-mizunagidori.</i>	...	5
				<i>Hashibuto-ajisashi.</i>	...	75
				<i>Hashibutogarasu.</i>	...	162
				<i>Hashibuto-goi.</i>	...	17
				<i>Hashibuto-hishikui.</i>	...	32
				<i>Hashibuto-hiyodori.</i>	...	118
				<i>Hashibuto-ikaru.</i>	...	172
				<i>Hashibuto-kogara.</i>	...	158
				<i>Hashibuto-ōyoshikiri.</i>	...	136
				<i>Hashibuto-umi-garasu.</i>	...	83
				<i>Hashijiro-abi.</i>	...	1
				<i>Hashinaga-hoshigarasu.</i>	...	164
				<i>Hashinaga-kuina.</i>	...	53
				<i>Hauchiwadori.</i>	...	137
				<i>Hayabusa.</i>	...	43
				<i>Hensongara.</i>	...	157
				<i>Henson-kogara.</i>	...	157
				<i>Herasagi.</i>	...	20
				<i>Hera-shigi.</i>	...	66
				<i>Hibari.</i>	...	112
				<i>Hibari-shigi.</i>	...	64
				<i>Hidori-gamo.</i>	...	26
				<i>Higara.</i>	...	156
				<i>Higegara.</i>	...	160
				<i>Hige-washi.</i>	...	35
				<i>Higoromo.</i>	...	160
				<i>Hi-kuina.</i>	...	55
				<i>Hime-ama-tsubame.</i>	...	103
				<i>Hime-fukurō.</i>	...	100
				<i>Hime-hishikui.</i>	...	32
				<i>Hime-isohiyo.</i>	...	127
				<i>Hime-iwa-tsubame.</i>	...	145

<i>Hime-kuina.</i>	54	<i>Iso-shigi.</i>	69
<i>Hime-kuroumitsubame.</i>	...	4		<i>Isuka.</i>	177
<i>Hime-mejiro.</i>	170	<i>Iwahibari.</i>	141
<i>Hime-mifuuizura.</i>	46	<i>Iwami-sekirei.</i>	115
<i>Hime-ōchū.</i>	148	<i>Iwa-tsubame.</i>	145
<i>Hime-sasagoi.</i>	16	<i>Iwōjima-mejiro.</i>	169
<i>Hime-shiroharanizunagidori</i>	7							
<i>Hime-tobi.</i>	38					
<i>Hime-u.</i>	12	J				
<i>Hime-uzura.</i>	50	<i>Jiu-ichi.</i>	93
<i>Hi-renjaku.</i>	148	<i>Jōbitaki.</i>	128
<i>Hishikui.</i>	32	<i>Jusukake-bato.</i>	89
<i>Hiyodori.</i>	117					
<i>Hō-aka.</i>	182	K				
<i>Hoibii.</i>	140	<i>Kagoshima-awogera.</i>	104
<i>Hōjiro.</i>	181	<i>Kagoshima-shijukara.</i>	153
<i>Hōjiro-gamo.</i>	24	<i>Kaitsuburi.</i>	2
<i>Hōjiro-sekirei.</i>	114	<i>Kakesu.</i>	166
<i>Hokuroku-kiji.</i>	47	<i>Kakkō.</i>	92
<i>Hontō-akahige.</i>	130	<i>Kamabane-raichō.</i>	51
<i>Hōroku-shigi.</i>	71	<i>Kammuri-hibari.</i>	112
<i>Hoshigarasu.</i>	164	<i>Kammuri-kaitsuburi.</i>	3
<i>Hoshi-hajiro.</i>	24	<i>Kammuri-tsukushigamo.</i>	28
<i>Hototogisu.</i>	92	<i>Kammuri-umi-suzume.</i>	84
				<i>Kammuritōashi.</i>	36
				<i>Kamome.</i>	79
				<i>Kanoko-bato.</i>	89
<i>Ijima-hōjīro.</i>	181	<i>Kanokoshōbin.</i>	93
<i>Ijima-meboso.</i>	132	<i>Kansai-ōakagera.</i>	106
<i>Ijima-mejiro.</i>	169	<i>Kara-akahara.</i>	125
<i>Ijima-uguisu.</i>	134	<i>Kara-atwoji.</i>	180
<i>Ikaru.</i>	172	<i>Kara-chinedori.</i>	138
<i>Ikaru-chidori.</i>	60	<i>Kara-mukudori.</i>	168
<i>Inu-washi.</i>	35	<i>Karasckka.</i>	138
<i>Ishigakigara.</i>	153	<i>Kara-shirasagi.</i>	15
<i>Ishigaki-hiyodori.</i>	118	<i>Karafuto-awo-ashi-shigi.</i>	67
<i>Iso-hiyodori.</i>	127	<i>Karafuto-fukuro.</i>	101

Karafuto-ginzan-mashiko	177	Kibane-daruma-enaga.	160
Karafuto-misosazai	142	Kibara-shijukara.	154
Karafuto-mujisckka	133	Kibara-uguisu.	134
Karafuto-mushikui.	133	Kibashiri.	171
Karafuto-ōmosu.	149	Kibitaki.	122
Karafuto-raichō.	53	Kihibari.	115
Karafuto-shime	172	Kiji.	47
Karafuto-suzume.	178	Kiji-bato.	88
Karafuto-tahibari.	116	Kikuchi-chimedori.	131
Karafuto-washi	36	Kikuchi-hitaki.	128
Karafuto-yezoraichō.	51	Kikuchi-mejiro.	170
Karasu-bato.	87	Kikuitadaki.	152
Karen.	168	Kimawari.	151
Karigane.	32	Kimayu-hōjirō.	182
Karu-gamo.	28	Kimayu-mushikui.	138
Kasasagi.	165	Kin-bato.	90
Kashidori.	166	Kinkuro-hajiro.	24
Kashiradaka.	182	Kinme-fukurō.	99
Kasuidori.	102	Ki-renjaku.	148
Katashiro-washi.	35	Kiriai.	66
Katsuwodori.	10	Kisckirei.	114
Kawa-aisa.	21	Kitu-akagera.	105
Kawa-bitaki.	128	Kita-awoiji.	181
Kawachō.	93	Kita-higara.	156
Kawagarasu.	143	Kita-kibashiri.	171
Kawarabato.	87	Kitataki.	110
Kawari-sankō-chō.	120	Kita-tsumenaga-sekirei.	113
Kawasemi.	94	Kiusiu-fukurō.	101
Kaw-u.	11	Kiusiu-kiji.	48
Kayakuguri.	142	Kiusiu-kogera.	108
Kayanobori.	119	Ko-ahōdori.	9
Keashi-nosuri.	39	Ko-ajisashi.	77
Keima-furi.	83	Ko-akagera.	107
Keri.	62	Ko-awo-ashi-shigi.	67
Kewata-gamo.	22	Kobane-chimedori.	131
Kiashi-shigi.	68	Kobashi-binzui.	116

<i>Kobashi-chidori.</i>	61	<i>Kōrai-hikuina.</i>	55
<i>Kobashi-kihibari.</i>	116	<i>Kōrai-kiji.</i>	47
<i>Kobashi-umi-suzume.</i>	84	<i>Kōrai-uguisu</i>	160
<i>Ko-beni-hitwa.</i>	174	<i>Kōri-gamo.</i>	23
<i>Kobu-hakuchō.</i>	34	<i>Kōruri.</i>	130
<i>Ko-chidori.</i>	60	<i>Kosagi.</i>	13
<i>Ko-chōgyōnbō.</i>	44	<i>Kosame-bitaki.</i>	121
<i>Ko-gamo.</i>	27	<i>Ko-shaku-shigi.</i>	72
<i>Kogera.</i>	108	<i>Koshi-aka-tsubame.</i>	144
<i>Ko-hibari.</i>	112	<i>Ko-shigi.</i>	73
<i>Ko-hōōka.</i>	182	<i>Koshijiro-ajisashi.</i>	76
<i>Ko-ikaru</i>	172	<i>Koshijiro-hitaki.</i>	124
<i>Ko-jurin.</i>	183	<i>Koshijiro-kinpara.</i>	161
<i>Ko-kanome.</i>	80	<i>Koshijiro-umitsubame.</i>	3
<i>Ko-karigane.</i>	32	<i>Koshijiro-yamadori.</i>	49
<i>Ko-katwaaisa.</i>	21	<i>Kō-shiroharamizunagidori.</i>	7
<i>Ko-kawarahitwa.</i>	173	<i>Ko-umi-suzume.</i>	85
<i>Ko-kewatagamo</i>	22	<i>Koyoshikiri.</i>	137
<i>Koku-gan.</i>	30	<i>Kubitwa-kanome.</i>	79
<i>Kokumaru-garasu.</i>	163	<i>Kuina.</i>	53
<i>Komadori.</i>	129	<i>Kumagera.</i>	110
<i>Komakarigane.</i>	33	<i>Kumataka.</i>	36
<i>Ko-mamijiro-tahibari.</i>	117	<i>Kuro-ajisashi.</i>	78
<i>Koma-onaga.</i>	165	<i>Kuroashi-ahōdori.</i>	9
<i>Ko-mejirō.</i>	170	<i>Kurobitac-hariwo-ama-</i>				
<i>Komimizuku.</i>	98	<i>tsubame.</i>	103
<i>Komomojirō.</i>	14	<i>Kuro-gamo.</i>	23
<i>Komon-shigi.</i>	65	<i>Kurogashira.</i>	119
<i>Ko-mukudori.</i>	168	<i>Kuroharā-ajisashi.</i>	75
<i>Komushikui.</i>	132	<i>Kuro-hiyodori.</i>	119
<i>Kon-hitaki.</i>	131	<i>Kurohoshi-kogera.</i>	109
<i>Konohazuku.</i>	98	<i>Kuroji.</i>	183
<i>Konori.</i>	42	<i>Kuro-jō-bitaki.</i>	129
<i>Kōnotori.</i>	19	<i>Kuro-koshijiro umitsubame.</i>	3
<i>Ko-oba-shigi.</i>	64	<i>Kuro-miyakodori.</i>	74
<i>Kōrai-bato.</i>	87	<i>Kurosagi.</i>	15

<i>Kuro-sankōchō</i>	120	<i>Mikadokiji</i>	49				
<i>Kurotoki</i>	20	<i>Miko-aisa</i>	22				
<i>Kuro-tōzoku-kamome</i>	82	<i>Mimijiro-chimedori</i>	140				
<i>Kuro-tsugumi</i>	125	<i>Mimi-kaitsuburi</i>	2				
<i>Kurotsura-herasagi</i>	20	<i>Minami ko-chidori</i>	60				
<i>Kurotsuru</i>	57	<i>Misago</i>	45				
<i>Kuro-umitsubame</i>	4	<i>Misosazai</i>	142				
<i>Kuro-ycrihitaki</i>	122	<i>Mitsuyubi-kamome</i>	81				
<i>Kusa-shigi</i>	68	<i>Miyake-kogera</i>	108				
<i>Kyōjo-shigi</i>	63	<i>Miyakodori</i>	74				
M														
<i>Madara-chūhi</i>	40	<i>Miyama-garasu</i>	163				
<i>Madara-umi-suzume</i>	83	<i>Miyama-hitaki</i>	121				
<i>Magamo</i>	28	<i>Miyama-hōhjirō</i>	180				
<i>Magan</i>	33	<i>Miyama-kakesu</i>	166				
<i>Mahirwa</i>	173	<i>Miyamashōbin</i>	95				
<i>Makino-senniu</i>	136	<i>Miyama-tckkei</i>	51				
<i>Mamichajinai</i>	125	<i>Miyama-uguisu</i>	135				
<i>Mami-hauchitwadori</i>	137	<i>Miyubigera</i>	109				
<i>Mamijiro</i>	124	<i>Miyubi-shigi</i>	65				
<i>Mamijiro-ajisashi</i>	77	<i>Mizo-goi</i>	17				
<i>Mamijiro-kibitaki</i>	123	<i>Mizunazidori</i>	6				
<i>Mamijiro-kuina</i>	54	<i>Mozu</i>	150				
<i>Mamijiro-tahibari</i>	117	<i>Mugimaki</i>	123				
<i>Mamijiro-tsumenaga-sekirei</i>	113	<i>Mujisekka</i>	133				
<i>Manazuru</i>	58	<i>Mukudori</i>	167				
<i>Matsushima-kawaralahitwa</i>	173	<i>Munaguro</i>	62				
<i>Meboso</i>	132	<i>Muneaka-awo-bato</i>	91				
<i>Medai-chidori</i>	61	<i>Muneaka-hayabusa</i>	43				
<i>Meguro</i>	119	<i>Muneaka-hitaki</i>	121				
<i>Mejiro</i>	168	<i>Muneaka-tahibari</i>	116				
<i>Mejiro-chimedori</i>	138	<i>Murasaki-sagi</i>	13				
<i>Men-fukuro</i>	102	N									
<i>Meriken-kiashi-shigi</i>	68	<i>Nabc-kō</i>	19				
<i>Mifu-uzura</i>	46	<i>Nabe-zuru</i>	58				

<i>Naki-isuka</i>	178	<i>Ōhamu</i>	1
<i>Namiye-gera</i>	107	<i>Ō-hashi-shigi</i>	66
<i>Namiye-yamagura</i>	154	<i>Ō-hayabusa</i>	43
<i>Nanyō-mamijiro-ajisashi</i>	78	<i>Ō-hibari</i>	111
<i>Nezumi-gan</i>	30	<i>Ō-hishikui</i>	31
<i>Niitaka-kikuitadaki</i>	152	<i>Ojiro-hitaki</i>	122
<i>Niunai-suzume</i>	179	<i>Ojiro-tōnen</i>	64
<i>Nobitaki</i>	127	<i>Ojiro-washi</i>	37
<i>Nodoguro-tsugumi</i>	125	<i>Ō-jishigi</i>	72
<i>No-gan</i>	59	<i>Ō-jin-ichi</i>	93
<i>Nogoma</i>	130	<i>Ō-juriu</i>	184
<i>Noguchi-gera</i>	105	<i>Ō-kammuriteashi</i>	36
<i>Nojiko</i>	181	<i>Ō-kara-mozu</i>	149
<i>Nosuri</i>	39	<i>Ō-kawarahirwa</i>	173

O

<i>Ō-ajisashi</i>	76	<i>Okayoshi-gamo</i>	28
<i>Ō-akagura</i>	106	<i>Okinawagara</i>	153
<i>Ōban</i>	57	<i>Ō-konohazuku</i>	97
<i>Ōba-shigi</i>	65	<i>Ō-koshiakatsubane</i>	145
<i>Ō-chidori</i>	61	<i>Ō-magan</i>	33
<i>Ōchu</i>	148	<i>Ō-mashiko</i>	176
<i>Ō-fukurō</i>	101	<i>Ō-medai-chidori</i>	61
<i>Ogasawara-gabichō</i>	126	<i>Ō-mizunagidori</i>	5
<i>Ogasawara-hiyodoai</i>	118	<i>Ō-mozu</i>	149
<i>Ogasawara-karasu-bato</i>	88	<i>Onaga</i>	165
<i>Ogasawara-kawarahirwa</i>	173	<i>Onaga-bato</i>	89
<i>Ogasawara-mashiko</i>	176	<i>Onaga-beni-mashiko</i>	175
<i>Ogasawara-mizunagidori</i>	6	<i>Onaga-fukurō</i>	100
<i>Ogasawara-uguisu</i>	134	<i>Onaga-hanchiwadodi</i>	137
<i>Ogawa-komadori</i>	129	<i>Onaga-gamo</i>	25
<i>Ogawa-misosazai</i>	143	<i>Onaga-mizunagidori</i>	5
<i>Ō-gunkandori</i>	12	<i>Oni-jisashi</i>	76
<i>Oguro-shigi</i>	70	<i>Oni-kakkō</i>	91
<i>Oguro-zuru</i>	57	<i>Oni-sanshokui</i>	146
<i>Ō-hakuchō</i>	34	<i>Ō-nosuri</i>	39
		<i>Ō-raichō</i>	52
		<i>Oruri</i>	123

Ōsagarasu.	162
Ō-seguro-kamome.	79	
Ō-sekka.	136
Oshidori.	30
Ō-sorihashi-shigi.	70	
Ōsoton-gera.	107	
Ōsoton-umitsubame.	4	
Ōsoton-yamagara.	155	
Ō-taka.	41	
Ōtora-tsugumi.	124	
Ō-tōzoku-kamome.	81	
Ō-washi.	37	
Ō-yoshigoi.	18	
Ōyoshikiri.	137	
Ō-yuri-kamome.	81	

R

Kaichō.	52
Renkaku.	74	
Riukiu-akashōbin.	95	
Riukiu-awo-bato.	90	
Riukiu-gamo.	29	
Riukiu-hashibutogarasu.	...	162		
Riukiu-hikuina.	55	
Riukiu-hiyodori.	118	
Riukiu-karasu-bato.	88	
Riukiu-kibitaki.	123	
Riukiu-kiji-bato.	89	
Riukiu-kogira.	109	
Riukiu-konohazuku.	98	
Riukiu-mejiro.	169	
Riukiu-mozu.	150	
Riukiu-ōkonohazuku.	97	
Riukiu-ōkuina.	53	
Riukiu-sankō-chō.	120	
Riukiu-sanshokui.	147	

Riukiu-shijūkara.	153	
Riukiu-tsubame.	145	
Riukiu-yoshišoi.	18	
Ruri-bitaki.	128	
Ruri-chō.	127	
Ruri-kakesu.	167	

S

Saisiu-ōakagera.	107	
Saisiu-yamagara.	155	
Sakatsura-gan.	33	
Sakei.	86	
Same-bitaki.	120	
Sameiro-komimizuku.	99	
Sameiro-ō-konohazuku.	97	
Sankano-goi.	18	
Sankei.	50	
Sankō-chō.	119	
Sanshokui.	147	
Saruhama-shigi.	63	
Sasagoi.	16	
Sashiba.	37	
Seguro-ajisashi.	77	
Seguro-kakkō.	92	
Seguro-kamome.	79	
Seitaka-shigi.	70	
Sejirō-tahibari.	117	
Sekka.	138	
Sendai-mushikui.	132	
Shiberia-chūhi.	40	
Shiberia-fukurō.	101	
Shiberia-iwatsubame.	145	
Shiberia-jurin.	183	
Shiberia-senniu.	136	
Shichitō-mejiro.	169	
Shijūkara.	152	

<i>Shijukara-gan.</i>	31	<i>Shirohara-tōzoku-kamome.</i>	82
<i>Shikoku-yamadori.</i>	48	<i>Shiro-hayabusa.</i>	44
<i>Shimaaji.</i>	26	<i>Shiro-kamome.</i>	80
<i>Shima-awoiji.</i>	180	<i>Shirokuro-hitaki.</i>	131
<i>Shima-enaga.</i>	158	<i>Shirokurosagi.</i>	15
<i>Shima-fukuro.</i>	96	<i>Shōdō-tsubame.</i>	146
<i>Shimagonma.</i>	129	<i>Shōjōsagi.</i>	16
<i>Shimakiji.</i>	48	<i>Sōdeguro-zuru.</i>	58
<i>Shima-kinpara.</i>	161	<i>Sorihashi-scitaka-shigi.</i>	70
<i>Shima-kuina.</i>	54	<i>Sorihashi-shigi.</i>	69
<i>Shima-majiro.</i>	169	<i>Suinbō-gura.</i>	159
<i>Shima-mozu.</i>	150	<i>Sunamuguri-tsubame.</i>	146
<i>Shima-nōjiko.</i>	180	<i>Suzu-gamo.</i>	24
<i>Shima-ōakagera.</i>	106	<i>Suzume.</i>	179
<i>Shima-senniu.</i>	135	<i>Suzume-fukuro.</i>	100
<i>Shima-shijūkara.</i>	153			
<i>Shima-yamagara.</i>	155	T		
<i>Shime.</i>	171	<i>Ta-gvri.</i>	62
<i>Shinori-gamo.</i>	22	<i>Tahibari.</i>	116
<i>Shiosazai.</i>	135	<i>Taiwan-awobato.</i>	90
<i>Shiraga-hōjirō.</i>	180	<i>Taiwan-haku-sekirei.</i>	115
<i>Shirahige-umi-suzume.</i>	85	<i>Taiwan-hashibutogarasu.</i>	163
<i>Shirako-bato</i>	89	<i>Taiwan-hibari.</i>	112
<i>Shirao-nettaichō.</i>	10	<i>Taiwan-higara.</i>	157
<i>Shirao-jisashi.</i>	79	<i>Taiwan-hikuina.</i>	55
<i>Shirobitac-gojūkara.</i>	150	<i>Taiwan-hoshigarasu.</i>	164
<i>Shiro-chidori.</i>	60	<i>Taiwan-isohiyo.</i>	127
<i>Shirocri-ōhamu.</i>	1	<i>Taiwan-juzukake-bato.</i>	87
<i>Shiro-fukuro.</i>	96	<i>Taiwan-kaitsubari.</i>	2
<i>Shirogashira.</i>	119	<i>Taiwan-kasuidori.</i>	102
<i>Shirohara.</i>	126	<i>Taiwan-kawasemi.</i>	94
<i>Shirohara-gojūgara.</i>	150	<i>Taiwan-kiji.</i>	47
<i>Shirohara-hōjirō.</i>	182	<i>Taiwan-kinpara.</i>	161
<i>Shirohara-isuka.</i>	177	<i>Taiwan-kogera.</i>	109
<i>Shirohara-kuina.</i>	56	<i>Taiwan-ko-hibari.</i>	112
<i>Shiroharamizunagidori.</i>	7	<i>Taiwan-konohazuku.</i>	98

Taiwan-ko-uguisu.	135	Tane-awogera.	105
Taiwan-kumataka.	36	Tane-yamagara.	155
Taiwan-mifu-uzura.	46	Ta-shigi.	72
Taiwan-mizogvi.	17	Tekkei.	50
Taiwan-ōakagera.	107	Tobi.	38
Taiwan-ōkonohazuku.	97	Tōgarasu.	153
Taiwan-ōkuina.	54	Toki.	19
Taiwan-onagadori.	166	Tomoc-gamo.	27
Taiwan-ōsekka.	136	Tōnen.	63
Taiwan-ōtaka.	41	Torafuzuku.	98
Taiwan-sekka.	138	Tora-mozu.	149
Taiwan-shijūkara.	154	Tora-tsugumi.	124
Taiwan-suzume.	179	Torishima-ajisashi.	78
Taiwan-tsugumi.	126	Tōzuku-kamome.	81
Taiwan-tsumitaka.	42	Tsubakuro.	144
Taiwan-tsutsudori.	92	Tsubame.	144
Taiwan-uguisu.	134	Tsubame-chidori.	59
Taiwan-yamagara.	156	Tsugumi.	125
Taiwan-yotaka.	102	Tsukushi-gamo.	29
Taiwan-znaka-awo-bato.	91	Tsumenaga-hōjirō.	184
Takabu-shigi.	68	Tsumenaga-s. kirci.	113
Takasago-ban.	56	Tsumi.	42
Takasago-daka.	41	Tsunomedori.	86
Takasago-daruma-enagu.	160	Tsuru-kuina.	56
Takasago-fukuro.	100	Tsuru-shigi.	66
Takasago-hitaki.	128	Tsushima-higara.	156
Takasago-kakesu.	167	Tsushima-iso-hiyo.	127
Takasago-kurosagi.	19	Tsushima-kakesu.	166
Takasago-mashiko.	176	Tsushima-kogera.	108
Takasago-misosazai.	143	Tsushima-misosazai.	143
Takasago-mozu.	149	Tsushima-yamagara.	15
Takasago-uso.	175	Tsutsudori.	2
Takedori.	140			
Takegarasu.	164	U		
Tama-shigi.	74	Uchida-uso.	175
Tanchō.	57	Uchiyam-a-sennin.	135

<i>Ugarasu.</i>	12	<i>Yamabato.</i>	90						
<i>Uguisu.</i>	134	<i>Yamadori.</i>	48						
<i>Umi-aisa.</i>	21	<i>Yamagura.</i>	154						
<i>Umi-bato.</i>	83	<i>Yamagura.</i>	104						
<i>Umi-garasu.</i>	82	<i>Yamahibari.</i>	142						
<i>Umi-neko.</i>	80	<i>Yamamusume.</i>	165						
<i>Umi-ōmu.</i>	84	<i>Yamasen'i.</i>	93						
<i>Umi-suzume.</i>	84	<i>Yama-shigi.</i>	72						
<i>Umi-u.</i>	11	<i>Yamashōbin.</i>	95						
<i>Uso.</i>	174	<i>Yanagi-mushikui.</i>	131						
<i>Usuaka-yamadori.</i>	49	<i>Yatsugashira.</i>	96						
<i>Utō.</i>	85	<i>Yezo-akagera.</i>	105						
<i>Utsuryō-shijukara.</i>	153	<i>Yezo-arisui.</i>	111						
<i>Uwo-mimizuku.</i>	97	<i>Yezo-bitaki.</i>	121						
<i>Uzura.</i>	50	<i>Yezo-fukuro.</i>	101						
<i>Uzura-shigi.</i>	64	<i>Yezo-gojūkara.</i>	151						
W																	
<i>Wakijiro-daka.</i>	42	<i>Yezo-kogera.</i>	107						
<i>Wakisuzi-hayabusa.</i>	43	<i>Yezo-ōakagera.</i>	106						
<i>Washi-kamome.</i>	80	<i>Yezo-raichō.</i>	51						
<i>Washi-mimizuku.</i>	96	<i>Yezo-senniu.</i>	135						
<i>Watarigarasu.</i>	161	<i>Yoshi-gamo.</i>	26						
Y																	
<i>Yabusame.</i>	135	<i>Yoshi-goi.</i>	18						
<i>Yairochō.</i>	111	<i>Yotaka.</i>	102						
<i>Yakushima-akakokko.</i>	126	<i>Yuki-hōhjirō.</i>	184						
<i>Yakushima-kibitaki.</i>	122	<i>Yuri-kamome.</i>	80						
<i>Yakushima-yamagara.</i>	155	Z											
<i>Sh</i>						<i>Zuaka-enoga.</i>	159						
<i>Shi</i>						<i>Zuakagara.</i>	159						
<i>Shi</i>						<i>Zuguro-kamome.</i>	81						

Corrigenda and Addenda.

Page 1. Read for *Colybmi* *Colymbi*.

Page 4. Read for *Oceanodroma melanina*

Oceanodroma melania matsudairae Kuroda.

1922.—*Oceanodroma melania matsudairae* Kuroda, Ibis, p. 311. (Off the coast of Saganui Bay, Hondo).

The following species on page 7 is to be erased :—

24. *Pterodroma brevipes* (Peale).

Page 24. Read for *Nyroca ferina ferinoides*

Nyroca ferina ferina (L.).

1758.—*Anas ferina* L., Syst. Nat., ed. X, I, p. 126 (Sweden).

Page 36. Read for *Spizaetus nipalensis nipalensis*

Spizaetus nipalensis fokiensis W. Sclater.

1919.—*Spizaetus nipalensis fokiensis* [W. Sclater], Swann, Syn. List. Accip., p. 72 (nom. nud.); W. Sclater, Bull. B. O. C., XL, p. 37 (1919) (Fokien, S. China).

Page 36. Read for *Spilornis cheela* (Latham), subsp.

Spilornis cheela perplexus Swann.

1922.—*Spilornis cheela perplexus* Swann, Syn. List. Accip., p. 135 (Yayeyama, S. Loo Choo Is.).

Page 37. Read for *Haliæetus albicilla*

Haliæetus albicilla albicilla (L.).

Page 39. Read for *Archibuteo* Brehm (1828) (nom. nud.)

Triorchis Kaup (1829).

Page 39. Read for *Archibuteo lagopus pallidus*

Triorchis lagopus pallidus (Menzbier).

Page 41. Read for *Astur soloensis soloensis* (Latham)

Astur soloensis (Horsfield).

1821.—*Accipiter soloensis* Horsfield, Trans. Linn. Soc., XIII,

p. 137 (Java).

Chinise Goshawk.

[Page 42. *Al. cuculoides* a synonym].

Page 49. Read for *Neocalophasis* Tak. & Kur. (1922)

Cyanophasis Buturlin (1908).

Page 49. Read for *Neocalophasis mikado*

Cyanophasis mikado (Ogilvie-Grant).

Page 51. Read for ? *Tetrastes bonasia amurensis*

Tetrastes bonasia coreensis Kuroda & Mori.

1922.—*Tetrastes bonasia coreensis* Kuroda & Mori, Auk, p. 365 (Kōgen Distr., Corea).

Page 64. 236½. **Heteropygia maculata** (Vieillot).....
..... Amerika-usurashigi.

1819.—*Tringa maculata* Vieill., Nouv. Dict. d'Hist. Nat., nouv. éd., XXXIV, p. 465 (Antilles or southern U. S.).
Tirolia maculata (Vieill.), Hartert.

American Pectoral Sandpiper.

Hab. Japan (Hartert).

Page 65. Read for *Arquatella maritima quarta*

Arquatella maritima couesi Ridgway.

1880.—*Arquatella couesi* Ridgw., Bull. Nutt. Orn. Club, V, p. 160 (Aleutian Islands).

Page 72. Read for *Gallinago* Koch (1816)

Capella Frenzel (1801).

Page 72. Read for *Gallinago gallinago raddei*

Capella gallinago raddei (Buturlin).

Page 73. Read for *Lymnocryptes gallinula*

Lymnocryptes minimus (Brünnich).

1764.—*Scolopax Minima* Brünnich, Orn. Bor., p. 49.

Page 74. Read for *Hæmatopus niger bachmani*

Hæmatopus niger niger (Gmelin).

1787.—*Scolopax nigra* Gmelin, Syst. Nat. I, 2, p. 659
(Aleutian Is.).

Page 80. Read for *Larus leucopterus*

Larus glaucopterus Meyer.

1822.—*Larus glaucopterus* Meyer, in Meyer & Wolf, Taschenb.
d. deutsch. Vögeln., p. 197 “Meere der arktischen Zone,
z. B. in Island, zuweilen im Herbst an den Küsten der
Ost—und Nordsee”.

Page 92. Read for *Cuculus intermedius intermedius*

Cuculus poliocephalus poliocephalus Latham.

1790.—*Cuculus poliocephalus* Lath., Ind. Orn. I, p. 214 (India).

Page 102. Read for *Apus pacificus*

Apus pacificus pacificus.

Page 105. 415½. **Dryobates major brevirostris** (Reichenbach)
..... *Hashibuto-akagera*.

1854.—*Picus brevirostris* (Reichenb.), Handb. d. Spec.
Orn., Scansores, p. 365, PL. DCXXXIII, fig. 4212
(“Irtisch,” Siberia).
Hab. Corea (straggler!).

Page 105. 417½. **Dryobates major seoulensis** Kuroda & Mori.
..... *Chōsen-akagera*.

1922.—*Dryobates major seoulensis* Kuroda & Mori, Auk,
p. 364 (Corea).

Hab. Corea.

Page 108. 430 $\frac{1}{2}$. **Yungipicus kizuki shikokuensis** Kuroda.

1922.—*Yungipicus kizuki shikokuensis* Kuroda, Annot.

Zool. Japon., X, Art. 4, p. 115 (Prov. Iyo, Shikoku).

Hab. S. Hondo (Yamato), Shikoku (Iyo).

Page 128. Read for *Chaimarronis*

Chaimarrornis.

Page 138. Read for *Mijirochimedori*

Mijirochimedori.

Page 148. Read for *Captia*

Chaptia.

Page 172. Read for *Korafuto-shime*

Karafuto-shime.

日本鳥學會規則摘要

大正十一年十一月二十一日印刷

定價 上製五圓
並製三圓

第二條 本會事務所は東京帝國大學理學部動物學教室内に置く
第四條 本會は前條の目的を達する爲め評議會の決議を經て隨時種
々の事業をなす

一當分一年に二回雜誌「鳥」を出版すること

一臨時刊行物を刊行すること

一毎年春秋二回會合し鳥類に關する講演談話なし同時に鳥類に
關する圖書標本其他の展覽會を催す

一鳥學的探檢を舉行すること

第五條 本會員を分ちて甲種會員と乙種會員の二とす

一甲種會員は會費として一箇年金五圓を納むること

一乙種會員は會費として一箇年金二圓五十錢を納むること

第六號 甲種會員には雜誌「鳥」及臨時刊行物を配布す

乙種會員には雜誌「鳥」を配布す臨時刊行物は配布せず定價の三割
引を以て講讀することを得

不許複製

東京市日本橋區兜町二番地
編輯者 兼 下 憲
發行者 木

東京市日本橋區兜町二番地
印刷人 神谷 岩次郎

東京市日本橋區兜町二番地
印刷所 東京印刷株式會社

帝國大學理學部
動物學教室内

日本鳥學會

會頭 公爵 司信輔
幹事 内田清之助

評議員
理學博士 飯塚 啓
黑田長禮 子爾松平
飯塚 啓 平 賴孝


發賣所


東京日本橋區
十軒店町

裳華房
振替口座東京一〇七番


□ 錄目物行刊時臨會學鳥本日 □

本會機關雜誌「鳥」二年回發行一價金圓稅六錢









A Hand-list of the Japanese

D. F. Garrison
D. F. Garrison

AMNH LIBRARY



100071523